

CLASSIFICATION OF THREE-DIMENSIONAL EXCEPTIONAL LOG CANONICAL HYPERSURFACE SINGULARITIES I

S. A. KUDRYAVTSEV

ABSTRACT. In this paper the three-dimensional exceptional strictly log canonical hypersurface singularities are described and the detailed classification of three-dimensional exceptional canonical hypersurface singularities is given under the condition of well-formedness.

Introduction

One of the main problems in log Minimal Model Program is the study of appeared extremal contractions (singularities), i.e. contractions $f: (X, D_X) \rightarrow (X', D_{X'})$, where $-(K_X + D_X)$ is f -ample divisor and $\rho(X/X') = 1$. For the three-dimensional varieties this problem is completely open except the classification of Fano manifolds obtained by V. A. Iskovskikh at the end of 70th. In solving this problem one of the main difficulties is the absence of geometrical description of singularities. The first step in Fano manifold classification is to find a "good" divisor in the anticanonical linear system $|-K_X|$. The existence of a "good" divisor for extremal contraction allows to understand the contraction structure (example 0.2). Therefore the main problem discussed in this paper can be formulated as the follows:

Problem 0.1. To find a "good" element in the multiple anticanonical linear system for a variety, extremal contraction or singularity.

Example 0.2. 1. Consider a small extremal contraction of three-dimensional terminal variety. Then the existence of a divisor with Du Val singularities in the linear system $|-2K_X|$ implies the flip existence [7].

2. If an extremal contraction of three-dimensional terminal variety is a conic bundle then the existence of a divisor with Du Val singularities in the anticanonical linear system $|-K_X|$ allows at once to obtain the complete local classification [16].

The presence of a "good" element is closely connected with the other problems, for example with the automorphisms of $K3$ surfaces [24, 1.12] and with the gorenstein indexes of strictly log canonical singularities (see example 1.8).

Recently V. V. Shokurov gave the approach to solve the problem 0.1 in [23],[24]. It consists of two parts:

1. The first idea is an inductive transfer from n -dimensional varieties to $(n - 1)$ -dimensional ones.
2. The second idea is to separate the varieties to the exceptional and nonexceptional ones.

Shortly, the first step is the following. First of all one have to find some divisor S on X or on some blow-up of X such that pair $(S, \text{Diff}_S(D_X))$ is klt and $-(K_S + \text{Diff}_S(D_X))$ is ample. Then a "good" element of this pair can be extended to a "good" divisor of K_X on X .

In studying varieties, extremal contractions, singularities the exceptional phenomenon study importance follows from the following observation:

1. If a variety or extremal contraction or singularity is nonexceptional then the linear system $| -nK_X |$ must have a "good" member for small n . For example, we can take $n \in \{1, 2\}$ for the two-dimensional singularities [23, 5.2] and $n \in \{1, 2, 3, 4, 6\}$ for the three-dimensional singularities [24, 7.1].
2. The exceptional singularities are "bounded" and can be classified. For example, in this paper it will be checked that the three-dimensional exceptional hypersurface singularities have the finite number of types.

Generally speaking, the *regular* (nonexceptional) extremal contractions (singularities) can not be completely classified even in the dimension three (for example, in [14] it was shown that it is impossible to enumerate all normal form equations of the three-dimensional terminal hypersurface singularities, which are the very simple three-dimensional singularities). On the other hand, the existence of small index complement allows to separate the extremal contractions (singularities) to the families with common properties.

The exceptional extremal contractions (singularities) can be completely classified but they can have the large minimal complement index.

In [18] it was proved that the exceptionality remains valid after an inductive transfer and conversely, if $(S, \text{Diff}_S(D_X))$ is exceptional then $(X/Z, D_X)$ is exceptional too. Therefore it became possible to speak about *the inductive method of the algebraic variety classification*.

This paper is devoted to the application of the inductive method to the hypersurface singularities. In this paper the three-dimensional exceptional hypersurface singularities are classified. For this purpose, at first the corresponding quasihomogeneous parts are described, then the purely log terminal blow-ups are constructed. Next the log Del Pezzo surfaces obtained are investigated on the exceptionality. In particular, the minimal index of complements is found for every singularity.

Also the description of three-dimensional exceptional strictly log canonical hypersurface singularities is given.

Main theorem. *Let $(X, 0) \subset (\mathbb{C}^4, 0)$ be a three-dimensional exceptional canonical (respectively strictly log canonical) hypersurface singularity defined by a polynomial f . Then there exists a biholomorphic coordinate change $\psi: (\mathbb{C}^4, 0) \rightarrow (\mathbb{C}_{t,z,x,y}^4, 0)$ and unique primitive vector $\mathbf{p} \in N_{\mathbb{R}}$ such that just one of the following two possibilities holds:*

1. *The quasihomogeneous polynomial $\tilde{f}_{\mathbf{p}} = (f \circ \psi)_{\mathbf{p}}$ defines an exceptional canonical (respectively strictly log canonical and canonical outside $\mathbf{0}$) singularity $(X_{\mathbf{p}}, 0) \subset (\mathbb{C}_{t,z,x,y}^4, 0)$. In this case $\mathbf{p}$ -blow-up of $\mathbb{C}^4$ induces purely log terminal blow-ups $\varphi: (Y, E) \rightarrow (X, 0)$ and $\varphi_{\mathbf{p}}: (Y_{\mathbf{p}}, E_{\mathbf{p}}) \rightarrow (X_{\mathbf{p}}, 0)$, where $(E, \text{Diff}_E(0)) = (E_{\mathbf{p}}, \text{Diff}_{E_{\mathbf{p}}}(0))$. That is, these singularities have the same type and in particular the same complement index.*

The canonical singularities satisfying the condition of well-formedness — $\text{Diff}_{E/\mathbb{P}(\mathbf{p})}(0) = 0$ are classified in the theorems 3.24, 3.28 and in the tables of chapter §4. The polynomial $\tilde{f}_{\mathbf{p}}$; $(E, \text{Diff}_E(0))$; minimal complement index are written in the tables.

The strictly log canonical and canonical outside $\mathbf{0}$ quasihomogeneous singularities are always exceptional (in any dimension) by theorem 2.13. In the three-dimensional case $(E, \text{Diff}_E(0)) = (\tilde{f}_{\mathbf{p}} \subset \mathbb{P}(\mathbf{p}), 0)$ is K3 surface with Du Val singularities and $(X, 0)$ is 1-complementary.

2. *$\tilde{f}_{\mathbf{p}} = t^3 + g_2^2(z, x, y)$, where g_2 is an irreducible homogeneous polynomial of degree two. In this case the purely log terminal blow-ups are constructed in the theorems 3.3 and 3.5. Also it was obtained the similar classification depending on the type of jets $\tilde{f}_5$ and $\tilde{f}_6$.*

For the non-degenerate strictly log canonical singularities it was proved that they are exceptional if and only if they have purely elliptic type $(0, d-1)$, where d is a dimension of the singularity [4]. Also for every non-degenerate strictly log canonical singularity the quasihomogeneous log canonical part can be found [6, 3.5]. Therefore, in our terms it is exceptional if and only if its quasihomogeneous part defines

a canonical outside $\mathbf{0}$ singularity (see corollary 2.18). Using this fact it is not difficult to get the list of all three-dimensional exceptional strictly log canonical hypersurface singularities. The mentioned one is not given in this paper. For the isolated singularities such list is given in [26]. This list corresponds to 95 families of weighted $K3$ surfaces obtained by M.Reid and I.Fletcher in [21, 4.5], [2]. Indeed, the single exceptional divisor of purely log terminal blow-up is $K3$ surface.

This paper is organized in the following way. In chapter §1 the main definitions and preliminary results are collected. In chapter §2 the quasihomogeneous singularities are studied. The main theorem is proved in chapter §3. Also the explanation of three-dimensional canonical singularities study on the exceptionality is given. In chapter §4 the summarizing tables are written.

I am grateful to Professor V.A.Iskovskikh, Professor Yu.G.Prokhorov and Professor V.V.Shokurov for useful discussions and valuable remarks.

The research was partially supported by a grant 99-01-01132 from the Russian Foundation of Basic Research and a grant INTAS-OPEN 2000#269.

1. Preliminary facts and results

All varieties are algebraic and are assumed to be defined over $\mathbb{C}$, the complex number field. We use the terminology, notations of log Minimal Model Program and the main properties of complements given in [8], [9], [20]. A *strictly lc singularity* is called lc singularity, but not klt singularity.

Definition 1.1. We say that the boundary $D = \sum d_i D_i$ has *standard coefficients* if $d_i \in \{1\} \cup \{1 - \frac{1}{m} | m \in \mathbb{N}\}$ for all i .

Definition 1.2. Let (X, D) be a log pair, where D is a subboundary. Then a $\mathbb{Q}$ -*complement* of $K_X + D$ is a log divisor $K_X + D'$ such that $D' \geq D$, $K_X + D'$ is lc and $n(K_X + D') \sim 0$ for some $n \in \mathbb{N}$.

Definition 1.3. Let X be a normal variety and let $D = S + B$ be a subboundary on X such that B and S have no common components, S is an effective integral divisor and $\perp B \perp \leq 0$. Then we say that $K_X + D$ is *n-complementary* if there is a $\mathbb{Q}$ -divisor D^+ such that

1. $n(K_X + D^+) \sim 0$ (in particular, nD^+ is an integral divisor);
2. $K_X + D^+$ is lc;
3. $nD^+ \geq nS + \perp(n+1)B \perp$.

In this situation the n -complement of $K_X + D$ is $K_X + D^+$. The divisor D^+ is called n -complement too.

The invariance of the complements with respect to log MMP and their inductive properties are shown in the following three theorems.

Theorem 1.4. [23, 5.4] *Let $f: X \rightarrow Y$ be a birational contraction and let D be a subboundary on X . Assume that $K_X + D$ is n -complementary for some $n \in \mathbb{N}$. Then $K_Y + f(D)$ is also n -complementary.*

Under additional assumptions we have the inverse statement:

Theorem 1.5. [24, 2.13] *Let $f: Y \rightarrow X$ be a birational contraction and let D be a subboundary on Y such that $K_Y + D$ is nef over X and $f_*(D)$ has standard coefficients. Assume that $K_X + f(D)$ is n -complementary. Then $K_Y + D$ is also n -complementary.*

Theorem 1.6. [17, 2.1] *Let $(X/Z \ni P, D = S + B)$ be a log variety. Put $S = \lfloor D \rfloor$ and $B = \{D\}$. Assume that*

1. $K_X + D$ is plt;
2. $-(K_X + D)$ is nef and big over Z ;
3. $S \neq 0$ near $f^{-1}(P)$;
4. D has standard coefficients.

Further, assume that near $f^{-1}(P) \cap S$ there exists an n -complement $K_S + \text{Diff}_S(B)^+$ of $K_S + \text{Diff}_S(B)$. Then near $f^{-1}(P)$ there exists an n -complement $K_X + S + B^+$ of $K_X + S + B$ such that $\text{Diff}_S(B)^+ = \text{Diff}_S(B^+)$.

Definition 1.7. Let $(X/Z \ni P, D)$ be a contraction of varieties, where D is a boundary.

1. Assume that Z is not a point (local case). Then $(X/Z \ni P, D)$ is said to be *exceptional* over P if for any $\mathbb{Q}$ -complement $K_X + D'$ of $K_X + D$ near the fiber over P there exists at most one (not necessarily exceptional) divisor E such that $a(E, D') = -1$.
2. Assume that Z is a point (global case). Then (X, D) is said to be *exceptional* if every $\mathbb{Q}$ -complement of $K_X + D$ is klt.

If $Z = X$ then $(X/X \ni P)$ is a singularity $(X \ni P)$.

Example 1.8. (1) Let $(X \ni P)$ be a singularity. Suppose that there is an effective divisor H such that (X, H) is lc and $\lfloor H \rfloor \neq 0$. Then the singularity is not exceptional. Therefore the three dimensional terminal singularity is not exceptional because there is a divisor having only Du Val singularities in the anticanonical linear system $|-K_X|$ [22, 6.4].

It should be noted that there are the examples of four-dimensional exceptional terminal singularities [11].

(2) Let $(X \ni P)$ be a strictly log canonical exceptional singularity. Then the minimal index of complements is equal to the gorenstein index of $(X \ni P)$ [12, 1.10]. The minimal index of complements is bounded for the three dimensional log canonical singularities [23, 7.1]. A hypothesis is that this index is not more then 66. For the strictly log canonical exceptional singularities it was proved in [5] and [3]. For the nonexceptional nonisolated strictly log canonical singularities the gorenstein index is not bounded [3, 5.1].

Definition 1.9. Let $(X \ni P)$ be a log canonical singularity and let $f: Y \rightarrow X$ be a blow-up such that the exceptional locus of f contains only one irreducible divisor E ($\text{Exc}(f) = E$). Then $f: (Y, E) \rightarrow X$ is called a *purely log terminal (plt) blow-up* if $K_Y + E$ is plt and $-E$ is f -ample.

Definition 1.10. Let $(X \ni P)$ be a log canonical singularity and let $f: Y \rightarrow X$ be a blow-up such that the exceptional locus of f contains only divisors E ($\text{Exc}(f) = E$). Then $f: (Y, E) \rightarrow X$ is called a *log canonical (lc) blow-up* if $K_Y + E$ is lc and $-E$ is f -ample.

Remark 1.11. In the definitions 1.9 and 1.10 it is demanded that divisor E must be $\mathbb{Q}$ -Cartier. Hence Y is a $\mathbb{Q}$ -gorenstein variety.

Remark 1.12. The notations are the same as in the definition 1.9. We have the following properties of plt blow-ups:

1. If X is klt then $-(K_Y + E)$ is f -ample and a plt blow-up always exists [12, 1.5].
2. If X is strictly lc then $a(E, 0) = -1$ and a plt blow-up always exists for the exceptional singularities [12, 1.9].
3. If X is $\mathbb{Q}$ -factorial then Y is also $\mathbb{Q}$ -factorial and $\rho(Y/X) = 1$ [16, 2.2]. Hence, in the definition 1.9 for a $\mathbb{Q}$ -factorial singularity it is not necessary to demand that the divisor $-E$ is f -ample because ampleness takes place always.
4. By the inversion of adjunction $K_E + \text{Diff}_E(0)$ is klt. If $K_E + \text{Diff}_E(0)$ is n -complementary then $K_Y + E$ is n -complementary and K_X is n -complementary too (see theorem 1.6 and theorem 1.4).
5. Let $f_i: (Y_i, E_i) \rightarrow (X \ni P)$ be two plt blow-ups. If E_1 and E_2 define the same discrete valuation of function field $\mathcal{K}(X)$ then f_1 and f_2 are isomorphic [16, 2.2].

Definition 1.13. Let $(X \ni P)$ be a log canonical singularity. It is said to be *weakly exceptional* if there exists only one plt blow-up (up to isomorphism).

The following two theorems 1.14, 1.15 demonstrate the inductive transfer from an n -dimensional exceptional (resp. weakly exceptional) klt singularity to the unique $(n - 1)$ -dimensional exceptional (resp. weakly exceptional) log Fano variety $(E, \text{Diff}_E(0))$ having the same indexes of complements. The inverse statements take place too.

Theorem 1.14. [18, 4.9] *Let $(X \ni P)$ be a klt singularity and let $f: (Y, E) \rightarrow X$ be a plt blow-up of P . Then the following conditions are equivalent:*

1. $(X \ni P)$ is an exceptional singularity;
2. $(E, \text{Diff}_E(0))$ is an exceptional log variety.

Theorem 1.15. [12, 2.1] *Let $(X \ni P)$ be a klt singularity and let $f: (Y, E) \rightarrow X$ be a plt blow-up of P . Then the following conditions are equivalent:*

1. $(X \ni P)$ is not a weakly exceptional singularity;
2. there exists an effective $\mathbb{Q}$ -divisor $D \geq \text{Diff}_E(0)$ such that $-(K_E + D)$ is ample and (E, D) is not klt;
3. there exists an effective $\mathbb{Q}$ -divisor $D \geq \text{Diff}_E(0)$ such that $-(K_E + D)$ is ample and (E, D) is not lc.

According to the example of weakly exceptional singularity but not exceptional singularity in any dimension [12, 2.4] we can construct the example of the exceptional singularity.

Example 1.16. Let $(X \ni P)$ be an n -dimensional canonical hypersurface singularity given by the equation $(x_1^n + x_2^{n+1} + x_3^{n+1} + \dots + x_{n+1}^{n+1} = 0) \subset (\mathbb{C}^{n+1}, 0)$. The weighted blow-up of $\mathbb{C}^{n+1}$ with weights $(n + 1, n, n, \dots, n)$ induces a plt blow-up of P . Using the basic theorems about the hypersurfaces in the weighted spaces [2] (see also §4), it is easy to calculate that the obtained log Fano variety $(E, \text{Diff}_E(0))$ is

$$\begin{aligned} (\tilde{x}_1 + \tilde{x}_2^{n+1} + \dots + \tilde{x}_{n+1}^{n+1} \subset \mathbb{P}(n + 1, 1, \dots, 1), \frac{n-1}{n}\{\tilde{x}_1 = 0\}) = \\ = (\mathbb{P}^{n-1}, \frac{n-1}{n}Q_{n+1}), \end{aligned}$$

where Q_{n+1} is a smooth hypersurface of degree $n + 1$ in $\mathbb{P}^{n-1}$. Applying the approach of [11] it is not difficult to check that given log variety is exceptional for $n \geq 3$. The divisor $\frac{n-1}{n}\{\tilde{x}_1 = 0\} + \frac{1}{n}\{\tilde{x}_2 = 0\}$

is n -complement of minimal index. Our singularity is exceptional by exceptionality criterion 1.14 and n -complementary by remark 1.12.

Proposition 1.17. *An exceptional singularity is weakly exceptional. Assume that there exists lc blow-up but not plt blow-up of $(X \ni P)$. Then $(X \ni P)$ is not exceptional.*

Proof. In general case the first statement was proved in [12, 1.12]. The second statement follows from the proof [6, 2.4]. In fact, let $f: (Y, E) \rightarrow (X \ni P)$ be lc blow-up but not plt blow-up. We may assume that $(X \ni P)$ is klt otherwise the proof is trivial. Since $-(K_Y + E)$ is f -ample then the linear system $|-n(K_Y + E)|$ is very ample over X for $n \gg 0$. Let $H \in |-n(K_Y + E)|$ be a general element and let $B = \frac{1}{n}f(H)$. By Bertini's theorem $K_Y + E + \frac{1}{n}H$ is lc but not plt. Since $K_Y + E + \frac{1}{n}H \sim_{\mathbb{Q}} 0$ then the pair (X, B) is lc but not exceptional. $\square$

The next proposition follows from the corollary [12, 1.6] and the theorem [12, 1.9].

Proposition 1.18. *Let $f: (Y, E) \rightarrow (X \ni P)$ be a plt blow-up of log canonical singularity and let $\dim f(E) \geq 1$. If $(X \ni P)$ is klt then it is not weakly exceptional. If $(X \ni P)$ is strictly lc then it is weakly exceptional but not exceptional.*

Example 1.19. [23, 5.2] A two dimensional klt singularity is exceptional if and only if it has type $\mathbb{E}_6, \mathbb{E}_7, \mathbb{E}_8$. Its plt blow-up is the extraction of central vertex of minimal resolution graph. The corresponding one-dimensional log Fano variety $(E, \text{Diff}_E(0))$ is $(\mathbb{P}^1, \frac{n_1-1}{n_1}P_1 + \frac{n_2-1}{n_2}P_2 + \frac{n_3-1}{n_3}P_3)$, where $(n_1, n_2, n_3) = (2, 3, 3), (2, 3, 4)$ or $(2, 3, 5)$ respectively.

Example 1.20. [20, §6] A two dimensional strictly lc singularity is exceptional if and only if it has one of the following types: simple elliptic singularity or singularities of $\tilde{\mathbb{D}}_4, \tilde{\mathbb{E}}_6, \tilde{\mathbb{E}}_7, \mathbb{E}_8$ types. Its plt blow-up is the extraction of central vertex of minimal resolution graph. The corresponding one-dimensional log Fano variety $(E, \text{Diff}_E(0))$ is a smooth elliptic curve, $(\mathbb{P}^1, \frac{1}{2}P_1 + \frac{1}{2}P_2 + \frac{1}{2}P_3 + \frac{1}{2}P_4)$ or $(\mathbb{P}^1, \frac{n_1-1}{n_1}P_1 + \frac{n_2-1}{n_2}P_2 + \frac{n_3-1}{n_3}P_3)$, where $(n_1, n_2, n_3) = (3, 3, 3), (2, 4, 4)$ or $(2, 3, 6)$ respectively.

Thus, there are only three types of induced log Fano varieties for the two-dimensional klt singularities and there are only five types for the two-dimensional strictly lc singularities. There is a conjecture that the

similar result about the finite number of types is true in any dimension. For the three-dimensional hypersurface singularities this conjecture is checked in this paper. The essential difference from the two-dimensional singularities is that the infinite number of nonisomorphic singularities exists in almost every type of them. But they have the same resolutions.

1.21. The main facts of toric geometry. We make use of toric geometry terminology (see [15]). Let N be a free abelian group $\mathbb{Z}^n$ and M is its dual $\text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$. Denote $N \otimes_{\mathbb{Z}} \mathbb{R}$ and $M \otimes_{\mathbb{Z}} \mathbb{R}$ by $N_{\mathbb{R}}$ and $M_{\mathbb{R}}$ respectively. We have a canonical pairing $\langle \cdot \rangle: N_{\mathbb{R}} \times M_{\mathbb{R}} \rightarrow \mathbb{R}$. Let σ be the positive quadrant $\mathbb{R}_{\geq 0}^n$ of $N_{\mathbb{R}} = \mathbb{R}^n$ and $\sigma^{\vee}$ its dual. Then $\mathbb{C}^n$ is the toric variety corresponding to the cone σ . For a fan Δ in N the corresponding toric variety is denoted by $T_N(\Delta)$. For a primitive element $\mathbf{p} \in N$ of a 1-dimensional cone $\tau = \mathbb{R}_{\geq 0}\mathbf{p}$ in Δ , the closure $\overline{\text{orb}(\mathbb{R}_{\geq 0}\mathbf{p})}$ is denoted by $D_{\mathbf{p}}$, where $D_{\mathbf{p}}$ is a divisor on $T_N(\Delta)$.

Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a hypersurface singularity and $\mathbf{p} = (p_1, \dots, p_n)$ a primitive element in N with $p_{i_1} > 0$ and $p_{i_2} > 0$ for some $i_1 \neq i_2$ (such $\mathbf{p}$ is called a *weight*). Let $\varphi: \mathbb{C}^n(\mathbf{p}) \rightarrow \mathbb{C}^n$ be the blow-up with a weight $\mathbf{p}$. Denote the proper transform of X on $\mathbb{C}^n(\mathbf{p})$ by $X(\mathbf{p})$. The blow-up $\varphi: \mathbb{C}^n(\mathbf{p}) \rightarrow \mathbb{C}^n$ and its restriction $\varphi|_{X(\mathbf{p})}: X(\mathbf{p}) \rightarrow X$ are called the *$\mathbf{p}$ -blow-ups*. The exceptional locus of $\varphi|_{X(\mathbf{p})}$ is denoted by $E_{\mathbf{p}}$. The exceptional locus $D_{\mathbf{p}}$ of φ is the weighted projective space $\mathbb{P}(\mathbf{p})$. These $\mathbf{p}$ -blow-ups are obtained by a subdivision of the cone σ . The corresponding fan consists of the faces of cones σ_i ($i = 1, \dots, n$), where σ_i is generated by $\mathbf{e}_1, \dots, \mathbf{e}_{i-1}, \mathbf{p}, \mathbf{e}_{i+1}, \dots, \mathbf{e}_n$. Here $\mathbf{e}_1, \dots, \mathbf{e}_n$ are the unit vectors $(1, 0, \dots, 0), \dots, (0, \dots, 0, 1)$ which generate σ .

A monomial $x_1^{m_1} \cdots x_n^{m_n} \in \mathbb{C}[[x_1, x_2, \dots, x_n]]$ is denoted by x^m , where $m = (m_1, \dots, m_n) \in \mathbb{Z}^n = M$. For a power series $f = \sum_m a_m x^m \in \mathbb{C}[[x_1, x_2, \dots, x_n]]$ we write $x^m \in f$, if $a_m \neq 0$. For $\mathbf{p} \in N_{\mathbb{R}}$ and a power series f we define

$$\mathbf{p}(f) = \min_{x^m \in f} \langle \mathbf{p}, m \rangle.$$

We denote the *leading term* $\sum_{\langle \mathbf{p}, m \rangle = \mathbf{p}(f)} a_m x^m$ of f with respect to $\mathbf{p}$ by $f_{\mathbf{p}}$.

Definition 1.22. For a power series $f = \sum_m a_m x^m \in \mathbb{C}[[x_1, x_2, \dots, x_n]]$ define *Newton polyhedron* $\Gamma_+(f)$ in $M_{\mathbb{R}}$ as follows:

$$\Gamma_+(f) = \text{the convex hull of } \bigcup_{x^m \in f} (m + \sigma^{\vee}).$$

The set of the interior points of $\Gamma_+(f)$ is denoted by $\Gamma_+(f)^0$. For each face γ of $\Gamma_+(f)$ we define the polynomial f_γ as follows:

$$f_\gamma = \sum_{m \in \gamma} a_m x^m.$$

A power series f is said to be *non-degenerate* if for every face γ the equation $f_\gamma = 0$ defines a smooth hypersurface in the complement of the hypersurface $x_1 \cdots x_n = 0$.

Theorem 1.23. [14] *Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a normal hypersurface singularity defined by a power series f and $\Gamma_+(f)$ is its Newton polyhedron. Let $(X, 0)$ be canonical (resp. log canonical). Then $\mathbf{1} = (1, 1, \dots, 1) \in \Gamma_+(f)^0$ (resp. $\mathbf{1} = (1, 1, \dots, 1) \in \Gamma_+(f)$). The inverse statements take place for the non-degenerate singularities.*

Now as a preliminary we discuss the *nonisolated hypersurface singularities study*.

Definition 1.24. Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a nonisolated hypersurface singularity (maybe nonnormal) defined by a polynomial f . By proposition 1.18 we can investigate this singularity on the exceptionality in the arbitrary small neighborhood $\mathcal{U}$ of the origin. For example, in the three-dimensional case it means that $(\text{Sing } X \setminus \mathbf{0}) \cap \mathcal{U} \cong \bigsqcup_i \mathbb{C}_i^1$ is not arcwise connected union. The variety $X \cap \mathcal{U}$ is isomorphic to $\mathbb{C}^1 \times (\text{two dimensional log canonical hypersurface singularity})$ along the every component. In an n -dimensional case let $(\text{Sing } X \setminus \mathbf{0}) \cap \mathcal{U} \cong \bigcup_{i=1}^m (\mathcal{A}_i^{n_i} \setminus \mathbf{0})$, where $\mathcal{A}_i^{n_i}$ are the irreducible affine varieties of the dimension n_i . Now we define the mutually different affine varieties $\{V_j^{n_j}\}$, where $n_j = \dim V_j^{n_j}$. The system of indexes $I_j \subset \{1, \dots, m\}$ is called *admissible* if

$$\emptyset \neq \bigcap_{i \in I_j} (\mathcal{A}_i^{n_i} \setminus \mathbf{0}) \not\subseteq \mathcal{A}_k^{n_k}, \text{ where } k \notin I_j.$$

For all admissible systems I_j we can put $V_j^{n_j} \stackrel{\text{def}}{=} \bigcap_{i \in I_j} \mathcal{A}_i^{n_i}$. Let $\mathfrak{a}_j$ be the ideal radical of the variety $V_j^{n_j}$ and let $f_1, \dots, f_r$ be the generators of $\mathfrak{a}_j$. Then we have the decomposition

$$f = \Phi_1(f_1, \dots, f_r)u_1(x_1, \dots, x_n) + \dots + \Phi_m(f_1, \dots, f_r)u_m(x_1, \dots, x_n),$$

where $u_i \notin \mathfrak{a}_j$ for all i . The variety $\{u_1 = \dots = u_m = 0\}$ is empty or nonsingular by the construction of $V_j^{n_j}$. Let $W_k = V_j \cap \{u_k = 0\}$. Repeat the similar process for the irreducible components of the nonempty sets W_k . Thus we obtain the sets $W_{k,j}^{n_{k,j}} \subset V_j^{n_j}$, where n_k are their dimension. There is a map $\psi: X \cap \mathcal{U} \rightarrow V_j^{n_j}$ for all j . By the

construction, $(n-1-n_j)$ -dimensional singularities $\psi^{-1}(P_1)$ and $\psi^{-1}(P_2)$ have the same type (resolution) with respect to Newton diagram, where $P_1, P_2 \in (V_j^{n_j} \setminus (\bigcup_k W_{k,j}^{n_{k,j}} \cup \mathbf{0})) \setminus (\bigcup_{i \neq j} V_i^{n_i})$. It should be noted that they can be not biholomorphic. This type is denoted by $\mathcal{F}_j$. The similar statement is true for the map $\psi: X \cap \mathcal{U} \rightarrow W_{k,j}^{n_{k,j}}$ and the set $(W_{k,j}^{n_{k,j}} \setminus \mathbf{0}) \setminus \bigcup_{l \neq k} W_{l,j}^{n_{l,j}}$. The corresponding type is denoted by $\mathcal{F}_{j,k}$.

Examples 1.25. 1). Let us consider the singularity $(x_1^2 + x_2^4 + (x_3^2 + x_4^3)^2 x_4 = 0, 0) \subset (\mathbb{C}^5, 0)$. Then $V_1^2 = \{x_1 = x_2 = x_3^2 + x_4^3 = 0\}$, $W_{1,1}^1 = V_1^2 \cap \{x_4 = 0\}$, $\mathcal{F}_1 = \{(y_1^2 + y_2^4 + y_3^2 = 0, 0) \subset (\mathbb{C}^3, 0)\}$, $\mathcal{F}_{1,1} = \{(y_1^2 + y_2^4 + y_3^2 y_4 = 0, 0) \subset (\mathbb{C}^4, 0)\}$.
2). Let us consider the singularity $(x_1^2 + x_2^3 + (x_3^5 + x_4^5)x_5^2 + x_3^3 x_4^3 x_5 = 0, 0) \subset (\mathbb{C}^5, 0)$. Then $V_1^1 = \{x_1 = x_2 = x_3 = x_4 = 0\}$, $V_2^1 = \{x_1 = x_2 = x_3 = x_5 = 0\}$, $V_3^1 = \{x_1 = x_2 = x_4 = x_5 = 0\}$, $\mathcal{F}_1 = \{(y_1^2 + y_2^3 + y_3^5 + y_4^5 + cy_3^3 y_4^3 = 0, 0) \subset (\mathbb{C}^4, 0)\}$ and for the different values of c the singularities can be not biholomorphic, $\mathcal{F}_2 = \mathcal{F}_3 = \{(y_1^2 + y_2^3 + y_4^2 + y_3^3 y_4 = 0, 0) \subset (\mathbb{C}^4, 0)\}$.

Proposition 1.26. *Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a normal nonisolated hypersurface singularity. It is terminal outside $\mathbf{0}$ (resp. canonical outside $\mathbf{0}$, log canonical outside $\mathbf{0}$) if and only if $\mathcal{F}_j$ and $\mathcal{F}_{j,k}$ are terminal (resp. canonical, log canonical) for all j, k .*

Proof. For instance, let us prove this theorem in the terminal case. In definition 1.24 we made the subdivision of $\text{Sing } X \setminus \mathbf{0}$ to the "cells". Every cell has a dimension not less than one. Along every cell X has the same singularity type. Let some $\mathcal{F}_j$ or $\mathcal{F}_{j,k}$ be not terminal singularity. Then there exists a blow-up with a center along the corresponding cell and we obtain an exceptional divisor with discrepancy ≤ 0 . Contradiction.

Conversely assume that $\mathcal{F}_j$ and $\mathcal{F}_{j,k}$ are the terminal singularities and $(X, 0)$ is not terminal outside $\mathbf{0}$. Consider the log-resolution ψ . Then there exists an exceptional divisor E with discrepancy $a(E, 0) \leq 0$ such that $\dim \psi(E) \geq 1$. The generic point $P \in \psi(E)$ is lying in the fixed cell or $P \notin \text{Sing } X$. In both cases we have a contradiction with $a(E, 0) \leq 0$. $\square$

Definition 1.27. Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a nonnormal hypersurface singularity. It is called a *log canonical singularity* if $(\mathbb{C}^n, X)$ is lc. It is called *log canonical outside $\mathbf{0}$ singularity* if $(\mathbb{C}^n, X)$ is log canonical outside $\mathbf{0}$. The last definition is equivalent to the conditions that $\mathcal{F}_j$ and $\mathcal{F}_{j,k}$ are lc.

2. Quasihomogeneous hypersurface singularities

Definition 2.1. Let $f: (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$ be a polynomial. It is called a *quasihomogeneous polynomial of degree d* if there exist the strictly positive rational numbers $p_1, \dots, p_n$ (which are called *the weights*) that

$$(1) \quad f(\lambda^{p_1}x_1, \dots, \lambda^{p_n}x_n) = \lambda^d f(x_1, \dots, x_n)$$

for any λ and all $x_1, \dots, x_n$.

For the quasihomogeneous singularities (polynomials) the nonnegative rational numbers $p_1, \dots, p_n$ satisfying condition (1) are also called *the weights* and $d \neq 0$ is called a *quasihomogeneous degree* too.

Convention. Without loss of generality it can be assumed that $p_1, \dots, p_n, d$ are the integer numbers in this chapter.

Lemma 2.2. Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a quasihomogeneous hypersurface singularity defined by a polynomial f . If $\frac{\partial f}{\partial x_1}(\mathbf{x}) = \dots = \frac{\partial f}{\partial x_n}(\mathbf{x}) = 0$ then $f(\mathbf{x}) = 0$.

Proof. The polynomial f is quasihomogeneous if and only if $f \in I_{\nabla f}$, where $I_{\nabla f} = (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$ is a gradient ideal [13]. This concludes the proof. $\square$

Proposition 2.3. Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a normal quasihomogeneous hypersurface singularity with weights $p_1, \dots, p_n$ and degree d defined by a polynomial f . Then

1. If $\sum p_i \geq d+1$ and $(X, 0)$ is canonical outside $\mathbf{0}$ then $\mathbf{1} \in \Gamma_+(f)^0$.
2. If $\sum p_i \geq d$ and $(X, 0)$ is log canonical outside $\mathbf{0}$ then $\mathbf{1} \in \Gamma_+(f)$.

Proof. Assume the converse. Then $\mathbf{1} \notin \Gamma_+(f)^0$ or $\mathbf{1} \notin \Gamma_+(f)$ respectively. Then there exists a primitive vector $\mathbf{p}'$ from $\sigma^\vee$ such that the plane $\Pi = \{l \in N_{\mathbb{R}} \mid \langle \mathbf{p}', l \rangle = \mathbf{p}'(f)\}$ is the plane of support for Newton polyhedron. This plane contains $\mathbf{1}$ or lies over $\mathbf{1}$. Let $\varphi: \mathbb{C}^n(\mathbf{p}') \rightarrow \mathbb{C}^n$ be a $\mathbf{p}'$ -blow-up. Since the plane of support Π is parallel to some coordinate axis then at least one coordinate of $\mathbf{p}'$ is equal to zero. Therefore $\dim \varphi(D_{\mathbf{p}'}) \geq 1$. Also the discrepancy $a(D_{\mathbf{p}'}, X) = \langle \mathbf{p}', \mathbf{1} \rangle - \mathbf{p}'(f) - 1$ is less than -1 in the first case and strictly less than -1 in the second case. Since the singularity is canonical outside $\mathbf{0}$ or lc outside $\mathbf{0}$ we obtain the contradiction respectively. $\square$

The next similar proposition for a nonnormal singularity is proved by the same arguments.

Proposition 2.4. Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a nonnormal quasihomogeneous hypersurface singularity with weights $p_1, \dots, p_n$ and degree d

defined by a polynomial f . Let $\sum p_i \geq d$ and $(X, 0)$ is log canonical outside $\mathbf{0}$ (see definition 1.27) then $\mathbf{1} \in \Gamma_+(f)$.

2.5. The quasihomogeneous singularity resolution construction. For every hypersurface singularity $(X, 0) \subset (\mathbb{C}^n, 0)$ defined by a polynomial f there exists a subdivision Δ of σ in N [25]) depending only on Newton polyhedron $\Gamma_+(f)$ with the following properties. The variety $T_N(\Delta)$ is nonsingular, therefore $\text{Diff}_{X(\Delta)}(0) = 0$, where $X(\Delta)$ denotes the proper transform of X on $T_N(\Delta)$. If $(X, 0)$ is a non-degenerate singularity then the birational morphism $\psi: T_N(\Delta) \rightarrow \mathbb{C}^n$ is the toric log-resolution of $(\mathbb{C}^n, X)$ [25]. The value of non-degenerate singularities is in the presence of such toric embedded log-resolution. In any case the birational morphism ψ is called a *partial log-resolution*. Thus $K_{T_N(\Delta)} + X(\Delta) + E(\Delta) = \psi^*(K_{\mathbb{C}^n} + X)$.

Convention. Let us assume in addition that $f(x_1, \dots, x_n)$ is a quasihomogeneous polynomial. Let us choose its weights $\mathbf{p} = (p_1, \dots, p_n)$ that $p_i \neq 0$ for any i .

Example 2.6. If $(X, 0)$ is an isolated singularity then ψ is its embedded toric log-resolution.

(1). Let $(X, 0)$ be a normal singularity. It is equivalent to demand that $\text{codim}_{\mathbb{C}^n} \text{Sing } X \geq 3$, i.e. $n_i \leq n - 3$ for all i (see definition 1.24). Then $X(\Delta)$ is normal and $K_{X(\Delta)} = \psi^*K_X + \sum a(E_i, 0)E_i$. As in the example 2.6 variety $X(\Delta)$ can't have the isolated singularities. Therefore $\text{Sing } X(\Delta) = \bigcup_{i \in I} \tilde{\mathcal{A}}_i^{n_i}$, where $\tilde{\mathcal{A}}_i^{n_i}$ are the irreducible components and $\psi(\tilde{\mathcal{A}}_i^{n_i}) = \mathcal{A}_i^{n_i}$ (see definition 1.24).

Definition 2.7. $\mathcal{A}_i^{n_i}$ is called a *non-degenerate component* of singularity if $i \notin I$. $\mathcal{A}_i^{n_i}$ is called a *degenerate component* of singularity if $i \in I$.

We can recognize these components in the following way. Let us slightly modify the coefficients of f in order that the singularity became non-degenerate. If the component $\mathcal{A}_i^{n_i}$ of our singularity remains the component of new singularity then it is non-degenerate. If it doesn't occur then this component is degenerate.

In other words the morphism ψ is the log-resolution only for the non-degenerate singularity component.

Let us define $\tilde{V}_j^{n_j}, \tilde{W}_{k,j}^{n_{k,j}}$ as before $V_j^{n_j}, W_{k,j}^{n_{k,j}}$ in the definition 1.24. Notice that $\psi(\tilde{V}_j^{n_j}) = V_j^{n_j}, \psi(\tilde{W}_{k,j}^{n_{k,j}}) = W_{k,j}^{n_{k,j}}$ and $\tilde{V}_j^{n_j}, \tilde{W}_{k,j}^{n_{k,j}} \not\subset \text{Exc}(\psi)$ for all j, k .

Lemma 2.8. *Along the sets $\widetilde{V}_j^{n_j} \setminus ((\bigcup_k \widetilde{W}_{k,j}^{n_{k,j}}) \cup (\bigcup_{i \neq j} \widetilde{V}_i^{n_i}))$, $\widetilde{W}_{k,j}^{n_{k,j}} \setminus \bigcup_{l \neq k} \widetilde{W}_{l,j}^{n_{l,j}}$ we have the same singularity types $\mathcal{F}_j, \mathcal{F}_{k,j}$ respectively (cf. definition 1.24).*

Proof. The exceptional divisor corresponding to $\mathbf{p}$ -blow-up is denoted by $E_{\mathbf{p}}$. We have to prove that the type of hypersurface singularities is the same one in the intersection point of these sets with $E_{\mathbf{p}}$. This means that the singularity given by a polynomial

$$\hat{f} = x_i^{-d} f(x_1 x_i^{p_1}, \dots, x_i^{p_i}, \dots, x_n x_i^{p_n}) = f(x_1, \dots, 1, \dots, x_n)$$

in the neighborhood of $E_{\mathbf{p}}$ coincides with the singularity given by a polynomial f in the neighborhood of $x_i = 1$. For this purpose it is enough to prove that $\frac{\partial \hat{f}}{\partial x_i}(\mathbf{x}) = 0$, where $\mathbf{x} \in S = \text{Sing}\{\hat{f} = 0\}$. Since

$$\begin{aligned} \frac{\partial(x_i^d \hat{f})}{\partial x_i} &= \sum_{j \neq i} p_j \frac{\partial f}{\partial x_j}(x_1 x_i^{p_1}, \dots) x_j x_i^{p_j-1} + p_i \frac{\partial f}{\partial x_i}(x_1 x_i^{p_1}, \dots) x_i^{p_i-1} = \\ &= \frac{\partial(x_i^d f(x_1, \dots, 1, \dots, x_n))}{\partial x_i} = d x_i^{d-1} f(x_1, \dots, 1, \dots, x_n), \end{aligned}$$

then substituting any $\mathbf{x} = (x_1^0, \dots, 1, \dots, x_n^0) \in S$ we get $p_i \frac{\partial f}{\partial x_i}(x_1^0, \dots, 1, \dots, x_n^0) = 0$. Q.E.D. $\square$

By the same argument $\bigcup \text{Sing } E_i \subset \text{Sing } X(\Delta)$. Therefore there exists a log-resolution $\varphi: \widetilde{X} \rightarrow X(\Delta)$ such that $\dim \varphi(E) \geq 1$, where E is any component of $\text{Exc}(\varphi)$. Moreover, the center of every exceptional divisor doesn't lie in $\bigcup \text{Sing } E_i$. Such resolution is called a *good one*.

(2). Let $(X, 0)$ be a nonnormal reduced singularity, i.e. $\text{codim}_{\mathbb{C}^n} \text{Sing } X \geq 2$ ($n_i \leq n - 2$ for all i). Similarly there is the same definition of the *non-degenerate and degenerate singularity components*. In the same way we construct a *good log-resolution* φ of $(T_N(\Delta), X(\Delta) + E(\Delta))$ with the same properties. In particular the center of every exceptional divisor doesn't lie in $\text{Sing } E(\Delta)$.

Theorem 2.9. *Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a normal quasihomogeneous hypersurface singularity with weights $p_1, \dots, p_n$ and degree d defined by a polynomial f . Then*

1. $\sum p_i \geq d+1$ and $(X, 0)$ is canonical outside $\mathbf{0}$ if and only if $(X, 0)$ is canonical.
2. $\sum p_i \geq d$ and $(X, 0)$ is log canonical outside $\mathbf{0}$ if and only if $(X, 0)$ is log canonical.

Proof. The sufficient conditions are obvious. Now we prove their necessity. Let $\psi: T_N(\Delta) \rightarrow \mathbb{C}^n$ be a partial log-resolution (see 2.5). Let us prove the first assertion. By proposition 2.3 we have $\mathbf{1} \in \Gamma_+(f)^0$. Hence $a_i(E_i, 0) \geq 0$ for all i [14]. Since $(X, 0)$ is canonical outside $\mathbf{0}$ then $\mathcal{F}_i, \mathcal{F}_{k,j}$ are canonical by proposition 1.26. This implies that $X(\Delta)$ is canonical. Therefore $(X, 0)$ is a canonical singularity too. The second assertion is proved similarly. Arguing as above, we see that $a(E_i, 0) \geq -1$, $\mathcal{F}_i, \mathcal{F}_{k,j}$ are the lc singularities and $X(\Delta)$ is lc. By taking a good log-resolution we conclude that $(X, 0)$ is lc since $\widetilde{V}_j^{n_j}, \widetilde{W}_{k,j}^{n_{k,j}} \not\subseteq \text{Exc}(\psi)$ for all j, k . $\square$

Remark 2.10. By theorem 2.9 and proposition 1.26 the question about the concrete quasihomogeneous singularity type is reduced to the same one about $\mathcal{F}_i$ and $\mathcal{F}_{i,k}$ of lesser dimension.

The next similar criterion for the nonnormal singularity is proved by the same arguments.

Theorem 2.11. *Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a nonnormal quasihomogeneous hypersurface singularity with weights $p_1, \dots, p_n$ and degree d . Then $\sum p_i \geq d$ and $(X, 0)$ is log canonical outside $\mathbf{0}$ if and only if $(X, 0)$ is log canonical.*

Example 2.12. Consider the singularities from the example 1.25. By theorem 2.9 the first singularity is canonical. The second one is terminal since it is terminal outside $\mathbf{0}$ and its hyperplane section $\{x_3 = x_5\}$ has the canonical singularities.

Theorem 2.13. *Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a normal log canonical quasihomogeneous hypersurface singularity with weights $p_1, \dots, p_n$ defined by a polynomial f . Then*

1. *If $(X, 0)$ is not canonical outside $\mathbf{0}$ then $\mathbf{p}$ -blow-up is lc, but not plt blow-up, except the case described in the example 2.15(1). In particular, $(X, 0)$ is always nonexceptional singularity.*
2. *(cf. [6, 3.3]) If $(X, 0)$ is canonical outside $\mathbf{0}$ then $\mathbf{p}$ -blow-up is plt one. Assume in addition that $(X, 0)$ is strictly log canonical then $(X, 0)$ is exceptional.*

Proof. Let $\psi: \mathbb{C}^n(\mathbf{p}) \rightarrow \mathbb{C}^n$ be a $\mathbf{p}$ -blow-up. Its exceptional locus is

$$\text{Exc}(\psi|_{X(\mathbf{p})}) = E_{\mathbf{p}} = D_{\mathbf{p}} \cap X(\mathbf{p}) = \{f(x_1, \dots, x_n) \subset \mathbb{P}(p_1, \dots, p_n)\}.$$

Since f is an irreducible polynomial then $E_{\mathbf{p}}$ is an irreducible variety. It is clear that $-E_{\mathbf{p}}$ is relatively ample and $X(\mathbf{p})$ is a normal variety.

Consider the partial log-resolution of X . We have the corresponding morphism $\varphi: T_N(\Delta) \rightarrow \mathbb{C}^n(\mathbf{p})$.

$$\begin{array}{ccccc} T_N(\Delta) & \xrightarrow{\varphi} & \mathbb{C}^n(\mathbf{p}) & \xrightarrow{\psi} & \mathbb{C}^n \\ \cup & & \cup & & \cup \\ \tilde{X} \longrightarrow X(\Delta) & \xrightarrow{\varphi|_{X(\Delta)}} & X(\mathbf{p}) & \xrightarrow{\psi|_{X(\mathbf{p})}} & X \end{array}$$

One has

(2)

$$K_{T_N(\Delta)} + X(\Delta) + D_{\mathbf{p}} = \varphi^*(K_{\mathbb{C}^n(\mathbf{p})} + X(\mathbf{p}) + D_{\mathbf{p}}) + \sum_{\mathbb{R}_{\geq 0} \mathbf{q} \in \Delta(1), \mathbf{q} \neq \mathbf{p}} \alpha_{\mathbf{q}} D_{\mathbf{q}}.$$

By abuse of notation the divisors corresponding to $\mathbf{p}$ on $T_N(\Delta)$ and on $\mathbb{C}^n(\mathbf{p})$ are both denoted by $D_{\mathbf{p}}$. Let $\mathbf{q} \in \sigma_i$ be a primitive vector. By the proof of lemma [6, 3.2] we have

$$\alpha_{\mathbf{q}} = a(D_{\mathbf{q}}, X(\mathbf{p})) - \frac{q_i}{p_i} = \langle \mathbf{q}, 1 \rangle - \mathbf{q}(f) - \frac{q_i}{p_i}(\langle \mathbf{p}, 1 \rangle - \mathbf{p}(f)) - 1.$$

Let us prove the first assertion. Obviously $\mathbf{1} \in \Gamma_+(f)$. If $q_i = 0$ then $\alpha_{\mathbf{q}} = \langle \mathbf{q}, 1 \rangle - \mathbf{q}(f) - 1 \geq -1$. If $q_i > 0$ then we can define $\mathbf{a} = (a_1, \dots, a_n)$ from the proportion $(q_1 : \dots : q_n) = (p_1 + a_1 : \dots : p_n + a_n)$. Then $a_i = 0$ and $a_j \geq 0$ for all $j \neq i$. Substitute $\mathbf{q} = \frac{q_i}{p_i}(\mathbf{p} + \mathbf{a})$. Then $\alpha_{\mathbf{q}} = \frac{q_i}{p_i}(\langle \mathbf{a}, 1 \rangle - \mathbf{a}(f)) - 1 \geq -1$.

Lemma 2.14. $\text{Diff}_{X(\mathbf{p})}(0) = 0$.

Proof. If $\text{codim}_{\mathbb{C}^n(\mathbf{p})} \text{Sing } \mathbb{C}^n(\mathbf{p}) > 2$ then there is nothing to prove. Let $\text{codim}_{\mathbb{C}^n(\mathbf{p})} \text{Sing } \mathbb{C}^n(\mathbf{p}) = 2$. There are only two cases.

(1) $(p_1, \dots, p_{j-1}, p_{j+1}, \dots, p_n) = d > 1$ and $p_j \neq 0$. Hence $\psi(D_{\mathbf{p}}) = 0$.

Consider any chart of $\mathbb{C}^n(\mathbf{p})$

$$U_k = \mathbb{C}_{y_1, \dots, y_n}^n / \mathbb{Z}_{p_k}(-p_1, \dots, -p_{k-1}, 1, -p_{k+1}, \dots, -p_n), \quad k \neq j.$$

The variety $X(\mathbf{p})$ is given by an equation $f(y_1, \dots, y_{k-1}, 1, y_{k+1}, \dots, y_n) = 0$ in this chart. If $\text{Diff}_{X(\mathbf{p})}(0) \neq 0$ then $f(\mathbf{y}) = 0$ for all $\mathbf{y} = (y_1, \dots, y_{k-1}, 1, y_{k+1}, \dots, y_{j-1}, 0, y_{j+1}, \dots, y_k)$. Thus $f = x_j \cdot f'$. It is impossible.

(2) $p_i \neq 0$ if $i \in I$ and $p_i = 0$ if $i \notin I$, where set $I \subset \{1, \dots, n\}$ and $2 \leq k = |I| \leq n - 1$. Hence $\dim \psi(D_{\mathbf{p}}) = n - k$.

In the small neighborhood $\mathcal{U}$ of the generic point $P \in \psi(D_{\mathbf{p}})$ the variety X is isomorphic to $\mathbb{C}^{n-k} \times X'$. The morphism ψ induces $\mathbf{p}'$ -blow-up

$\mathbb{C}^k(\mathbf{p}') \rightarrow \mathbb{C}^k$ and $X'(\mathbf{p}') \rightarrow X'$, where $p'_i = p_i$ for $i \in I$. By the above statement $\text{Diff}_{X'(\mathbf{p}')} (0) = 0$. Hence $\text{Diff}_{X(\mathbf{p})} (0) = 0$. $\square$

From this lemma and equality (2) we have

$$K_{X(\Delta)} + \tilde{E}_{\mathbf{p}} = \varphi^*(K_{X(\mathbf{p})} + E_{\mathbf{p}}) + \sum \alpha'_{\mathbf{q}} D'_{\mathbf{q}}, \text{ where } \alpha'_{\mathbf{q}} \geq -1.$$

Consider a good log-resolution $\tilde{X} \rightarrow X(\Delta)$. All exceptional divisors for the pair $(X(\Delta), \tilde{E}_{\mathbf{p}} - \sum \alpha'_{\mathbf{q}} D'_{\mathbf{q}})$ have the discrepancies not less than -1 since $\tilde{V}_j^{n_j}, \tilde{W}_{k,j}^{n_{k,j}} \not\subseteq \text{Exc}(\psi \circ \varphi)$. By assumption $(X, 0)$ is not canonical outside $\mathbf{0}$. Thus $(X(\mathbf{p}) + E_{\mathbf{p}})$ is not plt or the case described in the example 2.15(1) takes place. By propositions 1.17 and 1.18 $(X, 0)$ is not exceptional.

Let us prove the second assertion. Arguing as above it is enough to prove that $\alpha_{\mathbf{q}} = \langle \mathbf{q}, 1 \rangle - \mathbf{q}(f) - 1 > -1$ if $q_i = 0$ and $\langle \mathbf{a}, 1 \rangle - \mathbf{a}(f) > 0$ if $q_i > 0$. Actually, if $\alpha_{\mathbf{q}} = -1$ then by considering $\mathbf{q}$ -blow-up or $\mathbf{a}$ -blow-up we have the contradiction that our singularity is canonical outside $\mathbf{0}$. It can happen that $\mathbf{a}$ -blow-up is undefined (see 1.21). Then $\mathbf{a}(f) = 0$ and there is nothing to prove or f is not irreducible, i.e. we have the contradiction with the normality of $(X, 0)$.

Assume in addition that $(X, 0)$ is strictly lc. Since $(X, 0)$ is canonical outside $\mathbf{0}$ then $p_i \neq 0$ for all i . Thus $\dim \psi(E_{\mathbf{p}}) = 0$. By theorem [12, 1.9] $(X, 0)$ is exceptional. $\square$

Example 2.15. (1) Let $(X, 0)$ be a normal strictly log canonical quasihomogeneous hypersurface singularity with weights $p_1, \dots, p_n$. Let $\mathbf{p}$ -blow-up ψ is plt one and $p_i = 0$ for some i . It means there exists only one index k such that $\mathcal{F}_k$ is strictly lc. Note also that $\mathcal{F}_k$ is exceptional, the corresponding component $\mathcal{A}_k^{n_k}$ is non-degenerate and $\psi(E_{\mathbf{p}}) = \mathcal{A}_k^{n_k}$.

(2) Let $(X, 0)$ be a normal strictly log canonical quasihomogeneous hypersurface singularity with weights $p_1, \dots, p_n$ defined by a polynomial f . Let $(X_{\mathbf{p}}, 0) = (f_{\mathbf{p}} = 0, 0)$ be a singularity of part (1) for some $\mathbf{p} \in N_{\mathbb{R}}$. Then $\mathbf{p}$ -blow-up is also plt one.

Example 2.16. Consider the singularity $(x_1^2 + x_2^3 + x_3^6 = 0, 0) \subset (\mathbb{C}^4, 0)$ with weights $\mathbf{p} = (3, 2, 1, a)$. If $a = 0$ then the $\mathbf{p}$ -blow-up is a plt one. If $a \neq 0$ then the $\mathbf{p}$ -blow-up is a lc but not plt one.

For the nonnormal singularities there is the similar theorem.

Theorem 2.17. Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a nonnormal log canonical quasihomogeneous hypersurface singularity with weights $\mathbf{p} = (p_1, \dots, p_n)$ defined by a polynomial f . Consider $\mathbf{p}$ -blow-up $\psi: \mathbb{C}^n(\mathbf{p}) \rightarrow \mathbb{C}^n$. Then $K_{\mathbb{C}^n(\mathbf{p})} + X(\mathbf{p}) + D_{\mathbf{p}}$ is lc but not plt.

By theorem 2.13 we have the following important corollary.

Corollary 2.18. *Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a normal hypersurface singularity defined by a polynomial f and let $\mathbf{p} \in N_{\mathbb{R}}$. Put $(X_{\mathbf{p}}, 0) = (f_{\mathbf{p}} = 0, 0)$. Then*

1. *If $(X_{\mathbf{p}}, 0)$ is canonical (resp. log canonical) then $(X, 0)$ is also canonical (resp. log canonical).*
2. *Consider $\mathbf{p}$ -blow-ups $\varphi: (X(\mathbf{p}), E_{\mathbf{p}}) \rightarrow (X, 0)$ and $\varphi_{\mathbf{p}}: (X_{\mathbf{p}}(\mathbf{p}), E'_{\mathbf{p}}) \rightarrow (X_{\mathbf{p}}, 0)$. Then $(E_{\mathbf{p}}, \text{Diff}_{E_{\mathbf{p}}}(0)) = (E'_{\mathbf{p}}, \text{Diff}_{E'_{\mathbf{p}}}(0))$.*
3. *If $(X_{\mathbf{p}}, 0)$ is log canonical and not canonical outside $\mathbf{0}$ then $\mathbf{p}$ -blow-up is lc but not plt one of $(X, 0)$ except the case described in the example 2.15(2). In particular $(X, 0)$ is always not exceptional.*
4. *If $(X_{\mathbf{p}}, 0)$ is log canonical and canonical outside $\mathbf{0}$ then $\mathbf{p}$ -blow-up is plt one of $(X, 0)$. Assume in addition that $(X, 0)$ is strictly log canonical. Then $(X, 0)$ is exceptional.*

Proof. Consider the deformation $F_t = t^{-\mathbf{p}(f)} f(t^{p_1} x_1, \dots, t^{p_n} x_n)$. If $t = 0$ then $F_0 = f_{\mathbf{p}}$. For small $t \neq 0$ singularity F_t has the type f . Let $(X_{\mathbf{p}}, 0)$ is lc then $(\mathbb{C}^n, X_{\mathbf{p}})$ is lc by lemma [10, 7.1.3]. By Inversion of Adjunction and [10, 7.8] the first statement is true. The statements (2) and (4) are trivial. To prove the third statement it is enough to show that $(\mathbb{C}^n(\mathbf{p}), X(\mathbf{p}) + D_{\mathbf{p}})$ is lc. Consider the same deformation. If $t = 0$ then $(\mathbb{C}^n(\mathbf{p}), X_{\mathbf{p}}(\mathbf{p}) + D_{\mathbf{p}})$ is lc by the proof of theorem 2.13. Therefore corollary [10, 7.8] completes the proof. $\square$

Corollary 2.19. *Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be a normal log canonical hypersurface singularity defined by a polynomial f and $(f_{\mathbf{p}} = 0, 0)$ be a nonnormal log canonical singularity for some $\mathbf{p} \in N_{\mathbb{R}}$. Then $\mathbf{p}$ -blow-up is lc but not plt one of $(X, 0)$. In particular $(X, 0)$ is not exceptional.*

Proof. Since $f_{\mathbf{p}}$ is a reduced polynomial then it is easy to check that $\dim \text{Sing } X(\mathbf{p}) \leq n - 2$ in any chart for $\mathbf{p}$ -blow-up [20, 3.7]. Therefore $X(\mathbf{p})$ is normal. The rest is proved as above corollary 2.18 (applying theorem 2.17). $\square$

Remark 2.20. For every non-degenerate log canonical (resp. canonical) singularity there always exists $\mathbf{p} \in N_{\mathbb{R}}$ such that $f_{\mathbf{p}}$ defines a log canonical (resp. canonical) singularity [6, 3.5].

3. Three-dimensional log canonical hypersurface singularities

Now we prove the main theorem formulated in the introduction.

Proof of the main theorem. According to theorem 3.2 the exceptional strictly log canonical (resp. canonical) hypersurface singularities are divided into two types.

(1) There exists a primitive $\mathbf{p} \in N_{\mathbb{R}}$ that $\tilde{f}_{\mathbf{p}} = (f \circ \psi)_{\mathbf{p}}$ gives a strictly log canonical and canonical outside $\mathbf{0}$ (resp. canonical) singularity.

(2) $\tilde{f}_{\mathbf{p}} = t^3 + g_2^2(z, x, y)$, where g_2 is an irreducible homogeneous polynomial of degree 2.

The exceptional singularities of second type are classified in the theorems 3.3 and 3.5.

Consider the singularities of the first type. The primitive vector $\mathbf{p}$ is unique. Indeed, let there exist two different $\mathbf{p}_1$ and $\mathbf{p}_2$. Then by lemma [6, 4.1] and corollary 2.18 they give two different lc blow-ups. Therefore $(X, 0)$ is not exceptional by proposition 1.17. Thus $\mathbf{p}$ -blow-up $\varphi: (X(\mathbf{p}), E_{\mathbf{p}}) \rightarrow (X, 0)$ is plt one by corollary 2.18.

Let $\tilde{f}_{\mathbf{p}}$ defines a strictly log canonical singularity. By corollary 2.18 $(X, 0)$ is exceptional. Since $\dim \varphi(E_{\mathbf{p}}) = 0$ (theorem [12, 1.9]) and $\langle \mathbf{p}, 1 \rangle = \mathbf{p}(f)$ then it is easy to prove that $\text{Diff}_{E_{\mathbf{p}}/\mathbb{P}(\mathbf{p})}(0) = 0$ and $\text{Diff}_{E_{\mathbf{p}}}(0) = 0$. Hence $E_{\mathbf{p}}$ is a surface with Du Val singularities and $K_{E_{\mathbf{p}}} \sim 0$. By lemma [2, 1.4.1] $E_{\mathbf{p}}$ is a singular K3 surface.

Let $\tilde{f}_{\mathbf{p}}$ defines a canonical singularity. Then under condition — $\text{Diff}_{E_{\mathbf{p}}/\mathbb{P}(\mathbf{p})}(0) = 0$ such singularities are classified in the theorems 3.24, 3.28 and in the tables of chapter §4. The proof of this classification begins from the subsection 3.6. Q.E.D.

Definition 3.1. A quasihomogeneous hypersurface singularity f belongs to the type $\mathcal{M}_i$ if the maximal compact face of Newton polyhedron $\Gamma_+(f)$ has the dimension i .

The next theorem 3.2 divides the three-dimensional exceptional hypersurface singularities into two types.

Theorem 3.2. *Let $(Z, 0) \subset (\mathbb{C}^4, 0)$ be a three-dimensional exceptional canonical (resp. strictly log canonical) hypersurface singularity defined by a polynomial g . Then there exists a biholomorphic coordinate change $\psi: (\mathbb{C}^4, 0) \rightarrow (\mathbb{C}_{t,z,x,y}^4, 0)$ and unique primitive vector $\mathbf{p} \in N_{\mathbb{R}}$ such that just one of the following two possibilities holds:*

1. $\tilde{g}_{\mathbf{p}} = (g \circ \psi)_{\mathbf{p}}$ defines an exceptional canonical (resp. strictly log canonical and canonical outside $\mathbf{0}$) singularity.
2. $\tilde{g}_{\mathbf{p}} = t^3 + g_2^2(z, x, y)$, where g_2 is an irreducible homogeneous polynomial of degree 2.

Proof. At first consider the canonical singularity case. If g is a quasihomogeneous polynomial then there is nothing to prove. Take $\mathbf{p} \in N_{\mathbb{R}}$ such that $g_{\mathbf{p}} \in \mathcal{M}_3$, $\mathbf{1} \in \Gamma_+(g_{\mathbf{p}})^0$ and $\mathbf{p}(g) = 1$. It is clear that

2 or 3-jet of $g_{\mathbf{p}}$ is nonzero. Let $g_{\mathbf{p}}$ be not reduced polynomial, i.e. $g_{\mathbf{p}} = (g'_{\mathbf{p}})^k$, where $k \geq 2$. Without loss of generality it can be assumed that $g'_{\mathbf{p}} = t + g''$. After the quasihomogeneous coordinate change $\psi: t \mapsto t - g''$ we get $\tilde{g} = g \circ \psi = t^k + \tilde{g}'$ and $\tilde{g}_{\mathbf{p}} = t^k$. Repeat the described above process for $\tilde{g}$. Since $\mathbf{1} \in \Gamma_+(g)^0$ then this process will be finished. Hence we may assume that $g_{\mathbf{p}}$ is a reduced polynomial.

Let $(Z_{\mathbf{p}}, 0) = (\{g_{\mathbf{p}} = 0\}, 0)$ be a nonnormal singularity. Then all components $\mathcal{A}_i^2$ are degenerate (cf. [6, 3.3]). Indeed, let $\mathcal{A}_j^2$ be a non-degenerate component. Take the partial resolution ψ . Then there is an exceptional divisor E with discrepancy $a(E, 0) \leq -1$ such that $\psi(E) = \mathcal{A}_j^2$. We obtain the contradiction with $\mathbf{1} \in \Gamma_+(g_{\mathbf{p}})^0$. By corollary 2.19 we have a nonnormal, not log canonical singularity along some component $\mathcal{A}_k^2$. In the neighborhood of the generic point of $\mathcal{A}_k^2$ we have $Z_{\mathbf{p}} \cong \mathbb{C}_{f_1, f_2}^2 \times \{\Phi(f_1(t, z, x, y), f_2(t, z, x, y)) = 0\}$, $\deg \Phi \geq 2$. Since $\mathbf{1} \notin \Gamma_+(\Phi)$ and 2 or 3-jet of $g_{\mathbf{p}}$ is nonzero it follows that after the quasihomogeneous biholomorphic coordinate change we get $f_1 = t$. If $\text{mult}_0 f_2 = 1$ then it is not difficult to show that monomial x, y or z belongs to f_2 . Thus, after some biholomorphic coordinate change we can assume that $\text{mult}_0 f_2 \geq 2$. Let $\Phi_{\mathbf{q}}$ be a quasihomogeneous part of Φ ($\mathbf{q}(\Phi) = 1$), which defines an isolated singularity. Then $\text{wt } z + \text{wt } x + \text{wt } y \leq 3 \cdot \frac{1}{2} \text{wt } f_2$. The equality holds if and only if $f_2 = g_2$ is an irreducible homogeneous polynomial of degree 2. Using this fact in solving the following system

$$\begin{cases} \text{wt } t + \text{wt } z + \text{wt } x + \text{wt } y > 1 \\ \text{wt } f_1 + \text{wt } f_2 < 1 \\ \text{wt } f_1, \text{wt } f_2 \in \{\frac{1}{2}, \frac{1}{3}\} \cup (0, \frac{1}{3}) \end{cases}$$

we get $\Phi_{\mathbf{q}} = f_1^3 + f_2^2$. Therefore $g_{\mathbf{p}} = t^3 u_1(t, z, x, y) + g_2^2 u_2(t, z, x, y)$. Since $\mathbf{1} \in \Gamma_+(g_{\mathbf{p}})^0$ then u_1, u_2 are the constants and we have the second case in the theorem condition.

Let $(Z_{\mathbf{p}}, 0)$ be a normal, not canonical singularity. By theorem 2.9 $(Z_{\mathbf{p}}, 0)$ is not canonical outside $\mathbf{0}$. By the same argument we have not canonical singularity $\Phi(f_1, f_2, f_3) = 0$ along some degenerate component $\mathcal{A}_k^1$. If $\text{mult}_0 f_i = 1$ for all i then one can take the corresponding coordinate change and repeat the process. Let $\text{mult}_0 f_3 \geq 2$. Consider a quasihomogeneous part of Φ . Then $\text{wt } t + \text{wt } z + \text{wt } x + \text{wt } y \leq \text{wt } f_1 + \text{wt } f_2 + \frac{1}{2} \text{wt } f_3 + \frac{1}{2} \text{wt } f_3 \leq 1$. Contradiction with the construction of $g_{\mathbf{p}}$. Therefore this theorem is proved for the canonical singularities.

Consider the strictly log canonical singularity case. Similarly, there is $\mathbf{p} \in N_{\mathbb{R}}$ such that $g_{\mathbf{p}} \in \mathcal{M}_3$, $\mathbf{1} \in \Gamma_+(g_{\mathbf{p}})$ and $\mathbf{p}(g) = 1$. Moreover, if $g_{\mathbf{p}}$ defines a not log canonical outside $\mathbf{0}$ singularity then $\mathbf{1} \in \Gamma_+(g_{\mathbf{p}})^0$.


Therefore, applying the discussed above arguments we can prove this theorem. $\square$

The next theorem classifies all exceptional canonical singularities in the second case of theorem 3.2.

Theorem 3.3. *Let $(X, 0) = (f = t^3 + g_2^2(z, x, y) + f'_{\geq 5}(z, x, y) = 0, 0) \subset (\mathbb{C}^4, 0)$ be a canonical singularity. Then there exists a plt blow-up $\varphi: (Y, S) \rightarrow (X \ni 0)$ such that $(S, \text{Diff}_S(0)) = (\overline{\mathbb{F}}_{10}, \frac{1}{2}H_1 + \frac{2}{3}H_2)$, where $\overline{\mathbb{F}}_{10} = \mathbb{P}(10, 1, 1)$ is a cone, H_1 and H_2 are its hyperplane sections. Moreover, $(X, 0)$ is exceptional (resp. weakly exceptional) and 6-complementary if and only if any irreducible factor of the nonzero 5-jet f'_5 has the multiplicities at most 3 (resp. at most 4).*

Proof. Consider $\mathbf{p} = (4, 3, 3, 3)$ -blow-up $\phi: (X(\mathbf{p}), E) \rightarrow X$. Then $(E, \text{Diff}_E(0)) = (t + g_2^2 \subset \mathbb{P}(4, 1, 1, 1), \frac{2}{3}\{t = 0\}) = (\mathbb{P}^2, \frac{4}{3}C)$, where C is an irreducible conic (see construction 3.12). Also $a(E, 0) = 0$. If 5-jet of f' is zero then polynomial f has a view $t^3 + g_2^2(1, x', y') + z'^6 f''$ in the chart $\mathbb{C}_{t', z', x', y'}^4 / \mathbb{Z}_3(2, 1, 0, 0)$. This new singularity is not canonical, therefore 5-jet of f' is nonzero. Now our theorem is proved by exhaustion of all cases for 5-jets.

Let $f = t^3 + g_2^2 + z^5 + f'_{\geq 6}(z, x, y)$. We have the following picture



The variety $X(\mathbf{p})$ in the neighborhood of P_i (the points P_i have the coordinates $(0, 1, y_i)$ in $\mathbb{P}^2$, where $g_2(0, 1, y_i) = 0$) is

$$(t'^3 + \omega^2 + z'^5 x'^3 + x'^6 f'' = 0 \subset (\mathbb{C}_{t', \omega, z', x'}^4, 0)) / \mathbb{Z}_3(2, 0, 0, 1).$$


Depending on the type of f' this singularity can be nonisolated along the curves l_1 and l_2 not lying on E . By considering the invariants of

group action we get $X(\mathbf{p})$ to be isomorphic to

$$(x_1^2 x_4^5 + x_1 x_3^2 + x_2^3 + f''' = 0, 0) \subset (\mathbb{C}_{x_1, x_2, x_3, x_4}^4, 0)$$

in the neighborhoods of P_i .

The surface E is given by the equations $\{x_1 = x_3 = 0\}$ in these neighborhoods. Along the curve C the variety $X(\mathbf{p})$ has cE_6 singularity. Let H be a general hyperplane section of $X(\mathbf{p})$ passing through any point from $C \setminus \{P_1, P_2\}$. Let $C' = E \cap H$. Consider the minimal resolution of H . We have the following graph



where $\tilde{C}'$ is a proper transform of C' . The numbers near graph vertices (exceptional curves) denote their corresponding discrepancies for the pair (H, cC') . This implies that log canonical threshold of (H, cC') is equal to $1/2$ and the extraction of only central graph vertex is an inductive blow-up of $(H, \frac{1}{2}C')$ (see theorem [12, 1.5]). Let $c = c(X(\mathbf{p}), E)$ be a log canonical threshold. The pair $(X(\mathbf{p}), E)$ has the same singularities in the neighborhoods of the points P_1 and P_2 . Therefore by connectedness theorem [9, 17.4] $LCS(X(\mathbf{p}), cE) = C$. Hence $c = 1/2$.

Let us take a blow-up of $\mathbb{C}^4(\mathbf{p})$ along the curve C with the weights $(6, 4, 3, 0)$. Then this blow-up induces $\psi: \tilde{X}(\mathbf{p}) \rightarrow X(\mathbf{p})$. Thus $K_{\tilde{X}(\mathbf{p})} + \tilde{S} + \frac{1}{2}E_{\tilde{X}(\mathbf{p})} = \psi^*(K_{X(\mathbf{p})} + \frac{1}{2}E)$. It is obvious that f''' doesn't influence on the exceptional surface $\text{Exc } \psi = \tilde{S} \cong \mathbb{F}_{10}$. In what follows it will be demonstrated that the proper transform $E_{\tilde{X}(\mathbf{p})}$ of E can be contracted to the point. Therefore $E_{\tilde{X}(\mathbf{p})} \cap \tilde{S}$ is a minimal section of $\mathbb{F}_{10}$. One have $\text{Diff}_{\tilde{S}}(\frac{1}{2}E_{\tilde{X}(\mathbf{p})}) = \text{Diff}_{\tilde{S}}(0) + \frac{1}{2}E_{\tilde{X}(\mathbf{p})}|_{\tilde{S}} = \frac{2}{3}l_{\infty} + \frac{1}{2}l_0 + \frac{2}{3}l'_0 + \frac{1}{2} \cdot \frac{1}{3}l_{\infty} = \frac{5}{6}l_{\infty} + \frac{1}{2}l_0 + \frac{2}{3}l'_0$, where l_{∞} is a minimal section, l_0 and l'_0 are the different zero sections. The sections l_0 and l'_0 have the intersections in the two points with multiplicities 5. These points are lying over points P_1 and P_2 . It is clear that log surface $(\tilde{S}, \text{Diff}_{\tilde{S}}(\frac{1}{2}E_{\tilde{X}(\mathbf{p})}))$ is klt, i.e. ψ is an inductive blow-up of $(X(\mathbf{p}), \frac{1}{2}E)$. Let $L_{X(\mathbf{p})} \in |-n(K_{X(\mathbf{p})} + \frac{1}{2}E)|$ be a general element from the very ample linear system over X for $(n \gg 0)$. Then $K_{X(\mathbf{p})} + \frac{1}{2}E + \frac{1}{n}L_{X(\mathbf{p})}$ is lc and numerical trivial over X . One have $K_{\tilde{X}(\mathbf{p})} + \tilde{S} + \frac{1}{2}E_{\tilde{X}(\mathbf{p})} + \frac{1}{n}L_{\tilde{X}(\mathbf{p})} = \psi^*\phi^*(K_X + \frac{1}{n}L_X)$, where $L_X = \phi(L_{X(\mathbf{p})})$. Apply $K_{\tilde{X}(\mathbf{p})} + \tilde{S} + (\frac{1}{2} + \varepsilon)E_{\tilde{X}(\mathbf{p})} + \frac{1}{n}L_{\tilde{X}(\mathbf{p})} - \text{MMP}$ ($0 < \varepsilon \ll 1$). Since $\rho(E_{\tilde{X}(\mathbf{p})}) = 1$ then at the first and unique step of MMP we have a

divisorial contraction $E_{\tilde{X}(\mathbf{p})}$ to the point. Thus we get a plt blow-up $\varphi: (Y, S) \rightarrow (X \ni 0)$ and $(S, \text{Diff}_S(0)) = (\overline{\mathbb{F}}_{10}, \frac{1}{2}H_1 + \frac{2}{3}H_2)$, where H_1 and H_2 are the hyperplane sections of cone. Divisor $K_S + \text{Diff}_S(0) + \frac{1}{6}l$ is anti-ample and not klt, where l is a cone generator passing through the intersection point of H_1 and H_2 (of multiplicity 5). By theorem 1.15 $(X, 0)$ is not weakly exceptional.

In the same way one can consider the remaining cases for 5-jets of f' . In these cases the surface S doesn't change, H_1 and H_2 are the hyperplane sections too. Only the multiplicities of H_1 and H_2 intersection change. Now our theorem follows from the next trivial lemma. $\square$

Lemma 3.4. *Log Del Pezzo surface $(\overline{\mathbb{F}}_{10}, \frac{1}{2}H_1 + \frac{2}{3}H_2)$ is exceptional (resp. weakly exceptional) and 6-complementary if and only if its hyperplane sections H_1 and H_2 have the multiplicities at most 3 (resp. at most 4) in any intersection point.*

The next theorem classifies all exceptional strictly log canonical singularities in the second case of theorem 3.2.

Theorem 3.5. *Let $(X, 0) = (f = t^3 + g_2^2(z, x, y) + f'_{\geq 5}(z, x, y) = 0, 0) \subset (\mathbb{C}^4, 0)$ be a strictly log canonical singularity. Then $f'_5 = 0$ and $f'_6 \neq 0$. Moreover $(X, 0)$ is exceptional and 1-complementary if and only if f'_6 doesn't have an irreducible factor of multiplicity 6. In this case there exists a plt blow-up $\varphi: (Y, S) \rightarrow (X \ni 0)$ such that $(S, \text{Diff}_S(0)) = (S, 0)$, where S is a surface obtained by the section contraction of the elliptic surface $S' \rightarrow \mathbb{P}^1$ to Du Val singularity of type $\mathbb{A}_1$ (kodaira dimension $\kappa(S') = 0$). The degenerate elliptic fibers of S' are always elliptic cuspidal curves. The singular points of S are Du Val singularities. They lie only in the cusp vertexes. Depending on the irreducible factors of f'_6 we have the following cases: 12 (it is a number of degenerate fibers) $\mathbb{A}_0$ (it means the singularity type in the cusp vertex; $\mathbb{A}_0$ is a smooth point by the definition); $2\mathbb{A}_2 + 8\mathbb{A}_0$; $4\mathbb{A}_2 + 4\mathbb{A}_0$; $6\mathbb{A}_2$; $2\mathbb{D}_4 + 6\mathbb{A}_0$; $2\mathbb{D}_4 + 2\mathbb{A}_2 + 2\mathbb{A}_0$; $4\mathbb{D}_4$; $2\mathbb{E}_6 + 4\mathbb{A}_0$; $2\mathbb{E}_6 + 2\mathbb{A}_2$; $2\mathbb{E}_8 + 2\mathbb{A}_0$.*

Proof. The proof is similar to the theorem 3.3. Consider $\mathbf{p} = (4, 3, 3, 3)$ - blow-up $\phi: (X(\mathbf{p}), E) \rightarrow X$. Then $(E, \text{Diff}_E(0)) = (t + g_2^2 \subset \mathbb{P}(4, 1, 1, 1), \frac{2}{3}\{t = 0\}) = (\mathbb{P}^2, \frac{4}{3}C)$, where C is an irreducible conic. Also $a(E, 0) = 0$. By theorem 3.3 $f'_5 = 0$. If $f'_6 = 0$ then a polynomial f in the chart $\mathbb{C}_{t', z', x', y'}^4 / \mathbb{Z}_3(2, 1, 0, 0)$ is $t'^3 + g_2^2(1, x', y') + z'^9 f''$. This singularity is not log canonical therefore 6-jet of f' is nonzero. Now our theorem is proved by exhaustion of all cases for 6-jets. Let us prove it in two cases: $f'_6 = z^6 + f'_{\geq 7}(z, x, y)$, $f'_6 = z^5 x + f'_{\geq 7}(z, x, y)$. Other cases are considered likewise.

Along $C \setminus \bigcup P_i$ the variety $X(\mathbf{p})$ has nonisolated strictly log canonical singularity $(x_1^3 + x_1x_3^2 + x_2^3 = 0, 0) \subset (\mathbb{C}_{x_1, x_2, x_3, x_4}^4, 0)$.

Consider the first case. Then $i = 1, 2$ and the variety $X(\mathbf{p})$ in the neighborhood of P_i (the points P_i have coordinates $(0, 1, y_i)$ in $\mathbb{P}^2$, where $g_2(0, 1, y_i) = 0$) is

$$(x_1^3x_4^6 + x_1x_3^2 + x_2^3 + f''' = 0, 0) \subset (\mathbb{C}_{x_1, x_2, x_3, x_4}^4, 0).$$

That is, P_i are the centers of log canonical singularities of $X(\mathbf{p})$. Therefore $(X, 0)$ is not exceptional by the definition.

Consider the second case. Then $i = 1, 2, 3, 4$. The variety $X(\mathbf{p})$ in the neighborhoods of P_1, P_2 is

$$(x_1^3x_4^5 + x_1x_3^2 + x_2^3 + f''' = 0, 0) \subset (\mathbb{C}_{x_1, x_2, x_3, x_4}^4, 0),$$

but in the neighborhoods of P_3, P_4 is

$$(x_1^3x_4 + x_1x_3^2 + x_2^3 + f''' = 0, 0) \subset (\mathbb{C}_{x_1, x_2, x_3, x_4}^4, 0).$$

Let us take a blow-up of $\mathbb{C}^4(\mathbf{p})$ along the curve C with weights $(1, 1, 1, 0)$. Then this blow-up induces $\psi: \tilde{X}(\mathbf{p}) \rightarrow X(\mathbf{p})$. Thus, $K_{\tilde{X}(\mathbf{p})} + S' = \psi^*K_{X(\mathbf{p})}$. It is obvious that f''' doesn't influence on the exceptional elliptic surface $S' \rightarrow \mathbb{P}^1$. It is easy to check that the degenerate cuspidal fibers lie over P_i . The surface S' has the singularities $\mathbb{E}_8$ in the degenerate fibers for $i = 1, 2$. Divisor $K_{X(\mathbf{p})} + S'$ is plt. Apply $K_{\tilde{X}(\mathbf{p})} + S' + \varepsilon E_{\tilde{X}(\mathbf{p})} - \text{MMP}$ ($0 < \varepsilon \ll 1$). Since $\rho(E_{\tilde{X}(\mathbf{p})}) = 1$ then on the first and unique step of MMP we have a divisorial contraction $E_{\tilde{X}(\mathbf{p})}$ to the point. So we get a required plt blow-up $\varphi: (Y, S) \rightarrow (X \ni 0)$. $\square$

3.6. Classification of three-dimensional exceptional quasihomogeneous canonical singularities. According to the proof of the main theorem it remains to classify the quasihomogeneous exceptional singularities. The classification process consists of two parts:

1. At first we have to classify the quasihomogeneous polynomials. For any such polynomial one can correspond its weight $\mathbf{p}$. By the main theorem the vector $\mathbf{p}$ is unique. It will be shown that there is the finite number of weights and consequently the finite number of types of exceptional singularities.
2. The second part is the investigation of given singularities on the exceptionality. Also the minimal complement index will be found.

Theorem 3.7. [21] *Let $(X, 0) \subset (\mathbb{C}^4, 0)$ be a canonical hypersurface singularity defined by a polynomial f and (X, H) is not log canonical for any hyperplane section H . Then f belongs to one of the following types:*

$$(\Upsilon_1) \quad f = t^2 + z^3 + zf_1(x, y) + f_2(x, y), \text{ where } \deg f_1 \geq 5, \deg f_2 \geq 7.$$

- (Υ_2) $f = t^2 + f'(z, x, y)$, where $\deg f' \geq 4$ and 4-jet is nonzero.
(Υ_3) $f = t^2 + f'(z, x, y)$, where $\deg f' \geq 5$ and 5-jet is nonzero.
(Υ_4) The rank of quadratic part of f is equal to 0 and 3-jet is nonzero.

The further notations. Further we always work under the following notations: $(X, 0) = (f(t, z, x, y) = 0, 0) \subset (\mathbb{C}_{t,z,x,y}^4, 0)$ denotes a canonical quasihomogeneous hypersurface singularity. Since one have to classify exceptional singularities we can assume that (X, H) is not log canonical, where H is any hyperplane section (see example 1.8).

The classification of the quasihomogeneous parts will be made by "Newton line rotation" [1, §11]. For this purpose we first find the finite number of the parts of type $\mathcal{M}_2$ (see definition 3.1). After that we obtain the required list of quasihomogeneous singularities with the help of the simple computer program. Therefore the main problem is to find the parts of type $\mathcal{M}_2$. At first consider the singularities of type Υ_1 .

Singularities of type Υ_1 . The following theorem describes such singularities.

Theorem 3.8. *All exceptional canonical quasihomogeneous singularities of type Υ_1 can be obtained by the rotations of plane in $\mathbb{C}_{z,x,y}^3$ passing through the monomial z^3 and one of the following monomials: $x^7, x^8, x^7y, x^9, x^8y, x^7y^2, x^{10}, x^9y, x^8y^2, x^7y^3, x^{11}, x^{10}y, x^9y^2, x^8y^3, x^7y^4$ and $zx^5, zx^5y, zx^7, zx^6y, zx^5y^2$. The rotation condition is that the weight of x is greater then the weight of y ($\text{wt } x \geq \text{wt } y$) and the point $2 = (2, 2, 2)$ lies over the rotation plane (see theorem 2.9).*

Proof. Note that the monomial zx^6 is missing in the list since the monomials z^3, zx^6, x^9 are lying on the straight line. One can choose the quasihomogeneous weights of f such that the quasihomogeneous degree is equal to 1.

In the theorem statement the monomials x^m of type $zx^a y^b, x^a y^b$ satisfy the following conditions: $a \geq b$; $\text{wt } x \geq \text{wt } y$; $\text{wt } x + \text{wt } y > \frac{1}{6}$ (see theorem 2.9); singularity defined by $t^2 + z^3 + x^m$ is not lc.

Let us prove that these monomials give all singularities. Assume the converse. Let the quasihomogeneous polynomial f gives an exceptional singularity $(X, 0)$ of type Υ_1 and every its monomial doesn't belong to the list (in this case $\text{wt } x > \text{wt } y$). Let $zx^{a_1} y^{b_1}$ be a monomial of minimal usual degree from $f_1(x, y)$ and $x^a y^b$ be a monomial of minimal usual degree from $f_2(x, y)$. Since $\text{wt } x > \text{wt } y$ it follows that the degree of x in these two monomials is maximal then the one in the remaining monomials of f . Also the degree of y is minimal.

If $f_1 = 0$ or $f_2 = 0$ then $f' = t^2 + z^3 + x^a y^b$ (or $+zx^{a_1} y^{b_1}$) defines an exceptional log canonical singularity. In fact there are two different

vectors from $N_{\mathbb{R}}$. Hence, f' gives an exceptional singularity (see main theorem proof). By taking $\mathbf{p} = (\frac{1}{2}, \frac{1}{3}, \frac{1}{a})$ (or $(\frac{2}{3a_1}, 1)$) we have $f_{\mathbf{p}} = f'$. Hence $(X, 0)$ is not exceptional.

Therefore there are the following 4 cases. We prove that they are impossible.

Case 1. $f' = t^2 + z^3 + x^a y^b$ and $f'' = t^2 + z^3 + zx^{a_1} y^{b_1}$ give the not log canonical singularities. Then they are not log canonical outside $\mathbf{0}$ along $\{t = z = y = 0\}$. Therefore $(X, 0)$ is not log canonical outside $\mathbf{0}$ along $\{t = z = y = 0\}$. Contradiction.

Case 2. $f' = t^2 + z^3 + x^a y^b$ gives a not log canonical singularity (i.e. $a < b \geq 7$) and $f'' = t^2 + z^3 + zx^{a_1} y^{b_1}$ gives a log canonical singularity (i.e. $a_1 \leq 4$ and $b_1 \leq 4$). Thus if $b_1 = 0$ then by the definition of Υ_1 we have $a_1 \geq 5$. Contradiction. We have the following linear equation system on the weights:

$$\begin{cases} a \text{ wt } x + b \text{ wt } y &= 1 \\ a_1 \text{ wt } x + b_1 \text{ wt } y &= 2/3, \quad d = ab_1 - ba_1 \neq 0 \end{cases}$$

Let $a_1 \geq b_1$. To prove theorem in this case it is enough to find new weights $\text{wt } x, \text{wt } y$ such that $f_{\mathbf{p}} = f''$, where $\mathbf{p} = (1/2, 1/3, \text{wt } x, \text{wt } y)$. Decrease $\text{wt } x$ by ε and increase $\text{wt } y$ by $\sigma = a_1 \varepsilon / b_1$ (after it the second system equation remains valid). These new weights are required. Actually we have to check that $\langle \mathbf{p}, x^a y^b \rangle > 1$, i.e. $-d = -ab_1 + ba_1 > 0$. It is true since $a/b < 1$ and $a_1/b_1 \geq 1$.

Let $a_1 < b_1$. If $-d > 0$ then $(X, 0)$ is also nonexceptional as above. Let $d > 0$ then $\text{wt } x = (b_1 - \frac{2}{3}b)/d > 0$. Hence $b_1 > \frac{2}{3}b \geq 14/3$. Contradiction.

Case 3. $f' = t^2 + z^3 + x^a y^b$ gives a log canonical singularity (i.e. $a \leq 6$ and $b \leq 6$) and $f'' = t^2 + z^3 + zx^{a_1} y^{b_1}$ gives a not log canonical singularity (i.e. $a_1 < b_1 \geq 5$). For the same reason $b \neq 0$. If $d > 0$ then as before in the case 2) we can find $\mathbf{p}$ such that $f_{\mathbf{p}} = f'$. If $d < 0$ then as before in the case 2) we have $b_1 - \frac{2}{3}b < 0$, i.e. $b > \frac{3}{2}b_1 \geq 15/2$. Contradiction.

Case 4. $f' = t^2 + z^3 + x^a y^b$ and $f'' = t^2 + z^3 + zx^{a_1} y^{b_1}$ give the log canonical singularities. Similarly let us decrease $\text{wt } x$ and increase $\text{wt } y$ in order that the first or second system equation remains valid. In one of the cases we get $f_{\mathbf{p}} = f'$ or $f_{\mathbf{p}} = f''$. $\square$

The investigation of singularities on the exceptionality. At first let us recall the main facts about the hypersurfaces in the weighted projective spaces [2]. The greatest common divisor of the numbers $a_1, \dots, a_n$ is denoted by $(a_1, \dots, a_n)$.

Proposition 3.9. *Let $(a_1, \dots, a_n) = 1$ and $q_i = (a_1, \dots, \hat{a}_i, \dots, a_n)$, where $a_i \in \mathbb{N}$ for all i . Then $\mathbb{P} = \mathbb{P}(a_1, \dots, a_n) \cong \mathbb{P}(a_1/q_i, \dots, a_i, \dots, a_n/q_i)$. If $q_i = 1$ for all i then $K_{\mathbb{P}} \sim \mathcal{O}_{\mathbb{P}}(-\sum a_i)$ and $\mathcal{O}_{\mathbb{P}}(1)^{n-1} = 1/(a_1 \cdots a_n)$. In this case the weights of $\mathbb{P}$ are called well-defined.*

Proposition 3.10. *In the same notations let $(X_d \subset \mathbb{P}) = (g(x_1, \dots, x_n) \subset \mathbb{P}(a_1, \dots, a_n))$ be a hypersurface of degree d . Then $(X_d \subset \mathbb{P}) = (g(x_1, \dots, x_i^{1/q_i}, \dots, x_n) \subset \mathbb{P}(a_1/q_i, \dots, a_i, \dots, a_n/q_i))$. If $\text{Diff}_{X_d/\mathbb{P}}(0) = 0$ and the weights of $\mathbb{P}$ are well-defined then $K_{X_d} = (K_{\mathbb{P}} + X_d)|_{X_d} = \mathcal{O}_{X_d}(d - \sum a_i)$.*

Definition 3.11. Assume the weights of the weighted projective space $\mathbb{P}$ are well-defined. Then the hypersurface $X_d \subset \mathbb{P}$ is called *well-formed* if one of the following equivalent conditions holds:

1. $\text{codim}_{X_d}(X_d \cap \text{Sing } \mathbb{P}) \geq 2$,
2. $\text{Diff}_{X_d/\mathbb{P}}(0) = 0$,
3. for all $i \neq j$ we have $(a_1, \dots, \hat{a}_i, \dots, \hat{a}_j, \dots, a_n) | d$.

3.12. The structure of purely log terminal blow-up of well-formed singularity. Let $\mathbf{p} = (a_1, \dots, a_n)$ be the weights of the quasihomogeneous log canonical singularity $g(x_1, \dots, x_n) = 0$ and $a_i \in \mathbb{N}$ for all i . By the main theorem these weights are defined uniquely for the exceptional singularity (in particular, $a_i > 0$ for all i). A $\mathbf{p}$ -blow-up of $\mathbb{C}^n$ induces a plt blow-up $(Y, E) \rightarrow (\{g = 0\}, 0)$. Given log Fano variety is $(E, \text{Diff}_E(0)) = (E, D) = (\tilde{g}(x_1, \dots, x_n) \subset \mathbb{P}(\tilde{a}_1, \dots, \tilde{a}_n), D) = (g(x_1^{1/q_1}, \dots, x_n^{1/q_n}) \subset \mathbb{P}(a_1 q_1 / (q_1 \cdots q_n), \dots, a_n q_n / (q_1 \cdots q_n)), \sum \frac{q_i - 1}{q_i} \{x_i = 0\})$. When calculating (E, D) it is important that the hypersurface $E \subset \mathbb{P}$ be well-formed. If it is not well-formed (the weights $\tilde{a}_1, \dots, \tilde{a}_n$ are well-defined) then we have other different D (see example 3.27).

By abuse of notation the coordinates of $g = 0$ and the coordinates of the weighted projective space are both denoted equally.

The quasihomogeneous degrees of $f, \tilde{f}$ are denoted by d and $\tilde{d}$ respectively. In our three-dimensional case given log Del Pezzo surface is always denoted by (E, D) .

The following easy proposition allows to consider only finite number of cases when studying given log Del Pezzo surface on the exceptionality.

Proposition 3.13. [24, 2.3] *Let (S, Δ) be a weakly log Del Pezzo surface, i.e. $-(K_S + \Delta)$ is nef and big. Assume that all 1,2,3,4 or 6-complements Δ^+ are exceptional, i.e. (S, Δ^+) is klt. Then log surface is exceptional.*

Using definition 1.3 we get the next corollary:

Corollary 3.14. *Notation is as in 3.13. Let $K_S + \frac{\lceil (n+1)\Delta \rceil}{n}$ be ample for all $n = 1, 2, 3, 4, 6$. Then (S, Δ) is exceptional.*

Corollary 3.15. *Let $(X, 0)$ be a three-dimensional quasihomogeneous log canonical singularity. Assume its any hyperplane section is not log canonical. Let $(E, D = \sum d_i D_i)$ be a log Del-Pezzo surface obtained by a plt blow-up. If $d_k \geq \frac{6}{7}$ for some k and $\text{Diff}_{E/\mathbb{P}}(0) = 0$ (it is the condition of well-formedness) then $(X, 0)$ is exceptional.*

Proof. Let $(X, 0)$ is not exceptional. Then by proposition 3.13 there exists 1,2,3,4 or 6-complement D^+ . By the definition of complement $D^+ \geq D_k + \sum_{i \neq k} d_i D_i$, where $D_k = \{x_k = 0\}$. Since $\text{Diff}_{E/\mathbb{P}}(0) = 0$ then $\frac{1}{q_1 \dots q_4} (d - \sum_{i \neq k} a_i) = \tilde{d} - \sum_{i \neq k} \frac{\tilde{a}_i}{q_i} \leq 0$. Therefore by theorem 2.9 the general hyperplane section $x_k + h(x_1, \dots, x_4) = 0$ of $(X, 0)$ is lc. This contradiction concludes the proof. $\square$

Remark 3.16. The corollary 3.15 is hypothetically true in any dimension n . In the statement we need to assume $d_k \geq \frac{|RN_{n-1}|}{|RN_{n-1}|+1}$, where $|RN_n| = \max\{m | m \in RN_n\}$. The set RN_n is defined in the following way: $m \in RN_n$ if there exists the nonexceptional log Fano variety of dimension n such that the minimal index of nonexceptional complements is equal to m . A hypothesis is that $|RN_n| < \infty$ for all n . For $n \leq 3$ it was proved in [24], besides $RN_1 = \{1, 2\}$ and $RN_2 = \{1, 2, 3, 4, 6\}$ [24].

The common principles of investigation on the exceptionality.

(1) At first one can check that $\text{Diff}_{E/\mathbb{P}}(0) = 0$ (it is convenient to verify the third condition in the definition 3.11).

(2) If D contains a component with coefficient $\geq \frac{6}{7}$ then by corollary 3.15 the singularity is not exceptional.

Any complement D^+ of (E, D) can be extended to the complement $K_{\mathbb{P}} + E + \widehat{D}^+ \sim_{\mathbb{Q}} 0$, where $\widehat{D}^+|_E = D^+$. Hence, the structure of Weyl divisor group of weighted projective space $\mathbb{P}$ implies that $\text{Supp } D^+$ is cut by polynomials.

For any number $n \in \mathbb{N}$ satisfying the following property there exists an n -complement.

Property: Let $r_n = \sum_{i=1}^4 \tilde{a}_i - \tilde{d} - \frac{1}{n} \sum_{i=1}^4 \lceil \frac{q_i-1}{q_i} (n+1) \rceil \tilde{a}_i$, i.e. $K_E + \frac{\lceil (n+1)D \rceil}{n} \sim_{\mathbb{Q}} -r_n \mathcal{O}_E(1)$. Then $r_n \geq 0$ and $r_n = \frac{1}{n} (\sum \tilde{a}_i b_i)$, where $b_i \in \mathbb{N}$ for all i .

(3) By corollary 3.13 there are two cases: (E, D) is exceptional and there is no 1,2,3,4 or 6-complement or there exists 1,2,3,4 or 6-complement. Later on we always assume the second case to take place.

First we determine the singular points of $\tilde{f} \subset \mathbb{P}$ and the singular points of curves from $\text{Supp } D$. It is very convenient to apply lemma 2.2 for these purposes. The answer depends on the coefficients of f .

(4) Let D^+ be a complement of (E, D) . By theorem 1.14 the question about the exceptionality is equivalent to the next one: Is pair (E, D^+) klt? This question is local. Consider this pair in the chart $U_k = \mathbb{C}^3 / \mathbb{Z}_{a_k} (a_1, \dots, a_{k-1}, a_{k+1}, \dots, a_4)$ of $\mathbb{P}$. The klt property of pair remains true by the finite dominant morphisms [10, 3.16]. Hence our investigation reduces to the finite number of questions: Is pair $(\mathbb{C}_{u,v}^2, B = \sum c_i \{h_i(u, v) = 0\})$ klt? Is pair $(\mathbb{C}_{u,v,w}^3, B = \{h = 0\} + \sum c_i \{h_i(u, v) = 0\})$ plt, where $h = 0$ defines Du Val singularity (h is a local equation of E in the corresponding chart)? In solving these problems it is convenient to consider the analog of inductive blow-up [12, 1.5] (it is a blow-up with the unique exceptional divisor C such that $(C, \text{Diff}_C(\tilde{B}))$ is klt or plt, where $\tilde{B}$ is a proper transform of B) and apply lemma 3.19.

The singularity investigation of type Υ_1 on the exceptional-ity.

Further we consider the most interesting cases and the common examples illustrating the investigation of given log Del Pezzo surfaces on the exceptionality.

Example 3.17. $\mathbf{f} = \mathbf{t}^2 + \mathbf{z}^3 + \mathbf{x}^{11} + \mathbf{y}^{13}$. Then $(E, D) = (\mathbb{P}^2, \frac{1}{2}l_1 + \frac{2}{3}l_2 + \frac{10}{11}l_3 + \frac{12}{13}l_4)$, where l_i are the lines in the general position. By corollary 3.14 or by corollary 3.15 given log surface is exceptional. A minimal complement index is equal to 66. The corresponding complement is $D^+ = \frac{1}{2}l_1 + \frac{2}{3}l_2 + \frac{10}{11}l_3 + \frac{61}{66}l_4$. A conjecture is that 66 is maximal number for log surfaces with the standard coefficients and consequently for every three-dimensional local contraction (singularity) [24]. In other words $|RN_3| = 66$ (see remark 3.16).

Example 3.18. $\mathbf{f} = \mathbf{t}^2 + \mathbf{z}^3 + \mathbf{g}(\mathbf{z}, \mathbf{x}, \mathbf{y}) = \mathbf{t}^2 + \mathbf{z}^3 + \mathbf{z}\mathbf{y}\mathbf{f}'_2(\mathbf{x}^2, \mathbf{y}^3) + \mathbf{x}\mathbf{f}'_3(\mathbf{x}^2, \mathbf{y}^3)$. The lower indexes denote the binary form degrees. These forms are different although they are both denoted by $\mathbf{f}'$. Let us remark that we got rid of the monomial z^2xy^2 before writing f . Namely we

made the coordinate change $z \mapsto z + \alpha xy^2$ for some α . The rotation takes place through the unique part $t^2 + z^3 + x^7$ of type $\mathcal{M}_2$. Therefore the coefficient of x^7 is not equal 0. Put $f'_2 = ax^{2i}f_{2-i}(x^2, y^3)$, $f'_3 = x^{2j}f_{3-j}(x^2, y^3)$.

Proposition. *$ij = 0$ (if $a = 0$ then $i \neq 0$ by definition) if and only if $(X, 0)$ is exceptional and 2-complementary.*

Proof. A plt blow-up with weights $(21, 14, 6, 4)$ gives $(E, D) = (t + z^3 + g(z, x, y) \subset \mathbb{P}(21, 7, 3, 2), \frac{1}{2}\{t = 0\}) = (\mathbb{P}(7, 3, 2), \frac{1}{2}\{t = 0\})$. Let $i \geq 1$ and $j \geq 1$. Then $K_E + \frac{1}{2}\{t = 0\} + \frac{1}{2}\{x = 0\} \equiv 0$ is not klt hence singularity is not exceptional. Check it. Let $i = j = 1$. Then this statement is equivalent to the following one: $(\mathbb{C}_{u,v}^2, \frac{1}{2}\{(u^3 + a'u^2v + v^3) \cdot v = 0\})$ is not klt. Indeed, take a blow-up with weights $(1, 1)$. Then the exceptional divisor with discrepancy -1 appears. The cases with other i, j are considered similarly.

Now, let $i = 0$. Consider two more difficult cases:

1. $g(z, x, y) = zy^7 + x^7$
2. $g(z, x, y) = zy(x^2 + y^3)^2 + x(x^2 + y^3)^3$

The first case is exceptional obviously. In the second case it is enough to check that $K_E + \frac{1}{2}\{t = 0\} + \frac{1}{4}\{x^2 + y^3 = 0\} \equiv 0$ is klt. This statement is equivalent to $(\mathbb{C}_{u,v}^2, B)$ be klt, where $B = \frac{1}{2}\{u^3 + a'u^2v + v^3 = 0\} + \frac{1}{4}\{v = 0\}$. To prove this or analogous assertion the next easy lemma can be applied.

Lemma 3.19. *Let $(Z \ni P, D)$ be a log pair and let $f: Y \rightarrow Z$ be a birational morphism. Assume that the exceptional set of f consists of one irreducible exceptional divisor C . If $a(C, D) > -1$ (resp. $a(C, D) = -1$) and $K_Y + C + D_Y$ is plt (D_Y is a proper transform of D), then $(Z \ni P, D)$ is klt (resp. lc) and exceptional.*

Take the blow-up with the weights $(1, 1)$. Then the single exceptional divisor $C \cong \mathbb{P}^1$ with discrepancy $-3/4$ appears. The divisor $K_C + \text{Diff}_C(\tilde{B}) = K_C + \frac{1}{2}P_1 + \frac{1}{2}P_2 + \frac{1}{2}P_3 + \frac{1}{4}P_4$ is klt, where P_i are the distinct points and $\tilde{B}$ is a proper transform of B . The case $j = 0$ is considered similarly. Divisor $D^+ = \frac{1}{2}\{t = 0\} + \frac{1}{2}\{x = 0\}$ is 2-complement. $\square$

Consider quotient singularity $\mathbb{C}^3/H_{168}$, where H_{168} is a simple Klein's group of order 168. The singularity pointed out has the same type as in the previous example. It is exceptional (cf. [19]). Note also that it is a degenerate singularity.

Example 3.20. $\mathbf{f} = \mathbf{t}^2 + \mathbf{z}^3 + \underline{\mathbf{a}}\mathbf{z}\mathbf{x}^i\mathbf{f}'_{5-i}(\mathbf{x}, \mathbf{y}^2) + \underline{\mathbf{b}}\mathbf{y}\mathbf{x}^j\mathbf{f}'_{7-j}(\mathbf{x}, \mathbf{y}^2)$. The plane can rotate through two different parts of type $\mathcal{M}_2$ for this singularity. The first part is $t^2 + z^3 + x^7y$. The second one is $t^2 + z^3 + zx^5$. Therefore the coefficient of zx^5 or the coefficient of x^7y is not equal 0. On occasion of notation of $\mathbf{f}$ see the conventions in §4. A plt blow-up with weights $(15, 10, 4, 2)$ gives $(E, D) = (t + z^3 + \dots \subset \mathbb{P}(15, 5, 2, 1), \frac{1}{2}\{t = 0\}) = (\mathbb{P}(5, 2, 1), \frac{1}{2}\{t = 0\})$. Let $i \geq 3$ and $j \geq 5$. Then $K_E + \frac{1}{2}\{t = 0\} + \frac{1}{4}\{x = 0\} \equiv 0$ is not klt. Hence singularity is not exceptional. Indeed, consider (E, D) in the neighborhood of the point $(0 : 0 : 0 : 1)$. Then we have to show that $(\mathbb{C}^2, \frac{1}{2}\{u^3 + \underline{a}uv^i + \underline{b}v^j = 0\} + \frac{1}{4}\{v = 0\})$ is not klt. Take a blow-up with weights $(3, 2)$. The exceptional divisor with discrepancy -1 appears. Q.E.D.

If $i \leq 2$ or $j \leq 4$ then singularity is not exceptional and 2-complementary. The requirement $i \leq 2 \parallel j \leq 4$ also means that it is true for the multiplicities of other common components of type $x + cy^2$. We write this requirement only for x , because after making the quasi-homogeneous coordinate change $x \mapsto x' - cy^2$ we have the initial case. Consider two more difficult cases. Let $j = 4$ and $\underline{a} = 0$. It is enough to check that $(\mathbb{C}^2, B = \frac{1}{2}\{u^3 + v^4 = 0\} + \frac{1}{4}\{v = 0\})$ is klt. Take a blow-up with weights $(4, 3)$. The single exceptional divisor C with discrepancy $-3/4$ appears. The pair $(C, \text{Diff}_C(\tilde{B})) = (\mathbb{P}^1, \frac{1}{2}P_1 + \frac{7}{8}P_2 + \frac{2}{3}P_3)$ is klt, where P_i are the distinct points. Lemma 3.19 completes the proof. The remained difficult case $i = 2$ and $\underline{b} = 0$ is studied similarly.

Singularities of type Υ_2 . Let us consider the exceptional singularities of this type. By corollary 2.19 there is a biholomorphic map such that 4-jets can be only the following: z^4 (type $\Upsilon_2^{(1)}$), z^3x (type $\Upsilon_2^{(2)}$), z^3y (type $\Upsilon_2^{(3)}$).

Singularities of type $\Upsilon_2^{(1)}$ $(t^2 + z^4 + z^2f_1(x, y) + zf_2(x, y) + f_3(x, y))$. To rotate Newton line (the rotation condition is $\text{wt } x \geq \text{wt } y$) it is enough to fix the monomials of the following forms: x^ay^b and zx^ay^b . Using the same arguments just as for singularities of type Υ_1 (theorem 3.8) we get the following cases to rotate Newton line: $x^5, x^6, x^5y, x^7, x^6y, x^5y^2$ and zx^4, zx^5, zx^4y .

Example 3.21. $\mathbf{f} = \mathbf{t}^2 + \mathbf{z}^4 + \mathbf{z}\mathbf{x}^4\mathbf{y} + \mathbf{x}^3\mathbf{y}^{\mathbf{n}}, \mathbf{n} = 4, 5$. The weights of plt blow-up are $(2(4n - 3), (4n - 3), 3n - 4, 7)$. Besides $(E, D) = (f \subset \mathbb{P}(2(4n - 3), (4n - 3), 3n - 4, 7), 0)$ and $K_E = (n - 6)\mathcal{O}_E(1)$. If $n = 5$ then there is no 1, 2, 3, 4 and 6-complement. Therefore the singularity is exceptional and $\frac{1}{7}\{y = 0\}$ gives 7-complement of minimal index.

Let $n = 4$. There is 4-complement $D^+ = \frac{1}{4}\{x = 0\}$. Let us prove that (E, D^+) is not plt. For this purpose it is enough to prove that $(\mathbb{C}_{u,v,w}^3, \{u^2 + v^4 + w^3 = 0\} + \frac{1}{4}\{w = 0\})$ is not plt. Take the weighted blow-up with weights $(6, 3, 4)$. Then exceptional divisor with discrepancy -1 appears. Thus the singularity is not exceptional.

Singularities of type $\Upsilon_2^{(2)}(t^2 + z^3x + g(z, x, y))$. Using the same arguments just as for singularities of type Υ_1 (see theorem 3.8) we get the following cases to rotate Newton line ($\text{wt } x \geq \text{wt } y$): $x^5, x^6, x^5y, x^7, x^6y, x^5y^2, x^8, x^7y, x^6y^2, x^5y^3, x^9, x^8y, x^7y^2, x^6y^3, x^5y^4$ and $zx^4, zx^4y, zx^6, zx^5y, zx^4y^2$. Note that zx^5 is a subcase in x^7 case.

Example 3.22. $\mathbf{f} = \mathbf{t}^2 + \mathbf{z}^3\mathbf{x} + \mathbf{az}^2\mathbf{yx}^{i_1}(\mathbf{x} + \mathbf{y}^3)^{i_2}\mathbf{f}_{2-i_1-i_2}(\mathbf{x}, \mathbf{y}^3) + \mathbf{bzy}^2\mathbf{x}^{j_1}(\mathbf{x} + \mathbf{y}^3)^{j_2}\mathbf{f}_{3-j_1-j_2}(\mathbf{x}, \mathbf{y}^3) + \mathbf{x}^{l_1}(\mathbf{x} + \mathbf{y}^3)^{l_2}\mathbf{f}_{5-l_1-l_2}(\mathbf{x}, \mathbf{y}^3)$. The lower indexes denote the binary form degrees. After the coordinate change $x \mapsto x + \alpha y^3$ for some α we got rid of the monomial z^3y^3 . The rotation takes place through the unique part $t^2 + z^3x + x^5$ of type $\mathcal{M}_2$. Therefore the coefficient of x^5 is not equal 0. Since variable x is present in the monomial z^3x then we must select other common component $x + y^3$. A plt blow-up is the blow-up with weights $(15, 8, 6, 2)$. Thus $(E, D) = (\mathbb{P}(4, 3, 1), \frac{1}{2}\{t = 0\})$. There are the following 1, 2, 3, 4 and 6-complements: 2-complement $-D + \frac{1}{2}\{y = 0\}$ and 6-complement $-D + \frac{1}{6}\{x + \beta y^3 = 0\}$. Hence, the minimal complement index is equal to 2. Since the curve $\{y = 0\}$ doesn't pass through the singular points of $\text{Supp } D$ then the first complement is not considered. Let us consider the remaining cases $\beta = 0, \beta = 1$.

Let $\beta = 0$. If $i_1 \geq 2$ ($i_1 > 2$ denotes that $a = 0$), $j_1 \geq 3$ ($j_1 > 3$ denotes that $b = 0$), $l_1 = 3$ (if $l_1 \geq 4$ then our singularity is not canonical) then $(\mathbb{C}_{u,v}^2, B)$ is not klt, where $B = \frac{1}{2}\{u^3v + v^3 + au^2v^2 + buv^3 = 0\} + \frac{1}{6}\{v = 0\}$. Actually take a blow-up with weights $(2, 3)$. Then the exceptional divisor with discrepancy -1 appears. In other cases its complement is not exceptional. For example, let us prove it if $i_1 \geq 2, j_1 = 2, l_1 \geq 3$. In this case $B = \frac{1}{2}\{u^3v + au^2v^2 + uv^2 + v^{l_3} = 0\} + \frac{1}{6}\{v = 0\}$. Take a blow-up with weights $(1, 2)$. The discrepancy of the single exceptional divisor C is equal to $-\frac{5}{6}$. Besides $\text{Diff}_C(\tilde{B}) = \frac{3}{4}P_1 + \frac{2}{3}P_2 + \frac{1}{2}P_3$, where $\tilde{B}$ is a proper transform of B . Hence $K_C + \text{Diff}_C(\tilde{B})$ is klt, i.e. by lemma 3.19 pair $(\mathbb{C}^2, B)$ is klt.

Let $\beta = 1$. Then $B = \frac{1}{2}\{h = 0\} + \frac{1}{6}\{v = 0\}$, where $h(u, v) = u^3 + au^2v^{i_2} + buv^{j_2} + v^{l_2}$. It is easy to verify that $(\mathbb{C}^2, B)$ is klt if $b \neq 0$. Let $b = 0$. Similarly one can verify that $(\mathbb{C}^2, B)$ is always klt if $i_2 \leq 1$. Let $i_2 \geq 2$. If $l_2 \leq 4$ then this pair is klt. If $l_2 = 5$ then it is not klt.

Summary. The canonical singularity is exceptional and 2-complementary except the case $i_1 \geq 2$ & $j_1 \geq 3$ & $l_1 = 3$ and the case $i_2 \geq 2$ & $b = 0$ & $l_1 = 3$. These conditions are written taking into account other common components (see example 3.20).

Singularities of type $\Upsilon_2^{(3)}$. The following theorem takes place.

Theorem 3.23. *All exceptional quasihomogeneous singularities of this type can be obtained by the rotations of plane in $\mathbb{C}_{z,x,y}^3$ passing through the monomial z^3y and one of the following monomials: $x^7, x^8, x^7y, x^9, x^8y, x^7y^2$ and $zx^5, zx^6, zx^5y, z^2x^3$. The rotation condition is that the weight of x is greater than the weight of y ($\text{wt } x > \text{wt } y$) and the point $\mathbf{2} = (2, 2, 2)$ lies over the rotation plane.*

Proof. Let $f = t^2 + z^3y + z^2f_1(x, y) + zf_2(x, y) + z^2f_3(x, y)$. Denote the monomial of usual minimal degree from $z^2f_1(x, y)$ by $z^2x^ay^b$. If $b \neq 0$ then by taking the quasihomogeneous coordinate change $z \mapsto z - \alpha f_1(x, y)y^{-1}$ one can assume that $f_1(x, y) = 0$. Applying the same arguments as in the theorem 3.8 we complete the proof. Let $b = 0$. Then using $\text{wt } x + \text{wt } y + \text{wt } z > 1/2$ we get $a \leq 3$. Corollary 2.19 implies that $a = 3$. Q.E.D. $\square$

The singularities of type Υ_3 . For the exceptional singularities of this type there are three cases for 5-jets.

THE FIRST CASE. 5-jet is a point (type $\mathcal{M}_0$), i.e. $f_5 = z^ax^by^c$, where $a \geq b, a \geq c$ and $a + b + c = 5$. By corollary 2.19 we have $a \geq 3$. Thus 5-jet is one of the following: z^5 (type $\Upsilon_3^{(1)}$), z^4x (type $\Upsilon_3^{(2)}$), z^4y (type $\Upsilon_3^{(3)}$), z^3x^2 (type $\Upsilon_3^{(4)}$), z^3y^2 (type $\Upsilon_3^{(5)}$), z^3xy (type $\Upsilon_3^{(6)}$). As in the theorems 3.8, 3.23 there are the following cases to rotate Newton line ($\text{wt } x \geq \text{wt } y$):

Singularities of type $\Upsilon_3^{(1)}$. The rotation monomials are x^6, x^5y, x^4y^2 and zx^5, zx^4y, zx^3y^2 .

Singularities of type $\Upsilon_3^{(2)}$. The rotation monomials are x^6, x^5y, x^4y^2 and zx^5, zx^4y, zx^3y^2 .

Singularities of type $\Upsilon_3^{(3)}$. The rotation monomials ($\text{wt } x > \text{wt } y$) are x^6, x^5y and zx^5, zx^4y . All singularities in the case zx^4y are not well-formed.

Singularities of type $\Upsilon_3^{(4)}$. The rotation monomials are $x^6, x^5y, x^4y^2, x^3y^3, x^7, x^6y, x^5y^2, x^4y^3, x^3y^4$ and zx^5, zx^4y, zx^3y^2 .

Singularities of type $\Upsilon_3^{(5)}$. The rotation monomials ($\text{wt } x > \text{wt } y$) are x^7 and zx^5 .

Singularities of type $\Upsilon_3^{(6)}$. The rotation monomials are $x^5, x^6, x^5y, x^7, x^6y, x^5y^2$ and zx^5, zx^4y . All singularities in the cases x^5y, x^6y, zx^4y are not well-formed.

THE SECOND CASE. 5-jet is a line, i.e. $f_5 \in \mathcal{M}_1$ (type $\Upsilon_3^{(7)}$). To rotate Newton line ($\text{wt } x \geq \text{wt } y$) we have to fix the following parts: $g_5(z, x)$ is a binary form of degree 5, $z^5 + zx^3y, z^5 + x^4y, z^4x + zx^3y, z^4x + x^4y, z^4y + z^2x^3, z^4y + zx^4, z^3x^2 + zx^3y, z^3x^2 + x^3y^2, z^3y^2 + z^2x^3$. This list was obtained in the following way. The first monomial x^{m_1} contains a variable of maximal usual degree among other monomials. This variable is denoted by z . The second monomial $x^{m_2} = z^a x^b y^c$ satisfies the following conditions: $b \geq c$ and $t^2 + x^{m_1} + x^{m_2}$ gives not log canonical outside $\mathbf{0}$ singularity.

To prove that these cases describe all possibilities one may use the theorem 3.8 proof.

Consider one-dimensional 5-jet $f_5 = z^5 + \alpha z^3xy + \beta zx^2y^2$. Let us prove that this case doesn't realize. If $\beta \neq 0$ then $\exists \mathbf{p} \in N_{\mathbb{R}}$ such that $f_{\mathbf{p}} = t^2 + f_5$ is a log canonical outside $\mathbf{0}$ singularity. Therefore by corollary 2.19 our singularity is not exceptional. Actually let $x^n = z^c x^a y^b$ ($c + a + b \geq 6$) be another monomial and $d = b - a$ be a determinant of the following linear equation system on the weights.

$$\begin{cases} \text{wt } x + \text{wt } y &= 2/5 \\ a \text{ wt } x + b \text{ wt } y &= 1 - (1/5)c \end{cases}$$

Let $d < 0$. Since $\text{wt } x \geq \text{wt } y$ then $c + a + b \leq 5$. Contradiction. Let $d > 0$. Decrease $\text{wt } x$ on ε and increase $\text{wt } y$ on ε . Then for new vector $\mathbf{p}$ we have $\langle \mathbf{p}, x^n \rangle > 1$. Q.E.D.

Let $\beta = 0$ and $x^n = z^c x^a y^b$ ($c + a + b \geq 6$) be another monomial. As above we have $d > 0$. Using conditions $\text{wt } x > 0$ and $\text{wt } y > 0$ one have $b \geq 4$ for $c = 0, 1$ and one have $b \geq 3$ for $c = 2, 3$. That is, $t^2 + z^3xy + x^n$ gives a not log canonical outside $\mathbf{0}$ singularity along $\{t = z = x = 0\}$. Hence the original singularity is not log canonical.

Other cases are considered similarly except the case $f_5 = z^4y + z^3x^2$. In this case the reader will have no difficulty in proving by exhaustion that only two new singularities $t^2 + z^4y + z^3x^2 + xy^5$ and $t^2 + z^4y + z^3x^2 + az^2y^5 + bx^2y^4$ ($b \neq 0$) appear. It is easy to check that they are not exceptional.

THIRD CASE. 5-jet is a plane, i.e. $f_5 \in \mathcal{M}_2$ (type $\Upsilon_3^{(8)}$). The classification of such singularities reduces to the description of all curves of degree 5 in $\mathbb{P}^2$.

Theorem 3.24. *Let $f = t^2 + f_5(z, x, y)$ gives a canonical singularity $(X, 0)$, where f_5 is a homogeneous polynomial of degree 5. Then*

$\mathbf{p} = (5, 2, 2, 2)$ -blow-up induces a plt blow-up of $(X, 0)$. Besides $(E, D) = (t + f_5(z, x, y) \subset \mathbb{P}(5, 1, 1, 1), \frac{1}{2}\{t = 0\}) = (\mathbb{P}^2, \frac{1}{2}C)$, where $C = \{f_5 = 0\}$. Then $(X, 0)$ is exceptional if and only if one of the following possibilities holds.

1. The curve $C \subset \mathbb{P}^2$ is irreducible. Its singular points can be only ordinary cusps and ordinary double points.
2. The curve $C \subset \mathbb{P}^2$ is the union of irreducible cubic and irreducible conic which intersect in 6 distinct points. The singular point of cubic (if it exists) can be only ordinary cusp or ordinary double point.

Proof. Since $(X, 0)$ is normal singularity then each irreducible component of the curve $C = \{f_5 = 0\}$ has the multiplicity 1. Hence we have three cases: C is an irreducible curve, C is an union of irreducible cubic and irreducible conic, C is an union of line and quartic. The corresponding log Del Pezzo surface is $(E, D) = (\mathbb{P}^2, \frac{1}{2}C)$.

Consider the first case. Let the singularity of curve be worse then ordinary cusp. The reader will easily prove that $K_{\mathbb{P}^2} + \frac{1}{2}C + \frac{1}{2}l \sim_{\mathbb{Q}} 0$ is not klt, where the straight line l passes through the singularity (in the cases of singularities $(u^2 + v^n = 0, 0) \subset (\mathbb{C}^2, 0)$, where $n = 4, 5$ we take $l = \{u = 0\}$).

Consider the second case. If the cubic and conic intersect with multiplicity 2 in the point P then the same divisor is not klt, where the straight line l is a tangent of conic in the point P .

Consider the third case. If l is a straight line of $\text{Supp } C$ then the same divisor is not klt. $\square$

Example 3.25. $\mathbf{f} = \mathbf{t}^2 + \mathbf{z}^3\mathbf{xy} + \mathbf{x}^i(\mathbf{x} + \mathbf{y}^2)^j\mathbf{f}_{5-i-j}(\mathbf{x}, \mathbf{y}^2) = \mathbf{t}^2 + \mathbf{z}^3\mathbf{xy} + \mathbf{h}(\mathbf{x}, \mathbf{y})$. The rotation takes place through the unique part $t^2 + z^3xy + x^5$. Therefore the coefficient of x^5 is nonzero. A plt blow-up is the blow-up with weights $(15, 7, 6, 3)$. Then $(E, D) = (t^2 + zxy + h(x, y) \subset \mathbb{P}(5, 7, 2, 1), \frac{2}{3}\{z = 0\})$. There are the following 1, 2, 3, 4 and 6-complements: 3-complement $D + \frac{1}{3}\{y = 0\}$ and 6-complement $D + \frac{1}{6}\{x + \beta y^2 = 0\}$. If $i \geq 1$ then 6-complement for $\beta = 0$ is 3-complement $D + \frac{1}{3}C$, where $C = \{t = x = 0\}$. The minimal complement index is equal to 3. As in example 3.22 one can consider only the second complement.

Let $\beta = 0$. If $i = 3$ (if $i_1 \geq 4$ then the singularity is not canonical) then the singularity is not exceptional. Indeed, we have to show that $(\mathbb{C}_{u,v,w}^3, B)$ is not plt, where $B = \{u^2 + vw + w^3 = 0\} + \frac{2}{3}\{v = 0\} + \frac{1}{6}\{w = 0\}$. Take a blow-up with weights $(3, 4, 2)$. Then the exceptional divisor with discrepancy -1 appears. If $i \leq 2$ then $(E, D + \frac{1}{6}\{x = 0\})$ is klt.

For example, prove it for $i = 2$. We claim that pair (Z, B) is klt, where $Z = \{u^2 + vw + w^2 = 0\} \subset (\mathbb{C}^3, 0)$ and $B = \frac{2}{3}\{v = 0\}|_Z + \frac{1}{6}\{w = 0\}|_Z$. Take a blow-up with weights $(1, 1, 1)$. Then the single exceptional divisor C with the discrepancy $a(C, B) = -\frac{5}{6}$ appears. Also $\text{Diff}_C(\tilde{B}) = \frac{2}{3}P_1 + \frac{2}{3}P_2 + \frac{1}{3}P_3$. Hence $K_C + \text{Diff}_C(\tilde{B})$ is klt. Q.E.D.

Let $\beta = 1$. For the same reason the singularity is exceptional if and only if $j \leq 4$.

Answer. Canonical singularity is exceptional and 3-complementary if and only if $i \leq 2$ and $j \leq 4$.

Singularities of type Υ_4 . These singularities are considered likewise. To rotate Newton line ($\text{wt } x \geq \text{wt } y$) there are four cases. For underlined parts all singularities are not well-formed or they belong to other cases.

(A) $f_3 \in \mathcal{M}_0$. Then f_3 is equal to one of the following monomials: t^3, t^2z, t^2x, t^2y . Let us consider every monomial separately.

1. After coordinate change we get $f = t^3 + tg(z, x, y) + h(z, x, y)$. Since $\mathbf{1} \in \Gamma_+(f)^0$ then $f_4 \neq 0$. We have the following cases — $z^2x^2 + x^5, z^2x^2 + x^4y, z^2x^2 + x^3y^2, \underline{z^2x^2 + x^2y^3}, z^2xy + x^5, \underline{z^2xy + x^4y}, f_4 = \varphi_4(z, x)$ is a binary form of degree 4, $z^4 + zx^2y, z^4 + x^3y, z^4 + tx^3, z^4 + tx^2y, z^3x + zx^2y, z^3x + x^3y, z^3x + tx^3, z^3x + tx^2y, z^3y + z^2x^2, z^3y + tx^3, \underline{tz^3 + z^2x^2}, \underline{tz^3 + zx^3}, tz^3 + zx^2y, \underline{tz^3 + x^3y}, \underline{z^2x^2 + azx^2y + bx^2y^2}, z^2x^2 + tz^2x, z^2x^2 + tx^2y, \underline{tz^2x + zx^2y}, \underline{tz^2x + x^2y^2}, f_4 \in \mathcal{M}_2$. The exceptional singularities of last case are classified in the theorem 3.28.
2. After coordinate change we get $f = t^2z + tg(x, y) + h(z, x, y)$. Then also $f_4 \neq 0$. We have the following cases — $f_4 = \varphi_4(z, x)$ is a binary form of degree 4, $z^4 + x^3y, z^4 + x^5, \underline{z^4 + x^4y}, z^4 + x^3y^2, z^4 + tx^3, z^4 + tx^2y, z^3x + x^3y, z^3x + x^5, z^3x + x^4y, z^3x + x^3y^2, z^3x + tx^3, z^3x + tx^2y, z^3y + x^5, \underline{z^3y + x^4y}, \underline{z^3y + zx^3}, z^3y + z^2x^2, z^3y + tx^3, z^2x^2 + x^5, z^2x^2 + x^4y, \underline{z^2x^2 + x^3y^2}, \underline{z^2x^2 + tx^2y}, z^2y^2 + x^5, z^2xy + x^5, \underline{z^2xy + x^4y}$.
3. After coordinate change we get $f = t^2x + tg(z, y) + h(z, x, y)$. Then also $f_4 \neq 0$. We have the following cases — $z^4 + x^3y, z^4 + x^5, z^4 + x^4y, z^4 + x^3y^2, z^4 + zx^2y, z^4 + zx^4, z^4 + zx^3y, z^4 + zx^2y^2, \underline{z^3x + x^2y^2}, \underline{z^3x + x^5}, \underline{z^3x + x^4y}, \underline{z^3x + x^3y^2}, \underline{z^3x + x^2y^3}, \underline{z^3x + x^6}, \underline{z^3x + x^5y}, \underline{z^3x + x^4y^2}, \underline{z^3x + x^3y^3}, \underline{z^3x + x^2y^4}, \underline{z^3x + zx^2y}, \underline{z^3x + zx^4}, \underline{z^3x + zx^3y}, \underline{z^3x + zx^2y^2}, z^3y + x^5, z^3y + x^4y, z^3y + x^6, \underline{z^3y + x^5y}, \underline{z^3y + x^4y^2}, \underline{z^3y + zx^4}, \underline{z^3y + zx^3y}, z^3y + z^2x^2$.

4. After coordinate change we get $f = t^2y + tg(z, x) + h(z, x, y)$. Then also $f_4 \neq 0$. We have the following cases — $z^4 + x^5$, $z^4 + zx^4$; $z^3x + x^5$, $z^3x + x^6$, $\underline{z^3x + x^5y}$, $z^3x + zx^4$, $z^3x + tx^3$.

(B) $f_3 \in \mathcal{M}_1$. Then f_3 is one of the following polynomials:

1. $f_3 = g_3(t, z)$. The rotation monomials are x^4, x^5, x^4y, zx^3 .
2. $f_3 = t^3 + z^2x$. The rotation monomials are $x^4, x^5, x^4y, x^6, x^5y, x^4y^2, x^7, x^6y, x^5y^2, x^4y^3, x^8, x^7y, x^6y^2, x^5y^3, x^4y^4, tx^3, tx^3y, tx^5, tx^4y, tx^3y^2$. The monomials zx^3, zx^4, zx^3y, tx^4 are the subcases of monomials x^5, x^7, x^5y^2, x^6 respectively.
3. $f_3 = t^3 + z^2y$. The rotation monomials are $x^7, x^8, \underline{x^7y}, zx^4, tx^5$.
4. $f_3 = t^2z + z^2x$. The rotation monomials are $x^4, x^5, x^4y, x^6, x^5y, x^4y^2, tx^3, tx^4, tx^3y$. The monomial zx^3 is a subcase of monomial x^5 .
5. $f_3 = t^2z + z^2y$. The rotation monomials are $x^5, x^6, \underline{x^5y}, \underline{zx^3}, tx^4$.

(C) $f_3 \in \mathcal{M}_2$. Then after coordinate change f_3 belongs to one of the followings types: case (A), case (B) or a log canonical threshold $c(f_3) = 1$. In the last case by corollaries 2.18, 2.19 $(X, 0)$ is not exceptional.

(D) $f_3 \in \mathcal{M}_3$. Then after coordinate change f_3 belongs to one of the followings types: case (A), case (B), case(C) or the general hyperplane section has log canonical singularities, i.e. $(X, 0)$ is not exceptional.

Example 3.26. $\mathbf{f}$ has a type $\mathbf{f}_3(\mathbf{t}, \mathbf{z}) + \mathbf{y}^2\mathbf{f}_2(\mathbf{t}, \mathbf{z}) + (\mathbf{y}^4 + \mathbf{x}^2\mathbf{y})\mathbf{f}_1(\mathbf{t}, \mathbf{z}) + \mathbf{f}_2(\mathbf{x}^2, \mathbf{y}^3)$. In this case we must assume that the binary form $f_3(t, z)$ doesn't have the multiple irreducible factors. This singularity is obtained by Newton line rotation for the case $f_3(t, z) + x^4 \in \mathcal{M}_2$. After some quasihomogeneous coordinate change (see conventions in §4) we get $f = t^3 + ktz^2 + lz^3 + t(ax^2y + by^4) + z(cx^2y + dy^4) + etzy^2 + x^{2i}f_{2-i}(x^2, y^3)$. Besides $|k| + |l| \neq 0$. A plt blow-up is the blow-up with weights $(4, 4, 3, 2)$. Then $(E, D) = (f(t, z, x^{1/2}, y) \subset \mathbb{P}(2, 2, 3, 1), \frac{1}{2}\{x = 0\})$. There are the following 1,2,3,4,6 complements: $D + \frac{1}{2}\{y = 0\}$; $\frac{2}{3}\{x = 0\}$; $D + \frac{1}{4}\{\alpha t + \beta z + \gamma y^2 = 0\}$; $D + \frac{1}{6}\{y = 0\} + \frac{1}{6}\{\alpha t + \beta z = 0\}$. Since the coefficient of x^4 is nonzero then it is enough to consider the second and the third complement ($\gamma = 0$).

Let $e \neq 0$. Assume that $l = 0 (k \neq 0)$ & $z(cx^2y + dy^4) = 0$ & $i = 2$ then $D^+ = \frac{1}{4}\{t = 0\} + \frac{1}{2}\{x = 0\}$ is non-klt 4-complement.

Let $e = 0$. If $|b| + |d| = 0$ & $i \geq 1$ then $D^+ = \frac{2}{3}\{x = 0\}$ is non-klt 4-complement. For $e \neq 0$ the previous condition on the exceptionality remains only necessary one. Moreover $D^+ = \frac{1}{4}\{t + \alpha z = 0\} + \frac{1}{2}\{x = 0\}$ is non-klt 4-complement if $i = 2$ & $\text{G.C.D}(t^3 + ktz^2 + lz^3, t(ax^2y + by^4) + z(cx^2y + dy^4)) = t + \alpha z$.

Example 3.27. $\mathbf{f} = \mathbf{t}^2\mathbf{x} + \mathbf{z}^3\mathbf{y} + \mathbf{x}^5 + \mathbf{zxy}^3$. A plt blow-up is the blow-up with weights $(16, 11, 8, 7)$. Then $(E, D) = (f(t, z, x, y) \subset \mathbb{P}(16, 11, 8, 7), 0)$. Let us consider the neighborhood of the point $(0 : 0 : 0 : 1)$ in some chart. Then $(\mathbb{C}^3, S + \frac{1}{4}T) = (\mathbb{C}_{u,v,w}^3, \{u^2w + v^3 + w^5 + vw = 0\} + \frac{1}{4}\{w = 0\})$ is not plt. Indeed, take a blow-up ψ with the weights $(1, 2, 4)$. Hence the singularity is not exceptional. The singularities of such kind have the complex structure. For example, in our case $(S, 0)$ is Du Val point of type $\mathbb{A}_5$. So to prove the exceptionality we have to consider the inductive blow-up.

In the capacity of the example let us show that $(C, B) = (C, \text{Diff}_C(\frac{1}{4}\tilde{T}'))$ is klt in our case, where C is an exceptional curve of blow-up $\psi|_{\tilde{S}}: \tilde{S} \rightarrow S$ and $\tilde{T}'$ is a proper transform of $T' = T|_S$. The curve $(C \subset \mathbb{P}) = (uw + v^3 + vw \subset \mathbb{P}(1, 1, 2))$ is not well-formed. Therefore we can't use the results 3.12 to calculate (C, B) . The singularities of $\tilde{S}$ are $(u^2 + v^3 + v \subset \mathbb{C}_{u,v,w}^3)/\mathbb{Z}_4(-1, 2, 1) = \mathbb{A}_3$ and $(u^2w + 1 + w \subset \mathbb{C}_{u,v,w}^3)/\mathbb{Z}_2(1, 1, 0) = \mathbb{A}_1$. Since $\deg_C \tilde{T}' = 3$ and $\tilde{T}' \cap \text{Sing } \tilde{S} = \emptyset$ then $(C, B) = (\mathbb{P}^1, \frac{3}{4}P_1 + \frac{1}{2}P_2 + \frac{3}{4}P_3)$. Q.E.D.

Theorem 3.28. *Let $f = t^3 + f_4(z, x, y)$ gives a canonical singularity $(X, 0)$, where f_4 is a homogeneous polynomial of degree 4. Then $\mathbf{p} = (4, 3, 3, 3)$ -blow-up induces a plt blow-up of $(X, 0)$ and $(E, D) = (t + f_4(z, x, y) \subset \mathbb{P}(4, 1, 1, 1), \frac{2}{3}\{t = 0\}) = (\mathbb{P}^2, \frac{2}{3}C)$, where $C = \{f_4 = 0\}$. Then $(X, 0)$ is exceptional if and only if one of the following possibilities holds.*

1. *The curve $C \subset \mathbb{P}^2$ is irreducible. Its singular points can be only ordinary double points.*
2. *Curve $C \subset \mathbb{P}^2$ is the union of two irreducible conics which intersect in 4 distinct points.*

Proof. This theorem is proved as theorem 3.24 □

4. Classification of three-dimensional exceptional canonical hypersurface singularities

The general comment to the tables. The classification of exceptional singularities is given in the tables of the following view. In the first column the singularity number is written. In the second column the singularity equation and the exceptionality condition are written. Almost in all cases the polynomials, that are grouped according to the similar log Del Pezzo surface (E, D) (see 3.12), are written. In the third column corresponding log Del Pezzo surfaces are written. In the upper row the well-formed hypersurface $E \subset \mathbb{P}$ is written. If E is a linear

cone (i.e. $\tilde{d} = \tilde{a}_k$ for some k) then $E = \mathbb{P}(\tilde{a}_1, \dots, \tilde{a}_{k-1}, \tilde{a}_{k+1}, \dots, \tilde{a}_n)$ is written in the second row. As well the different D is written in the second row. The notation is $\text{Diff} = (\frac{q_1-1}{q_1}, \frac{q_2-1}{q_2}, \frac{q_3-1}{q_3}, \frac{q_4-1}{q_4})$. In the fourth column the minimal complement index is written. Let us consider the second column again. Different cases in it are separated by a comma or by a semicolon. If it is a comma then the following singularity has the same minimal complement index as the previous one. If it is a semicolon then the following singularity has the next minimal complement index. Also in the description of different cases we use usual logical symbols: $\&$ – "and", $\parallel$ – "or". For example, the notation $|(n = 4; 5) \& b \neq 0 | \dots | 2, 4|$ means that in the case $n = 4$ and $b \neq 0$ there is 2-complement, but in the case $n = 5$ and $b \neq 0$ there is 4-complement. The symbol $|||$ also means the logical "or", but it is used to separate the polynomials (monomials) only. For example, the notation $h + (x^8 ||| x^7 y)$ means that we have to consider two cases: $h + x^8$, $h + x^7 y$. The exceptionality conditions are also separated by the symbol $|||$.

Very often the conditions that the singularity is canonical are not written. The reader can find them himself. The nonisolated singularity components can be founded with the help of lemma 2.2. By theorem 2.9 the singularity is canonical if and only if it has Du Val singularities along all components. As well the requirement that singularity is normal – $\text{codim}_{\mathbb{C}^4} \text{Sing } X \geq 3$ must be taken into account. For example, the singularity given by $(t^3 - tg(z, x, y) + ag^3(z, x, y) = 0, 0)$ is nonnormal if $a = \frac{4}{3\sqrt{3}}, \frac{2}{3\sqrt{3}}$.

If we know the minimal complement index then the required complement can be easily found.

Quasihomogeneous transformations. Let the quasihomogeneous singularity with weights $\mathbf{p}$ and degree d is given. It can happen that there exist a lot of monomials corresponding to this weighting. On the quasihomogeneous part of the weighting the group of quasihomogeneous biholomorphic maps naturally acts. It is denoted by $H(\mathbf{p})$. Therefore the classification problem is reduced to describe the orbits of action and to find most simple element of every orbit. The general method that can be applied is given in [1]. The group $H(\mathbf{p})$ is generated by the toric and unipotent parts. The toric part is the linear transformations of coordinates. The unipotent part is the quasihomogeneous coordinate change of form $\alpha \mapsto \alpha + h(\beta, \gamma, \delta)$. In our three-dimensional case the singularities don't have the complex structures. Therefore it is easy to find which transformations must be applied. In the tables the most convenient form of singularity is given. After

this we separate to those orbits which have different conditions on the exceptionality.

Conventions. According to described above we introduce the following conventions.

(A) If $g(z, x, y)$ doesn't depend on z then sometimes we use the abridged notation $g(x, y)$. In the third column $g(z^{1/q_2}, x^{1/q_3}, y^{1/q_4})$ is not often given in details because it takes much place in the table.

(B) In the second column the notation $n = 3 \ i \leq 2, \ 4 \ i \leq 1$ means the following condition. If $n = 3$ then the condition on the exceptionality is $i \leq 2$, but if $n = 4$ then the condition on the exceptionality is $i \leq 1$ (let us remember that they have the same complement index).

(C) The notation $i, j \leq 1$ is equivalent to $i \leq 1 \ \& \ j \leq 1$.

(D) $f_n(x, y)$ is a binary form of degree n . In the notation of singularity equation the several binary forms are both denoted by the same one. We can consider that they are always different, i.e. doesn't depend on each other.

Consider notation $x^{ai} f_{n-i}(x^a, y^b)$. One always assumes that the coefficient of monomial x^{an} is nonzero. If there is no coefficient before a binary form then we always suppose that the coefficient of x^{an} is nonzero. Let $a = 1$ and the remaining part of f' doesn't depend on x (i.e. $\text{G.C.D.}(f', x) = 1$). Then we suppose that the irreducible factors of $f_{n-i}(x, y^b)$ have the multiplicities at most i .

(E) Consider notation $x^i(x+y^b)^j f_{n-i-j}(x, y^b)$. Let the remaining part of f' depends on x and doesn't depend on $x + y^b$ (i.e. $\text{G.C.D.}(f', x + y^b) = 1$). Then we suppose that the irreducible factors of $f_{n-i-j}(x, y^b)$ have the multiplicities at most j .

(F) Consider notation $cx^{ai}(x^a + y^b)^j f_{n-i-j}(x^a, y^b)$. The presence of coefficient c means that the coefficients of binary form are arbitrary. If $i + j > n$ then $c = 0$. If $i + j \leq n$ then $c \neq 0$. If there is $c \neq 0$ in the exceptionality condition then we always suppose that the coefficient c is absent (in other words the coefficient of x^{an} is nonzero). Let $a = 1$. If the remaining part of singularity doesn't depend on x then the irreducible factors of $f_{n-i}(x, y^b)$ have the multiplicities at most i . If it depends on x , but not on $x + y^b$ then the irreducible factors have the multiplicities at most j .

(I) It can happen that the quasihomogeneous polynomial has more then one part of type $\mathcal{M}_2$, which belongs to Newton line rotation list. They have the common part of type $\mathcal{M}_1$ and differ by some monomials. In the nontrivial cases their coefficients will be underlined. Assume that k coefficients are underlined. Then we must consider k cases. In each one the corresponding coefficient is absent. It is of a great importance for the binary forms (see (D)–(F)).

(J) If there is a footnote in the first column and in the second row then the exceptionality condition is given at the end of the corresponding table.

(K) The condition on the exceptionality is written for all common components.

Example 4.1. (1) $f = t^2 + z^3 + \underline{a}zx^i f_{5-i}(x, y^2) + \underline{b}yx^j f_{7-j}(x, y^2)$. The exceptionality condition $i \leq 2 \parallel j \leq 4$ (see example 3.20) has the following meaning. Assume that $x + cy^2$ has the multiplicity i_1 in $f_{5-i}(x, y^2)$ and the multiplicity j_1 in $f_{7-j}(x, y^2)$. Then the exceptionality condition means that $i_1 \leq 2 \parallel j_1 \leq 4$. Recall that if $\underline{a} = 0$ or $\underline{b} = 0$ then according to the point (F) one assume that $i > 5$ or $j > 8$ respectively.

(2) $f = t^2 + z^3x + x^i(x + y^2)^j f_{5-i}(x, y^2)$. The exceptionality condition is $i \leq 1 \& j \leq 3$. Here the remaining part $t^2 + z^3x$ depends on x . Therefore the condition $i \leq 1$ is fixed and the requirement $j \leq 3$ as in the previous point means that the multiplicity of any irreducible factor of $f_{5-i}(x, y^2)$ is at most 3.

1. Singularity $-t^2 + z^3 + g(z, x, y)$

No.	$g(z, x, y)$	log Del Pezzo surface	Index
1	$x^7 + azy^{4n} + by^{6n}$ $n = 2, 3, 4, 5, 6$	$t^2 + z^3 + x + azy^4 + by^6 \subset \mathbb{P}(3, 2, 6, 1)$ $\mathbb{P}(3, 2, 1)$, Diff= $(0, 0, \frac{6}{7}, \frac{n-1}{n})$	7
2	$x^7 + azy^{2n} + by^{3n}$ $n = 3; 5; 9, 11, 13$	$t + z^3 + x + azy^2 + by^3 \subset \mathbb{P}(3, 1, 3, 1)$ $\mathbb{P}(3, 1, 1)$, Diff= $(\frac{1}{2}, 0, \frac{6}{7}, \frac{n-1}{n})$	7,8,14
3	$x^7 + y^{2n}$, $n = 4, 5; 8;$ 10, 11, 13; 16, 17, 19, 20	$t^2 + z + x + y^2 \subset \mathbb{P}(1, 2, 2, 1)$ $\mathbb{P}(1, 2, 1)$, Diff= $(0, \frac{2}{3}, \frac{6}{7}, \frac{n-1}{n})$	7,9,15, 21
4	$x^7 + y^n$ $n = 11; 13; 17; 19;$ 23;25;29;31;37,41	$t + z + x + y \subset \mathbb{P}(1, 1, 1, 1)$ $\mathbb{P}^2$, Diff= $(\frac{1}{2}, \frac{2}{3}, \frac{6}{7}, \frac{n-1}{n})$	12,14,18, 21,24,28, 30,36,42
5	$x^i f_{7-i}(x, y^2)$ $i \leq 3$	$t^2 + z + x^i f_{7-i}(x, y^2) \subset \mathbb{P}(7, 14, 2, 1)$ $\mathbb{P}(7, 2, 1)$, Diff= $(0, \frac{2}{3}, 0, 0)$	3
6	$azy^2 x^i f_{4-i}(x, y^3) +$ $x^j f_{7-j}(x, y^3)$, $i \leq 3 j \leq 4$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21, 7, 3, 1)$ $\mathbb{P}(7, 3, 1)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	2
7	$x^i f_{7-i}(x, y^4)$ $i \leq 5$	$t^2 + z + x^i f_{7-i}(x, y^2) \subset \mathbb{P}(7, 14, 2, 1)$ $\mathbb{P}(7, 2, 1)$, Diff= $(0, \frac{2}{3}, 0, \frac{1}{2})$	3
8	$x^i f_{7-i}(x, y^5)$ $i \leq 5$	$t + z + x^i f_{7-i}(x, y) \subset \mathbb{P}(7, 7, 1, 1)$ $\mathbb{P}(7, 1, 1)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, \frac{4}{5})$	6
9	$x^7 + ax^4 y^n + bxy^{2n}$ $n = 5 \text{ } b \neq 0; 7, 11; 13, 17$	$t + z + x^7 + ax^4 y + bxy^2 \subset \mathbb{P}(7, 7, 1, 3)$ $\mathbb{P}(7, 1, 3)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, \frac{n-1}{n})$	6,12,18
10	$x^7 + ax^4 y^{2n} + bxy^{4n}$ $n = 2 \text{ } b \neq 0; 4, 5, 7, 8$	$t^2 + z + g(z, x, y^{1/n}) \subset \mathbb{P}(7, 14, 2, 3)$ $\mathbb{P}(7, 2, 3)$, Diff= $(0, \frac{2}{3}, 0, \frac{n-1}{n})$	3,9
11	$x^7 + xy^n$ $n = 7; 11; 13; 17; 19;$ 23; 25,29;31,35	$t + z + x^7 + xy \subset \mathbb{P}(7, 7, 1, 6)$ $\mathbb{P}(7, 1, 6)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, \frac{n-1}{n})$	8,12,15, 18,22,24, 30,36
12	$x^{2i+1} f_{3-i}(x^2, y^n)$ $n = 5 \text{ } i \leq 1; 7, 11$	$t + z + g(z, x, y^{1/n}) \subset \mathbb{P}(7, 7, 1, 2)$ $\mathbb{P}(7, 1, 2)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, \frac{n-1}{n})$	6,12
13	$azy^n x^{2i} f_{2-i}(x^2, y^{3n}) +$ $+ x^{2i+1} f_{3-j}(x^2, y^{3n})$, $n = 1$ $ij = 0; n = 3 \text{ } i \leq 1 j \leq 2$	$t + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(21, 7, 3, 2)$ $\mathbb{P}(7, 3, 2)$, Diff= $(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	2,4
14	$x^7 + x^2 y^n$, $n = 7; 11;$ 13; 17; 19, 23; 29	$t + z + x^7 + x^2 y \subset \mathbb{P}(7, 7, 1, 5)$ $\mathbb{P}(7, 1, 5)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, \frac{n-1}{n})$	9,12,16, 18,24,30
15	$x^7 + azx^3 y^{2n} + bx^2 y^{6n}$ $n = 2 \text{ } b \neq 0; 3, 4$	$t^2 + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(21, 14, 6, 5)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	4,5
16	$x^7 + azx^3 y^n + bx^2 y^{3n}$ $n = 3 \text{ } b \neq 0; 7, 9$	$t + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(21, 7, 3, 5)$ $\mathbb{P}(7, 3, 5)$, Diff= $(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	4,10
17	$x^7 + x^2 y^{2n}$ $n = 4, 7, 8; 11, 13, 14$	$t^2 + z + x^7 + x^2 y^2 \subset \mathbb{P}(7, 14, 2, 5)$ $\mathbb{P}(7, 2, 5)$, Diff= $(0, \frac{2}{3}, 0, \frac{n-1}{n})$	9,15
18	$x^7 + azx^2 y^{2n} +$ $+ bx^3 y^{3n}$, $n = 3, 5, 7$	$t + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(21, 7, 3, 4)$ $\mathbb{P}(7, 3, 4)$, Diff= $(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	8
19	$x^7 + x^3 y^n$ $n = 7; 11; 13, 17; 19, 23$	$t + z + x^7 + x^3 y \subset \mathbb{P}(7, 7, 1, 4)$ $\mathbb{P}(7, 1, 4)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, \frac{n-1}{n})$	10,12,18, 24
20	$x^7 + zy^n$, $n = 5; 9; 11,$ 13; 15; 17, 19; 23, 25, 27	$t + z^3 + x + zy \subset \mathbb{P}(3, 1, 3, 2)$ $\mathbb{P}(1, 3, 2)$, Diff= $(\frac{1}{2}, 0, \frac{6}{7}, \frac{n-1}{n})$	7,10,14, 16,22,28
21	$x^7 + zxy^{2n}$, $n = 3, 4, 5;$ 6, 7, 8, 9, 10	$t^2 + z^3 + x^7 + zxy^2 \subset \mathbb{P}(21, 14, 6, 11)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	5,11
22	$x^7 + zxy^n$, $n = 5, 7; 9;$	$t + z^3 + x^7 + zxy \subset \mathbb{P}(21, 7, 3, 11)$	8,10,16,

	13, 15; 17, 19, 21	$\mathbb{P}(7, 3, 11)$, Diff= $(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	22
23	$x^7 + zx^2y^n, n = 5, 7, 9;$ 11, 13, 15	$t + z^3 + x^7 + zx^2y \subset \mathbb{P}(21, 7, 3, 8)$ $\mathbb{P}(7, 3, 8)$, Diff= $(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	10, 16
24	$x^i y^j f_{6-i-j}(x, y)$ $i = j = 1$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(4, 8, 1, 1)$ $\mathbb{P}(4, 1, 1)$, Diff= $(0, \frac{2}{3}, 0, 0)$	3
25	$x^8 + az y^{2n} + by^{3n}$ $n = 3, 5; 7$	$t^2 + z^3 + x^2 + az y^2 + by^3 \subset \mathbb{P}(3, 2, 3, 2)$ Diff= $(0, 0, \frac{3}{4}, \frac{n-1}{n})$	5, 8
26	$x^8 + y^n$ $n = 11; 13; 17; 19; 23$	$t^2 + z + x^2 + y \subset \mathbb{P}(1, 2, 1, 2)$ $\mathbb{P}(1, 1, 2)$, Diff= $(0, \frac{2}{3}, \frac{3}{4}, \frac{n-1}{n})$	12, 13, 18, 21, 24
27	$az y^3 x^{4i} f_{1-i}(x^4, y^9) +$ $x^{4j} f_{2-j}(x^4, y^9), i = 0 j = 0$	$t^2 + z^3 + g(z, x^{1/4}, y^{1/3}) \subset \mathbb{P}(3, 2, 3, 1)$ Diff= $(0, 0, \frac{3}{4}, \frac{2}{3})$	4
28	$x^8 + ax^4 y^n + by^{2n}$ $n = 5 \neq 0; 7; 11$	$t^2 + z + x^2 + axy + by^2 \subset \mathbb{P}(1, 2, 1, 1)$ $\mathbb{P}^2$, Diff= $(0, \frac{2}{3}, \frac{3}{4}, \frac{n-1}{n})$	6, 9, 12
29	$az y^2 x^{2i} f_{2-i}(x^2, y^3) +$ $x^{2j} f_{4-j}(x^2, y^3), i \leq 1 j \leq 1$	$t^2 + z^3 + g(z, x^{1/2}, y) \subset \mathbb{P}(6, 4, 3, 1)$ Diff= $(0, 0, \frac{1}{2}, 0)$	2
30	$x^i f_{8-i}(x, y^2)$ $i \leq 4$	$t^2 + z + x^i f_{8-i}(x, y) \subset \mathbb{P}(4, 8, 1, 1)$ $\mathbb{P}(4, 1, 1)$, Diff= $(0, \frac{2}{3}, 0, \frac{1}{2})$	3
31	$x^{2i} f_{4-i}(x^2, y^5)$ $i \leq 2$	$t^2 + z + x^i f_{4-i}(x, y) \subset \mathbb{P}(2, 4, 1, 1)$ $\mathbb{P}(2, 1, 1)$, Diff= $(0, \frac{2}{3}, \frac{1}{2}, \frac{4}{5})$	6
32	$x^8 + xy^n, n = 8; 10;$ 11; 13; 16; 17; 19, 20	$t^2 + z + x^8 + xy \subset \mathbb{P}(4, 8, 1, 7)$ $\mathbb{P}(4, 1, 7)$, Diff= $(0, \frac{2}{3}, 0, \frac{n-1}{n})$	9, 10, 12 13, 18, 21
33	$x^8 + azx^3 y^n + bxy^{3n}$ $b \neq 0 \& (n = 3, 4); 5, 6$	$t^2 + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(12, 8, 3, 7)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	4, 7
34	$x^8 + x^2 y^n$ $n = 7; 11; 13; 17$	$t^2 + z + x^4 + xy \subset \mathbb{P}(2, 4, 1, 3)$ $\mathbb{P}(2, 1, 3)$, Diff= $(0, \frac{2}{3}, \frac{1}{2}, \frac{n-1}{n})$	7, 12, 15, 18
35	$x^8 + ax^5 y^n + bx^2 y^{2n}$ $b \neq 0 \& (n = 4, 5); 7, 8$	$t^2 + z + x^8 + ax^5 y + bx^2 y^2 \subset \mathbb{P}(4, 8, 1, 3)$ $\mathbb{P}(4, 1, 3)$, Diff= $(0, \frac{2}{3}, 0, \frac{n-1}{n})$	6, 9
36	$x^8 + x^3 y^n$ $n = 7; 8; 11; 13, 14$	$t^2 + z + x^8 + x^3 y \subset \mathbb{P}(4, 8, 1, 5)$ $\mathbb{P}(4, 1, 5)$, Diff= $(0, \frac{2}{3}, 0, \frac{n-1}{n})$	7, 9, 12 15
37	$x^8 + azx^2 y^{2n} +$ $+bx^3 y^{3n}, n = 3, 4$	$t^2 + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(12, 8, 3, 5)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	5
38	$x^8 + zy^n$ $n = 7; 9; 11, 13; 15$	$t^2 + z^3 + x^2 + zy \subset \mathbb{P}(3, 2, 3, 4)$ Diff= $(0, 0, \frac{3}{4}, \frac{n-1}{n})$	7, 10, 13, 16
39	$x^8 + zxy^n, n = 5; 6, 7;$ 8, 9, 10; 11, 12	$t^2 + z^3 + x^8 + zxy \subset \mathbb{P}(12, 8, 3, 13)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	5, 7, 10, 13
40	$x^8 + zx^2 y^n$ $n = 7; 9$	$t^2 + z^3 + x^4 + zxy \subset \mathbb{P}(6, 4, 3, 5)$ Diff= $(0, 0, \frac{1}{2}, \frac{n-1}{n})$	7, 10
41	$x^7 y + y^n$ $n = 11; 13; 17; 19; 23;$ 25; 31; 35	$t + z + xy + y^n \subset \mathbb{P}(n, n, n-1, 1)$ $\mathbb{P}(n, n-1, 1)$, Diff= $(\frac{1}{2}, \frac{2}{3}, \frac{6}{7}, 0)$	12, 15, 21, 22, 28, 30, 36, 42
42	$x^7 y + az y^{4n} +$ $+by^{6n}, n = 2, 3, 4, 5$	$t^2 + z^3 + g(z, x^{1/7}, y) \subset \mathbb{P}(3n, 2n, 6n-1, 1)$ Diff= $(0, 0, \frac{6}{7}, 0)$	7
43	$x^7 y + az y^{2n} +$ $+by^{3n}, n = 3; 7; 9, 11$	$t + z^3 + g(z, x^{1/7}, y) \subset \mathbb{P}(3n, n, 3n-1, 1)$ $\mathbb{P}(n, 3n-1, 1)$, Diff= $(\frac{1}{2}, 0, \frac{6}{7}, 0)$	7, 8, 14
44	$x^7 y + y^{2n}, n = 5; 7;$ 8; 10, 13; 14, 16, 17	$t^2 + z + xy + y^{2n} \subset \mathbb{P}(n, 2n, 2n-1, 1)$ $\mathbb{P}(n, 2n-1, 1)$, Diff= $(0, \frac{2}{3}, \frac{6}{7}, 0)$	7, 9, 15, 21
45	$ax x^i f_{5-i}(x, y^2) +$ $+by x^j f_{7-j}(x, y^2)$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(15, 5, 2, 1)$ $\mathbb{P}(5, 2, 1)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	2

	$i \leq 2 j \leq 4$		
46	$yx^i f_{7-i}(x, y^3)$ $i \leq 5$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(11, 22, 3, 1)$ $\mathbb{P}(11, 3, 1), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	3
47	$yx^i f_{7-i}(x, y^4)$ $i \leq 5$	$t + z + g(z, x, y) \subset \mathbb{P}(29, 29, 4, 1)$ $\mathbb{P}(29, 4, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
48	$x^i y + xy^n, n = 8; 12;$ $14; 18; 20; 24, 26; 30$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 1, 7n - 1, n - 1, 6)$ $\mathbb{P}(7n - 1, n - 1, 6), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	9,15,16, 22,24,30, 36
49	$x^i y + ax^4 y^{n+1} + bxy^{2n+1}$ $n = 4 \ b \neq 0; 8, 10; 14$	$t + z + g(z, x, y) \subset \mathbb{P}(7n + 3, 7n + 3, n, 3)$ $\mathbb{P}(7n + 3, n, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6,12,18
50	$x^i y + ax^4 y^{2n+2} + bxy^{4n+3}$ $n = 2 \ b \neq 0; 3, 5, 6$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(7n + 5, 14n + 10, 2n + 1, 3)$ $\mathbb{P}(7n + 5, 2n + 1, 3), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	3,9
51	$yx^{2i+1} f_{3-i}(x^2, y^n)$ $(n = 3, 5) \& i \leq 1; 9$	$t + z + g(z, x, y) \subset \mathbb{P}(7n + 2, 7n + 2, n, 2)$ $\mathbb{P}(7n + 2, n, 2), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6,12
52	$azy^3 x^{2i} f_{2-i}(x^2, y^i) +$ $+yx^{2j+1} f_{3-j}(x^2, y^7)$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(51, 17, 7, 2)$ $\mathbb{P}(17, 7, 2), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
53	$x^i y + x^2 y^n$ $n = 7; 9; 13; 15; 19; 25$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 2, 7n - 2, n - 1, 5)$ $\mathbb{P}(7n - 2, n - 1, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	9,10,16, 18,24,30
54	$x^i y + x^2 y^{2n}$ $n = 5, 6; 9, 11, 12$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(7n - 1, 14n - 2, 2n - 1, 5)$ $\mathbb{P}(7n - 1, 2n - 1, 5), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	9,15
55	$x^i y + azx^3 y^{2n+2} +$ $+bx^2 y^{6n+5}, n = 2, 3$	$t + z^3 + g \subset \mathbb{P}(42n + 33, 14n + 11, 6n + 4, 5)$ $\mathbb{P}(14n + 11, 6n + 4, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
56	$x^i y + azx^3 y^{2n+1} +$ $+bx^2 y^{6n+2}, n = 1 \ b \neq 0; 2, 3$	$t^2 + z^3 + g \subset \mathbb{P}(21n + 6, 14n + 4, 6n + 1, 5)$ $\text{Diff} = \emptyset$	4,5
57	$x^i y + x^3 y^n$ $n = 8; 10; 14, 16; 20$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 3, 7n - 3, n - 1, 4)$ $\mathbb{P}(7n - 3, n - 1, 4), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,12,18, 24
58	$x^i y + azx^2 y^{2n} +$ $+bx^3 y^{3n}, n = 4, 6$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21n - 3, 7n - 1, 3n - 1, 4)$ $\mathbb{P}(7n - 1, 3n - 1, 4), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8
59	$x^i y + zy^n, n = 7; 9;$ $11; 13; 15, 19; 21, 23$	$t + z^3 + xy + zy^n \subset \mathbb{P}(3n, n, 3n - 2, 2)$ $\mathbb{P}(n, 3n - 2, 2), \text{Diff}=(\frac{1}{2}, 0, \frac{6}{7}, 0)$	8,10,14, 16,22,28
60	$x^i y + xzy^n$ $n = 6; 10, 12, 14; 16, 18$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21n - 3, 7n - 1, 3n - 2, 11)$ $\mathbb{P}(7n - 1, 3n - 2, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,16,22
61	$x^i y + xzy^{2n+1}$ $n = 2, 3, 4; 5, 6, 7, 8$	$t^2 + z^3 + g \subset \mathbb{P}(21n + 9, 14n + 6, 6n + 1, 11)$ $\text{Diff} = \emptyset$	5,11
62	$x^i y + zx^2 y^n$ $n = 5, 7, 9; 11, 13$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21n - 6, 7n - 2, 3n - 2, 8)$ $\mathbb{P}(7n - 2, 3n - 2, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,16
63	$\underline{a}zx^i y^{j_1} f_{6-i-i_1}(x, y) +$ $+ \underline{b}x^j y^{j_1} f_{9-j-j_1}(x, y)$ $(i = 1 j \leq 2) \& (i_1 = 1 j_1 \leq 2)$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(9, 3, 1, 1)$ $\mathbb{P}(3, 1, 1), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	2
64	$ax^9 + bx^6 + y^{2n}$ $n = 5; 7; 8$	$t^2 + z^3 + ax^3 + bx^2 + y^2 \subset \mathbb{P}(3, 2, 2, 3)$ $\text{Diff}=(0, 0, \frac{2}{3}, \frac{n-1}{n})$	5,7,9
65	$ax^9 + bx^6 + y^n$ $n = 11; 13; 17$	$t + z^3 + ax^3 + y + bx^2 \subset \mathbb{P}(3, 1, 1, 3)$ $\mathbb{P}(1, 1, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, \frac{n-1}{n})$	12,14,18
66	$\underline{a}zx^{3i} f_{2-i}(x^3, y^4) +$ $+ \underline{b}x^{3j} f_{3-j}(x^3, y^4), i = 0 j \leq 1$	$t^2 + z^3 + g(z, x^{1/3}, y^{1/2}) \subset \mathbb{P}(3, 2, 2, 1)$ $\text{Diff}=(0, 0, \frac{2}{3}, \frac{1}{2})$	3
67	$\underline{a}zx^{3i} f_{2-i}(x^3, y^5) +$ $+ \underline{b}x^{3j} f_{3-j}(x^3, y^5), i \leq 1 j \leq 1$	$t + z^3 + g(z, x^{1/3}, y^{1/5}) \subset \mathbb{P}(3, 1, 1, 1)$ $\mathbb{P}^2, \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, \frac{4}{5})$	6
68	$ax^9 + bx^6 + xy^n$	$t + z^3 + ax^9 + xy + bx^6 \subset \mathbb{P}(9, 3, 1, 8)$	10,12,14,

	$n = 9; 11; 13; 15$	$\mathbb{P}(3, 1, 8), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	16
69	$\underline{a}zx^{2+4i}f_{1-i}(x^4, y^n) + \underline{b}x^{4j+1}f_{2-j}(x^4, y^n)$ $n = 5 \ i = 0 j = 0; 7$	$t + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(9, 3, 1, 4)$ $\mathbb{P}(3, 1, 4), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	6,8
70	$\underline{a}zx^{2i}f_{3-i}(x^2, y^3) + \underline{b}x^{2j+1}f_{4-j}(x^2, y^3), i \leq 1 j \leq 1$	$t + z^3 + g(z, x, y^{1/3}) \subset \mathbb{P}(9, 3, 1, 2)$ $\mathbb{P}(3, 1, 2), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	4
71	$ax^9 + bx^6 + x^2y^{2n}$ $n = 4, 5; 6$	$t^2 + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(9, 6, 2, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5,7
72	$ax^9 + bx^6 + x^2y^n$ $n = 9; 11; 13$	$t + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(9, 3, 1, 7)$ $\mathbb{P}(3, 1, 7), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	10,12,14
73	$\underline{a}zx^{6i}f_{1-i}(x^6, y^n) + \underline{b}x^{6j+3}f_{1-j}(x^6, y^n), n = 7; 11$	$t + z^3 + g(z, x, y^{1/3}, y^{1/n}) \subset \mathbb{P}(3, 1, 1, 2)$ $\mathbb{P}(1, 1, 2), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, \frac{n-1}{n})$	8,12
74	$dx^9 + axxy^{2n} + bx^4y^{2n} + czx^6, (n = 3 \ a \neq 0; 4) \& d + c \neq 0$	$t^2 + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(9, 6, 2, 5)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	3,5
75	$dx^9 + axxy^n + bx^4y^n + czx^6, (n = 7; 9) \& (a + b \neq 0 \ d + c \neq 0)$	$t + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(9, 3, 1, 5)$ $\mathbb{P}(3, 1, 5), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	8,10
76	$x^8y + y^{2n}$ $n = 5, 8$	$t^2 + z + xy + y^{2n} \subset \mathbb{P}(n, 2n, 2n - 1, 1)$ $\mathbb{P}(n, 2n - 1, 1), \text{Diff}=(0, \frac{2}{3}, \frac{7}{8}, 0)$	9
77	$x^8y + azx^{4n} + by^{6n}$ $n = 2, 3$	$t^2 + z^3 + g(z, x^{1/8}, y) \subset \mathbb{P}(3n, 2n, 6n - 1, 1)$ $\text{Diff}=(0, 0, \frac{7}{8}, 0)$	8
78	$x^8y + y^{2n}$ $n = 7; 10$	$t^2 + z + xy + y^{2n} \subset \mathbb{P}(n, 2n, 2n - 1, 1)$ $\mathbb{P}(n, 2n - 1, 1), \text{Diff}=(0, \frac{2}{3}, \frac{7}{8}, 0)$	9,24
79	$yx^{4i}f_{2-i}(x^4, y^{n-1})$ $n = 6 \ i = 0; n = 10 \ i \leq 1$	$t^2 + z + g(z, x^{1/2}, y) \subset \mathbb{P}(2n - 1, 4n - 2, n - 1, 2)$ $\mathbb{P}(2n - 1, n - 1, 2), \text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	3,12
80	$azy^3x^{4i}f_{1-i}(x^4, y^7) + yx^{4j}f_{2-j}(x^4, y^7), i = 0 j \leq 1$	$t^2 + z^3 + g(z, x^{1/2}, y) \subset \mathbb{P}(15, 10, 7, 2)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	4
81	$yx^{2i}f_{4-i}(x^2, y^3)$ $i \leq 2$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(13, 26, 3, 2)$ $\mathbb{P}(13, 3, 2), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	3
82	$yx^if_{8-i}(x, y^2)$ $i \leq 5$	$t + z + g(z, x, y) \subset \mathbb{P}(17, 17, 2, 1)$ $\mathbb{P}(17, 2, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
83	$x^8y + xy^n$ $n = 9; 10; 12; 16; 13; 18$	$t + z + g(z, x, y) \subset \mathbb{P}(8n - 1, 8n - 1, n - 1, 7)$ $\mathbb{P}(8n - 1, n - 1, 7), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,12,18, 30,42
84	$x^8y + axx^3y^{n+1} + bxy^{3n+2}$ $n = 3 \ b \neq 0; 4, 5$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(24n + 15, 8n + 5, 3n + 1, 7)$ $\mathbb{P}(8n + 5, 3n + 1, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4,14
85	$x^8y + x^2y^n$ $n = 8, 12, 14$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(4n - 1, 8n - 2, n - 1, 6)$ $\mathbb{P}(4n - 1, n - 1, 6), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
86	$x^8y + ax^5y^{n+1} + bx^2y^{2n+1}$ $(n = 4, 5) \& b \neq 0; 7$	$t + z + g(z, x, y) \subset \mathbb{P}(8n + 3, 8n + 3, n, 3)$ $\mathbb{P}(8n + 3, n, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6,18
87	$x^8y + x^3y^n$ $n = 7, 8, 10; 13$	$t + z + g(z, x, y) \subset \mathbb{P}(8n - 3, 8n - 3, n - 1, 5)$ $\mathbb{P}(8n - 3, n - 1, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12,30
88	$x^8y + axx^2y^{2n} + bx^3y^{3n}$ $n = 3, 4$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(24n - 3, 8n - 1, 3n - 1, 5)$ $\mathbb{P}(8n - 1, 3n - 1, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
89	$x^8y + zy^n$ $n = 7, 9; 11, 13$	$t^2 + z^3 + x^2y + zy^n \subset \mathbb{P}(3n, 2n, 3n - 2, 4)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	5,16
90	$x^8y + zxy^n$ $n = 6, 7, 8, 9; 10, 11$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(24n - 3, 8n - 1, 3n - 2, 13)$ $\mathbb{P}(8n - 1, 3n - 2, 13), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,26
91	$x^8y + zx^2y^n$ $n = 5, 7$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(12n - 3, 8n - 2, 3n - 2, 10)$ $\text{Diff} = \emptyset$	10

92	$x^7 y^2 + y^n$ $n = 11; 13; 17; 19; 25; 29$	$t + z + xy^2 + y^n \subset \mathbb{P}(n, n, n - 2, 1)$ $\mathbb{P}(n, n - 2, 1)$, Diff= $(\frac{1}{2}, \frac{2}{3}, \frac{6}{7}, 0)$	15,18,24, 28,36,42
93	$x^7 y^2 + y^{2n}$ $n = 5; 7; 11; 13$	$t^2 + z + xy^2 + y^{2n} \subset \mathbb{P}(n, 2n, 2n - 2, 1)$ $\mathbb{P}(n, 2n - 2, 1)$, Diff= $(0, \frac{2}{3}, \frac{6}{7}, 0)$	7,9,15, 21
94	$x^7 y^2 + y^{4n}$ $n = 5; 7$	$t^2 + z + xy^2 + y^{2n} \subset \mathbb{P}(n, 2n, 2n - 1, 1)$ $\mathbb{P}(n, 2n - 1, 1)$, Diff= $(0, \frac{2}{3}, \frac{6}{7}, \frac{1}{2})$	15,21
95	$x^7 y^2 + azy^{2n} + by^{3n}$ $n = 5; 7, 9$	$t + z^3 + g(z, x^{1/7}, y) \subset \mathbb{P}(3n, n, 3n - 2, 1)$ $\mathbb{P}(n, 3n - 2, 1)$, Diff= $(\frac{1}{5}, 0, \frac{6}{7}, 0)$	8,14
96	$x^7 y^2 + azy^{8n} + by^{12n}$ $n = 1, 2$	$t^2 + z^3 + g(z, x^{1/7}, y^{1/2}) \subset \mathbb{P}(3n, 2n, 6n - 1, 1)$ Diff= $(0, 0, \frac{6}{7}, \frac{1}{2})$	7
97	$x^7 y^2 + azy^{12} + by^{18}$	$t^2 + z^3 + g(z, x^{1/7}, y) \subset \mathbb{P}(9, 6, 16, 1)$ Diff= $(0, 0, \frac{6}{7}, 0)$	7
98	$y^2 x^i f_{7-i}(x, y^2)$ $i \leq 4$	$t^2 + z + yx^i f_{7-i}(x, y) \subset \mathbb{P}(4, 8, 1, 1)$ $\mathbb{P}(4, 1, 1)$, Diff= $(0, \frac{2}{3}, 0, \frac{1}{2})$	3
99	$y^2 x^i f_{7-i}(x, y^3)$ $i \leq 5$	$t + z + g(z, x, y) \subset \mathbb{P}(23, 23, 3, 1)$ $\mathbb{P}(23, 3, 1)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
100	$x^7 y^2 + xy^n$ $n = 9; 13; 15; 19, 21; 25$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 2, 7n - 2, n - 2, 6)$ $\mathbb{P}(7n - 2, n - 2, 6)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12,18,22, 30,36
101	$x^{3i+1} y^2 f_{2-i}(x^3, y^{2n})$ $n = 2$ $i = 0; 4$ $i \leq 1$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(7n + 3, 14n + 6, 2n, 3)$ $\mathbb{P}(7n + 3, 2n, 3)$, Diff= $(0, \frac{2}{3}, 0, 0)$	3,9
102	$x^7 y^2 + ax^4 y^{12} + bxy^{22}$	$t^2 + z + x^7 y + ax^4 y^6 + bxy^{11} \subset \mathbb{P}(19, 38, 5, 3)$ $\mathbb{P}(19, 5, 3)$, Diff= $(0, \frac{2}{3}, 0, \frac{1}{2})$	9
103	$x^7 y^2 + ax^4 y^{n+1} + bxy^{2n}$ $n = 6, 8; 12$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 1, 7n - 1, n - 1, 3)$ $\mathbb{P}(7n - 1, n - 1, 3)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12,18
104	$y^2 x^{2i+1} f_{3-i}(x^2, y^n)$ $n = 3$ $i \leq 1; 7$ $i \leq 2$	$t + z + g(z, x, y) \subset \mathbb{P}(7n + 4, 7n + 4, n, 2)$ $\mathbb{P}(7n + 4, n, 2)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6,12
105	$azy^3 x^{2i} f_{2-i}(x^2, y^5) + y^2 x^{2j+1} f_{3-j}(x^2, y^5)$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(39, 13, 5, 2)$ $\mathbb{P}(13, 5, 2)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	4
106	$x^7 y^2 + x^2 y^n$ $n = 9; 11; 15; 21$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 4, 7n - 4, n - 2, 5)$ $\mathbb{P}(7n - 2, n - 2, 5)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12,16,24, 30
107	$x^7 y^2 + x^2 y^{4n}$ $n = 2; 5$	$t^2 + z + x^7 y + x^2 y^{2n} \subset \mathbb{P}(7n - 1, 14n - 2, 2n - 1, 5)$ $\mathbb{P}(7n - 1, 2n - 1, 5)$, Diff= $(0, \frac{2}{3}, 0, \frac{1}{2})$	9,15
108	$x^7 y^2 + x^2 y^{2n}$ $n = 7; 9$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(7n - 2, 14n - 4, 2n - 2, 5)$ $\mathbb{P}(7n - 2, 2n - 2, 5)$, Diff= $(0, \frac{2}{3}, 0, 0)$	9,15
109	$x^7 y^2 + ax^3 y^{n+1} + bx^2 y^{3n+1}$, $n = 4, 6$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21n + 3, 7n + 1, 3n - 1, 5)$ $\mathbb{P}(7n + 1, 3n - 1, 5)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	10
110	$x^7 y^2 + ax^3 y^4 + x^2 y^{10}$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(33, 22, 8, 5)$ Diff= $\emptyset$	4
111	$x^7 y^2 + ax^3 y^6 + bx^2 y^{16}$	$t^2 + z^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(27, 18, 7, 5)$ Diff= $(0, 0, 0, \frac{1}{2})$	5
112	$x^7 y^2 + x^3 y^n$ $n = 7; 11, 13; 17$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 6, 7n - 6, n - 2, 4)$ $\mathbb{P}(7n - 6, n - 2, 4)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,18,24
113	$x^7 y^2 + ax^2 y^{2n} + bx^3 y^{3n}$, $n = 3, 5$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21n - 6, 7n - 2, 3n - 2, 4)$ $\mathbb{P}(7n - 2, 3n - 2, 4)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	8
114	$x^7 y^2 + zy^n$ $n = 7; 9; 11; 15; 17, 19$	$t + z^3 + xy^2 + zy^n \subset \mathbb{P}(3n, n, 3n - 4, 2)$ $\mathbb{P}(n, 3n - 4, 2)$, Diff= $(\frac{1}{2}, 0, \frac{6}{7}, 0)$	10,14,16, 22,28
115	$x^7 y^2 + zxy^{4n+2}$ $n = 1; 2, 3$	$t^2 + z^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(21n + 9, 14n + 6, 6n + 1, 11)$ Diff= $(0, 0, 0, \frac{1}{2})$	5,11

116	$x^7y^2 + zxy^n$ $n = 7; 9, 11; 13, 15$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21n - 6, 7n - 2, 3n - 4, 11)$ $\mathbb{P}(7n - 2, 3n - 4, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,16,22
117	$x^7y^2 + zxy^{4n}$ $n = 2; 3$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(42n - 3, 28n - 2, 12n - 4, 11)$ $\text{Diff} = \emptyset$	5,11
118	$x^7y^2 + zx^2y^n$ $n = 5, 7; 9, 11$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21n - 12, 7n - 4, 3n - 4, 8)$ $\mathbb{P}(7n - 4, 3n - 4, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,16
119	$x^iy^jf_{10-i-j}(x, y)$ $i \leq 3 \& j \leq 3$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(5, 10, 1, 1)$ $\mathbb{P}(5, 1, 1), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	3
120	$x^{10} + y^n$ $n = 11; 13$	$t^2 + z + x^2 + y \subset \mathbb{P}(1, 2, 1, 2)$ $\mathbb{P}(1, 1, 2), \text{Diff}=(0, \frac{2}{3}, \frac{4}{5}, \frac{n-1}{n})$	12,15
121	$zy^2(ax^5 + by^6) +$ $\underline{c}x^{5j}f_{2-j}(x^5, y^6), b \neq 0 j = 0$	$t^2 + z^3 + g(z, x^{1/5}, y^{1/2}) \subset \mathbb{P}(3, 2, 3, 1)$ $\text{Diff}=(0, 0, \frac{4}{5}, \frac{1}{2})$	5
122	$x^{10} + ax^5y^7 + by^{14}$	$t^2 + z + x^2 + axy + by^2 \subset \mathbb{P}(1, 2, 1, 1)$ $\mathbb{P}^2, \text{Diff}=(0, \frac{2}{3}, \frac{4}{5}, \frac{6}{7})$	15
123	$x^{10} + xy^n$ $n = 10, 11; 13$	$t^2 + z + x^{10} + xy \subset \mathbb{P}(5, 10, 1, 9)$ $\mathbb{P}(5, 1, 9), \text{Diff}=(0, \frac{2}{3}, 0, \frac{n-1}{n})$	12,27
124	$x^{3i+1}f_{3-i}(x^3, y^4)$ $i = 0, 1$	$t^2 + z + x^{3i+1}f_{3-i}(x^3, y) \subset \mathbb{P}(5, 10, 1, 3)$ $\mathbb{P}(5, 1, 3), \text{Diff}=(0, \frac{2}{3}, 0, \frac{3}{4})$	9
125	$x^{10} + ax^4y^3 +$ $+x^2y^9$	$t^2 + z^3 + x^{10} + ax^4y + x^2y^3 \subset \mathbb{P}(15, 10, 3, 8)$ $\text{Diff}=(0, 0, 0, \frac{2}{5})$	4
126	$x^{10} + ax^6y^5 + x^2y^{10}$	$t^2 + z + x^{10} + ax^6y + x^2y^2 \subset \mathbb{P}(5, 10, 1, 4)$ $\mathbb{P}(5, 1, 4), \text{Diff}=(0, \frac{2}{3}, 0, \frac{4}{5})$	6
127	$x^{10} + x^2y^{11}$	$t^2 + z + x^{10} + x^2y \subset \mathbb{P}(5, 10, 1, 8)$ $\mathbb{P}(5, 1, 8), \text{Diff}=(0, \frac{2}{3}, 0, \frac{10}{11})$	12
128	$x^{10} + x^3y^n$ $n = 8; 10$	$t^2 + z + x^{10} + x^3y \subset \mathbb{P}(5, 10, 1, 7)$ $\mathbb{P}(5, 1, 7), \text{Diff}=(0, \frac{2}{3}, 0, \frac{n-1}{n})$	9,21
129	$x^{10} + ax^2y^6 +$ $+bx^3y^9$	$t^2 + z^3 + g(z, x, y^{1/3}) \subset \mathbb{P}(15, 10, 3, 7)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	7
130	$x^{10} + x^4y^7$	$t^2 + z + x^{10} + x^4y \subset \mathbb{P}(5, 10, 1, 6)$ $\mathbb{P}(5, 1, 6), \text{Diff}=(0, \frac{2}{3}, 0, \frac{6}{7})$	9
131	$x^{10} + zy^n$ $n = 7; 9$	$t^2 + z^3 + zy + x^2 \subset \mathbb{P}(3, 2, 3, 4)$ $\text{Diff}=(0, 0, \frac{4}{5}, \frac{n-1}{n})$	7,10
132	$x^{10} + zxy^n$ $n = 6, 7; 8$	$t^2 + z^3 + zxy + x^{10} \subset \mathbb{P}(15, 10, 3, 17)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	7,17
133	$x^{10} + zx^2y^5$	$t^2 + z^3 + zx^2y + x^{10} \subset \mathbb{P}(15, 10, 3, 14)$ $\text{Diff}=(0, 0, 0, \frac{4}{5})$	7
134	$x^{10} + zx^3y^5$	$t^2 + z^3 + zx^3y + x^{10} \subset \mathbb{P}(15, 10, 3, 11)$ $\text{Diff}=(0, 0, 0, \frac{4}{5})$	11
135	$x^9y + azx^3y^4 + y^{11}$	$t + z^3 + g(z, x^{1/3}, y) \subset \mathbb{P}(33, 11, 10, 3)$ $\mathbb{P}(11, 10, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	4
136	$x^9y + azy^8 + by^{12}$	$t^2 + z^3 + azy^8 + xy + by^{12} \subset \mathbb{P}(6, 4, 11, 1)$ $\text{Diff}=(0, 0, \frac{8}{9}, 0)$	9
137	$\underline{a}zy^2x^{3i+2}f_{1-i}(x^3, y^4) +$ $+byx^{3j}f_{3-j}(x^3, y^4)$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(39, 13, 4, 3)$ $\mathbb{P}(13, 4, 3), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
138	$x^9y + azx^3y^5 + by^{14}$	$t^2 + z^3 + g(z, x^{1/3}, y) \subset \mathbb{P}(21, 14, 13, 3)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	9
139	$x^9y + azy^{10} + by^{15}$	$t + z^3 + g(z, x^{1/9}, y) \subset \mathbb{P}(15, 5, 14, 1)$ $\mathbb{P}(5, 14, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{8}{9}, 0)$	18

140	$x^9y + xy^n$ $n = 10; 12; 14$	$t + z + g(z, x, y) \subset \mathbb{P}(9n - 1, 9n - 1, n - 1, 8)$ $\mathbb{P}(9n - 1, n - 1, 8), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12,30,48
141	$x^9y + ax^5y^6 + xy^{11}$	$t + z + g(z, x, y) \subset \mathbb{P}(49, 49, 5, 4)$ $\mathbb{P}(49, 5, 4), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
142	$yx^{2i+1}f_{4-i}(x^2, y^3)$ $i \leq 2$	$t + z + g(z, x, y) \subset \mathbb{P}(29, 29, 3, 2)$ $\mathbb{P}(29, 3, 2), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12
143	$x^9y + x^2y^n$ $n = 9, 11$	$t + z + g(z, x, y) \subset \mathbb{P}(9n - 2, 9n - 2, n - 1, 7)$ $\mathbb{P}(9n - 2, n - 1, 7), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12
144	$x^9y + x^2y^{2n}$ $n = 5; 6$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(9n - 1, 18n - 2, 2n - 1, 7)$ $\mathbb{P}(9n - 1, 2n - 1, 7), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	12,21
145	$x^9y + azx^4y^3 + x^3y^8$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(69, 23, 7, 6)$ $\mathbb{P}(23, 7, 6), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
146	$x^9y + x^4y^n$ $n = 7; 9$	$t + z + g(z, x, y) \subset \mathbb{P}(9n - 4, 9n - 4, n - 1, 5)$ $\mathbb{P}(9n - 4, n - 1, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18,30
147	$x^9y + x^4y^8$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(34, 68, 7, 5)$ $\mathbb{P}(34, 7, 5), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	15
148	$x^9y + zy^n$ $n = 7, 9$	$t + z^3 + xy + zy^n \subset \mathbb{P}(3n, n, 3n - 2, 2)$ $\mathbb{P}(n, 3n - 2, 2), \text{Diff}=(\frac{1}{2}, 0, \frac{8}{9}, 0)$	10
149	$x^9y + zxy^n$ $n = 6, 8$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(27n - 3, 9n - 1, 3n - 2, 15)$ $\mathbb{P}(9n - 1, 3n - 2, 15), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
150	$x^9y + zxy^7$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(93, 62, 19, 15)$ $\text{Diff} = \emptyset$	5
151	$x^9y + zx^2y^n$ $n = 5; 7$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(27n - 6, 9n - 2, 3n - 2, 12)$ $\mathbb{P}(9n - 2, 3n - 2, 12), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14,24
152	$x^8y^2 + y^n$ $n = 11; 13; 17$	$t^2 + z + x^2y^2 + y^n \subset \mathbb{P}(n, 2n, n - 2, 2)$ $\mathbb{P}(n, n - 2, 2), \text{Diff}=(0, \frac{2}{3}, \frac{3}{4}, 0)$	16,18,24
153	$x^8y^2 + azy^{10} + by^{15}$	$t^2 + z^3 + g(z, x^{1/4}, y) \subset \mathbb{P}(15, 10, 13, 2)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	8
154	$azy^3x^{4i}f_{1-i}(x^4, y^5) + y^2x^{4j}f_{2-j}(x^4, y^5)$	$t^2 + z^3 + g(z, x^{1/4}, y) \subset \mathbb{P}(6, 4, 5, 1)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	4
155	$x^8y^2 + ax^4y^9 + by^{16}$	$t^2 + z + x^2y^2 + axy^9 + by^{16} \subset \mathbb{P}(8, 16, 7, 1)$ $\mathbb{P}(8, 7, 1), \text{Diff}=(0, \frac{2}{3}, \frac{3}{4}, 0)$	12
156	$y^2x^{2i}f_{4-i}(x^2, y^3)$ $i \leq 2$	$t^2 + z + y^2x^if_{4-i}(x, y^3) \subset \mathbb{P}(7, 14, 3, 1)$ $\mathbb{P}(7, 3, 1), \text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	6
157	$x^8y^2 + azx^3y^{n+1} + bxy^{3n+1}, n = 3 \text{ } b \neq 0; 4$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(12n + 3, 8n + 2, 3n - 1, 7)$ $\text{Diff} = \emptyset$	4,7
158	$x^8y^2 + xy^n$ $n = 11, 12; 14; 15$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(4n - 1, 8n - 2, n - 2, 7)$ $\mathbb{P}(4n - 1, n - 2, 7), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	15,18,21
159	$x^8y^2 + x^2y^n$ $n = 9; 13$	$t^2 + z + x^4y^2 + xy^n \subset \mathbb{P}(2n - 1, 4n - 2, n - 2, 3)$ $\mathbb{P}(2n - 1, n - 2, 3), \text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	12,18
160	$x^{3i+2}y^2f_{2-i}(x^3, y^{n-2})$ $n = 6 \text{ } i = 0; \text{ } 7 \text{ } i \leq 1$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(4n - 5, 8n - 10, n - 2, 3)$ $\mathbb{P}(4n - 5, n - 2, 3), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	6,9
161	$x^8y^2 + x^3y^n$ $n = 8; 10; 11$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(4n - 3, 8n - 6, n - 2, 5)$ $\mathbb{P}(4n - 3, n - 2, 5), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	9,12,15
162	$x^8y^2 + azx^2y^6 + bx^3y^9$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(33, 22, 7, 5)$ $\text{Diff} = \emptyset$	5
163	$x^8y^2 + zy^n$ $n = 7; 9; 11$	$t^2 + z^3 + x^2y^2 + zy^n \subset \mathbb{P}(3n, 2n, 3n - 4, 4)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	10,13,16

164	$x^8y^2 + zxy^n$ $n = 6; 7, 8; 9$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(12n - 3, 8n - 2, 3n - 4, 13)$ $\text{Diff} = \emptyset$	7, 10, 13
165	$x^8y^2 + zx^2y^n$ $n = 5; 7$	$t^2 + z^3 + x^4y^2 + zxy^n \subset \mathbb{P}(6n - 3, 4n - 2, 3n - 4, 5)$ $\text{Diff} = (0, 0, \frac{1}{2}, 0)$	7, 10
166	$x^7y^3 + y^n$ $n = 11; 13; 19; 23$	$t + z + xy^3 + y^n \subset \mathbb{P}(n, n, n - 3, 1)$ $\mathbb{P}(n, n - 3, 1), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, \frac{6}{7}, 0)$	18, 22, 36, 42
167	$x^7y^3 + y^{2n}$ $n = 7, 8; 10, 11$	$t^2 + z + xy^3 + y^{2n} \subset \mathbb{P}(n, 2n, 2n - 3, 1)$ $\mathbb{P}(n, 2n - 3, 1), \text{Diff} = (0, \frac{2}{3}, \frac{6}{7}, 0)$	15, 21
168	$x^7y^3 + az y^{2n} + by^{3n}$ $n = 5; 7$	$t + z^3 + g(z, x^{1/7}, y) \subset \mathbb{P}(3n, n, 3n - 3, 1)$ $\mathbb{P}(n, 3n - 3, 1), \text{Diff} = (\frac{1}{2}, 0, \frac{6}{7}, 0)$	8, 14
169	$x^7y^3 + az y^8 + by^{12}$	$t^2 + z^3 + g(z, x^{1/7}, y) \subset \mathbb{P}(6, 4, 9, 1)$ $\text{Diff} = (0, 0, \frac{6}{7}, 0)$	7
170	$x^7y^3 + az y^{12} + by^{18}$	$t^2 + z^3 + g(z, x^{1/7}, y^{1/3}) \subset \mathbb{P}(3, 2, 5, 1)$ $\text{Diff} = (0, 0, \frac{6}{7}, \frac{2}{3})$	7
171	$y^3x^i f_{7-i}(x, y^2)$ $i \leq 5$	$t + z + g(z, x, y) \subset \mathbb{P}(17, 17, 2, 1)$ $\mathbb{P}(17, 2, 1), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
172	$x^7y^3 + xy^n$ $n = 10; 14; 16; 20$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 3, 7n - 3, n - 3, 6)$ $\mathbb{P}(7n - 3, n - 3, 6), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	16, 24, 30 36
173	$x^7y^3 + ax^4y^{n+2} + bxy^{2n+1}$ $n = 5; 9$	$t + z + g(z, x, y) \subset \mathbb{P}(7n + 2, 7n + 2, n - 1, 3)$ $\mathbb{P}(7n + 2, n - 1, 3), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	12, 18
174	$x^7y^3 + ax^4y^{2n+2} + bxy^{4n+1}$ $n = 3, 4$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(7n + 1, 14n + 2, 2n - 1, 3)$ $\mathbb{P}(7n + 1, 2n - 1, 3), \text{Diff} = (0, \frac{2}{3}, 0, 0)$	9
175	$y^3x^{2i+1}f_{3-i}(x^2, y^3)$ $i = 0, 1$	$t + z^3 + yx^{2i+1}f_{3-i}(x^2, y) \subset \mathbb{P}(9, 3, 1, 2)$ $\mathbb{P}(3, 1, 2), \text{Diff} = (\frac{1}{2}, 0, 0, \frac{2}{3})$	4
176	$y^3x^{2i+1}f_{3-i}(x^2, y^5)$ $i \leq 2$	$t + z + g(z, x, y) \subset \mathbb{P}(41, 41, 5, 2)$ $\mathbb{P}(41, 5, 2), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	12
177	$x^7y^3 + azx^3y^4 + x^2y^9$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(57, 19, 6, 5)$ $\mathbb{P}(19, 6, 5), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	4
178	$x^7y^3 + x^2y^{2n}$ $n = 5; 7, 8$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(7n - 3, 14n - 6, 2n - 3, 5)$ $\mathbb{P}(7n - 3, 2n - 3, 5), \text{Diff} = (0, \frac{2}{3}, 0, 0)$	9, 15
179	$x^7y^3 + x^2y^n$ $n = 11; 17$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 6, 7n - 6, n - 3, 5)$ $\mathbb{P}(7n - 6, n - 3, 5), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	18, 30
180	$x^7y^3 + azx^3y^5 + bx^2y^{12}$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(39, 26, 9, 5)$ $\text{Diff} = \emptyset$	5
181	$x^7y^3 + azx^3y^6 + bx^2y^{15}$	$t + z^3 + g(z, x, y^{1/3}) \subset \mathbb{P}(33, 11, 4, 5)$ $\mathbb{P}(11, 4, 5), \text{Diff} = (\frac{1}{2}, 0, 0, \frac{2}{3})$	10
182	$x^7y^3 + x^3y^n$ $n = 8; 10; 14$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 9, 7n - 9, n - 3, 4)$ $\mathbb{P}(7n - 9, n - 3, 4), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	12, 18, 24
183	$x^7y^3 + azx^2y^8 + bx^3y^{12}$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(75, 25, 9, 4)$ $\mathbb{P}(25, 9, 4), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	8
184	$x^7y^3 + zy^n$ $n = 7; 11, 13$	$t + z^3 + xy^3 + zy^n \subset \mathbb{P}(3n, n, 3n - 6, 2)$ $\mathbb{P}(n, 3n - 6, 2), \text{Diff} = (\frac{1}{2}, 0, \frac{6}{7}, 0)$	10, 22
185	$x^7y^3 + zy^{15}$	$t + z^3 + xy + zy^5 \subset \mathbb{P}(15, 5, 13, 2)$ $\mathbb{P}(5, 13, 2), \text{Diff} = (\frac{1}{2}, 0, \frac{6}{7}, \frac{2}{3})$	28
186	$x^7y^3 + zxy^{3n}$ $n = 2; 4$	$t + z^3 + x^7y + zxy^n \subset \mathbb{P}(21n - 3, 7n - 1, 3n - 2, 11)$ $\mathbb{P}(7n - 1, 3n - 6, 11), \text{Diff} = (\frac{1}{2}, 0, 0, \frac{2}{3})$	10, 22
187	$x^7y^3 + zxy^n$ $n = 7; 11$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(21n - 9, 14n - 6, 6n - 12, 22)$ $\text{Diff} = \emptyset$	5, 11

188	$x^7y^3 + zxy^{2n}$ $n = 4, 5$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(42n - 9, 14n - 3, 6n - 6, 11)$ $\mathbb{P}(14n - 3, 6n - 6, 11)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	16
189	$x^7y^3 + zxy^9$	$t^2 + z^3 + x^7y + zxy^3 \subset \mathbb{P}(30, 20, 7, 11)$ Diff= $(0, 0, 0, \frac{2}{3})$	11
190	$x^7y^3 + zx^2y^n$ $n = 5, 7$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21n - 18, 7n - 6, 3n - 6, 8)$ $\mathbb{P}(7n - 6, 3n - 6, 8)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	10
191	$x^7y^3 + zx^2y^9$	$t + z^3 + x^7y + zx^2y^3 \subset \mathbb{P}(57, 19, 7, 8)$ $\mathbb{P}(19, 7, 8)$, Diff= $(\frac{1}{2}, 0, 0, \frac{2}{3})$	16
192	$x^i y^j f_{11-i-j}(x, y)$ $i \leq 4 \& j \leq 4$	$t + z + g(z, x, y) \subset \mathbb{P}(11, 11, 1, 1)$ $\mathbb{P}(11, 1, 1)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
193	$x^{11} + azy^8 + by^{12}$	$t^2 + z^3 + x + azy^4 + by^6 \subset \mathbb{P}(3, 2, 6, 1)$ $\mathbb{P}(3, 2, 1)$, Diff= $(0, 0, \frac{10}{11}, \frac{1}{2})$	11
194	$x^{11} + y^{13}$	$t + z + x + y \subset \mathbb{P}(1, 1, 1, 1)$ $\mathbb{P}^2$, Diff= $(\frac{1}{2}, \frac{2}{3}, \frac{10}{11}, \frac{12}{13})$	66
195	$x^{11} + xy^{11}$	$t + z + x^{11} + xy \subset \mathbb{P}(11, 11, 1, 10)$ $\mathbb{P}(11, 1, 10)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, \frac{10}{11})$	12
196	$x^{11} + x^2y^{10}$	$t^2 + z + x^{11} + x^2y^2 \subset \mathbb{P}(11, 22, 2, 9)$ $\mathbb{P}(11, 2, 9)$, Diff= $(0, \frac{2}{3}, 0, \frac{4}{5})$	15
197	$x^{11} + azx^2y^6 + bx^3y^9$	$t + z^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(33, 11, 3, 8)$ $\mathbb{P}(11, 3, 8)$, Diff= $(\frac{1}{2}, 0, 0, \frac{2}{3})$	16
198	$x^{11} + x^4y^8$	$t^2 + z + x^{11} + x^4y^2 \subset \mathbb{P}(11, 22, 2, 7)$ $\mathbb{P}(11, 2, 7)$, Diff= $(0, \frac{2}{3}, 0, \frac{3}{4})$	21
199	$x^{11} + x^5y^7$	$t + z + x^{11} + x^5y \subset \mathbb{P}(11, 11, 1, 6)$ $\mathbb{P}(11, 1, 6)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, \frac{6}{7})$	36
200	$x^{11} + zxy^7$	$t + z^3 + x^{11} + zxy \subset \mathbb{P}(33, 11, 3, 19)$ $\mathbb{P}(11, 3, 19)$, Diff= $(\frac{1}{2}, 0, 0, \frac{6}{7})$	14
201	$x^{11} + zx^3y^5$	$t + z^3 + x^{11} + zx^3y \subset \mathbb{P}(33, 11, 3, 13)$ $\mathbb{P}(11, 3, 13)$, Diff= $(\frac{1}{2}, 0, 0, \frac{4}{5})$	26
202	$x^{10}y + azy^8 + by^{12}$	$t^2 + z^3 + g(z, x^{1/10}, y) \subset \mathbb{P}(6, 4, 11, 1)$ Diff= $(0, 0, \frac{9}{10}, 0)$	10
203	$x^{10}y + ax^5y^7 + by^{13}$	$t + z + x^2y + axy^7 + by^{13} \subset \mathbb{P}(13, 13, 6, 1)$ $\mathbb{P}(13, 6, 1)$, Diff= $(\frac{1}{2}, \frac{2}{3}, \frac{4}{5}, 0)$	30
204	$x^{10}y + xy^n$ $n = 11; 12$	$t + z + g(z, x, y) \subset \mathbb{P}(10n - 1, 10n - 1, n - 1, 9)$ $\mathbb{P}(10n - 1, n - 1, 9)$, Diff= $(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12, 54
205	$x^{10}y + x^2y^{10}$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(49, 98, 9, 8)$ $\mathbb{P}(49, 9, 8)$, Diff= $(0, \frac{2}{3}, 0, 0)$	12
206	$x^{10}y + azx^2y^6 + bx^3y^9$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(87, 29, 8, 7)$ $\mathbb{P}(29, 8, 7)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	14
207	$x^{10}y + x^4y^8$	$t^2 + z + x^5y + x^2y^8 \subset \mathbb{P}(19, 38, 7, 3)$ $\mathbb{P}(19, 7, 3)$, Diff= $(0, \frac{2}{3}, \frac{1}{2}, 0)$	18
208	$x^{10}y + zxy^7$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(207, 69, 19, 17)$ $\mathbb{P}(69, 19, 17)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	24
209	$x^{10}y + zx^3y^5$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(141, 47, 13, 11)$ $\mathbb{P}(47, 13, 11)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	22
210	$x^9y^2 + azy^8 + by^{12}$	$t^2 + z^3 + xy + azy^4 + by^6 \subset \mathbb{P}(3, 2, 5, 1)$ Diff= $(0, 0, \frac{8}{9}, \frac{1}{2})$	9
211	$x^9y^2 + azx^3y^5 + by^{13}$	$t + z^3 + g(z, x^{1/3}, y) \subset \mathbb{P}(39, 13, 11, 3)$ $\mathbb{P}(13, 11, 3)$, Diff= $(\frac{1}{2}, 0, \frac{2}{3}, 0)$	18

212	$x^9y^2 + xy^{11}$	$t + z + g(z, x, y) \subset \mathbb{P}(97, 97, 9, 8)$ $\mathbb{P}(97, 9, 8), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	30
213	$x^9y^2 + ax^5y^7 + bxy^{12}$	$t + z + g(z, x, y) \subset \mathbb{P}(53, 53, 5, 4)$ $\mathbb{P}(53, 5, 4), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	24
214	$x^9y^2 + x^2y^{10}$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(43, 86, 8, 7)$ $\mathbb{P}(43, 8, 7), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
215	$x^9y^2 + x^2y^{11}$	$t + z + g(z, x, y) \subset \mathbb{P}(95, 95, 9, 7)$ $\mathbb{P}(95, 9, 7), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	42
216	$x^9y^2 + ax^2y^6 + bx^3y^9$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(75, 25, 7, 6)$ $\mathbb{P}(25, 7, 6), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	12
217	$x^9y^2 + x^4y^8$	$t^2 + z + x^9y + x^4y^4 \subset \mathbb{P}(16, 32, 3, 5)$ $\mathbb{P}(16, 3, 5), \text{Diff}=(0, \frac{2}{3}, 0, \frac{1}{2})$	15
218	$x^9y^2 + zy^9$	$t + z^3 + xy^2 + zy^9 \subset \mathbb{P}(27, 9, 23, 2)$ $\mathbb{P}(9, 23, 2), \text{Diff}=(\frac{1}{2}, 0, \frac{8}{9}, 0)$	36
219	$x^9y^2 + zxy^7$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(183, 61, 17, 15)$ $\mathbb{P}(61, 17, 15), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
220	$x^8y^3 + azy^8 + by^{12}$	$t^2 + z^3 + g(z, x^{1/8}, y) \subset \mathbb{P}(6, 4, 9, 1)$ $\text{Diff}=(0, 0, \frac{7}{8}, 0)$	8
221	$x^8y^3 + ax^4y^8 + by^{13}$	$t^2 + z + x^4y^3 + ax^2y^8 + by^{13} \subset \mathbb{P}(13, 26, 5, 2)$ $\mathbb{P}(13, 5, 2), \text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	12
222	$x^8y^3 + y^{14}$	$t^2 + z + xy^3 + y^{14} \subset \mathbb{P}(7, 14, 11, 1)$ $\mathbb{P}(7, 11, 1), \text{Diff}=(0, \frac{2}{3}, \frac{7}{8}, 0)$	24
223	$x^8y^3 + xy^n$ $n = 11; 13$	$t + z + g(z, x, y) \subset \mathbb{P}(8n - 3, 8n - 3, n - 3, 7)$ $\mathbb{P}(8n - 3, n - 3, 7), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18, 42
224	$x^8y^3 + ax^3y^5 + bxy^{12}$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(93, 31, 9, 7)$ $\mathbb{P}(31, 9, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
225	$x^8y^3 + x^2y^{10}$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(37, 74, 7, 6)$ $\mathbb{P}(37, 7, 6), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
226	$x^8y^3 + ax^5y^7 + bx^2y^{11}$	$t + z + g(z, x, y) \subset \mathbb{P}(41, 41, 4, 3)$ $\mathbb{P}(41, 4, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18
227	$\underline{d}x^8y^3 + z(\underline{a}x^7 + bx^2y^6) +$ $+cx^3y^9, b + c \neq 0$	$t + z^3 + g(z, x, y^{1/3}) \subset \mathbb{P}(21, 7, 2, 5)$ $\mathbb{P}(7, 2, 5), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	10
228	$x^8y^3 + x^3y^{10}$	$t + z + g(z, x, y) \subset \mathbb{P}(71, 71, 7, 5)$ $\mathbb{P}(71, 7, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	30
229	$x^8y^3 + zy^9$	$t^2 + z^3 + x^2y + zy^3 \subset \mathbb{P}(9, 6, 7, 4)$ $\text{Diff}=(0, 0, \frac{3}{4}, \frac{2}{3})$	16
230	$x^8y^3 + zxy^n$ $n = 7; 8$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(24n - 9, 8n - 3, 3n - 6, 13)$ $\mathbb{P}(8n - 3, 3n - 6, 13), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10, 26
231	$\underline{d}x^7y^4 + z(\underline{a}x^7 + by^8) +$ $+cy^{12}, b + c \neq 0$	$t^2 + z^3 + g(z, x^{1/7}, y^{1/2}) \subset \mathbb{P}(3, 2, 4, 1)$ $\text{Diff}=(0, 0, \frac{6}{7}, \frac{1}{2})$	7
232	$\underline{d}x^7y^4 + y^n$ $n = 13; 17$	$t + z + xy^4 + y^n \subset \mathbb{P}(n, n, n - 4, 1)$ $\mathbb{P}(n, n - 4, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, \frac{6}{7}, 0)$	30, 42
233	$x^7y^4 + y^{14}$	$t^2 + z + xy^4 + y^{14} \subset \mathbb{P}(7, 14, 10, 1)$ $\mathbb{P}(7, 10, 1), \text{Diff}=(0, \frac{2}{3}, \frac{6}{7}, 0)$	15
234	$x^7y^4 + azy^{10} + by^{15}$	$t + z^3 + g(z, x^{1/7}, y) \subset \mathbb{P}(15, 5, 11, 1)$ $\mathbb{P}(5, 11, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{6}{7}, 0)$	14
235	$x^7y^4 + y^{16}$	$t^2 + z + xy + y^4 \subset \mathbb{P}(2, 4, 3, 1)$ $\mathbb{P}(2, 3, 1), \text{Diff}=(0, \frac{2}{3}, \frac{6}{7}, \frac{3}{4})$	21

236	$x^r y^4 + xy^n$ $n = 11; 15$	$t + z + g(z, x, y) \subset \mathbb{P}(7n - 4, 7n - 4, n - 4, 6)$ $\mathbb{P}(7n - 4, n - 4, 6), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	24,36
237	$x^r y^4 + ax^4 y^8 + bxy^{12}$	$t^2 + z + x^r y + ax^4 y^2 + bxy^3 \subset \mathbb{P}(5, 10, 1, 3)$ $\mathbb{P}(5, 1, 3), \text{Diff}=(0, \frac{2}{3}, 0, \frac{3}{4})$	9
238	$x^r y^4 + ax^4 y^9 + bxy^{14}$	$t + z + g(z, x, y) \subset \mathbb{P}(47, 47, 5, 3)$ $\mathbb{P}(47, 5, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18
239	$x^{2i+1} y^4 f_{3-i}(x^2, y^3)$ $i \leq 2$	$t + z + g(z, x, y) \subset \mathbb{P}(29, 29, 3, 2)$ $\mathbb{P}(29, 3, 2), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12
240	$x^r y^4 + x^2 y^{10}$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(31, 62, 6, 5)$ $\mathbb{P}(31, 6, 5), \text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
241	$x^r y^4 + ax^3 y^5 + bx^2 y^{11}$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(69, 23, 7, 5)$ $\mathbb{P}(23, 7, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
242	$x^r y^4 + x^2 y^{12}$	$t^2 + z + x^r y^2 + x^2 y^6 \subset \mathbb{P}(19, 38, 4, 5)$ $\mathbb{P}(19, 4, 5), \text{Diff}=(0, \frac{2}{3}, 0, \frac{1}{2})$	15
243	$x^r y^4 + x^2 y^{13}$	$t + z + g(z, x, y) \subset \mathbb{P}(83, 83, 9, 5)$ $\mathbb{P}(83, 9, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	30
244	$\underline{dx}^r y^4 + z(\underline{ax}^6 y + bx^2 y^6) +$ $+cx^3 y^9, b + c \neq 0$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(51, 17, 5, 4)$ $\mathbb{P}(17, 5, 4), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8
245	$x^r y^4 + x^3 y^{11}$	$t + z + g(z, x, y) \subset \mathbb{P}(65, 65, 7, 4)$ $\mathbb{P}(65, 7, 4), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	24
246	$x^r y^4 + zy^n$ $n = 9; 11$	$t + z^3 + xy^4 + zy^n \subset \mathbb{P}(3n, n, 3n - 8, 2)$ $\mathbb{P}(n, 3n - 8, 2), \text{Diff}=(\frac{1}{2}, 0, \frac{6}{7}, 0)$	22,28
247	$x^r y^4 + zxy^n$ $n = 7; 9$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(21n - 12, 7n - 4, 3n - 8, 11)$ $\mathbb{P}(7n - 4, 3n - 8, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16,22
248	$x^r y^4 + zxy^8$	$t^2 + z^3 + x^r y^2 + zxy^4 \subset \mathbb{P}(39, 26, 8, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
249	$x^r y^4 + zx^2 y^r$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(123, 41, 13, 8)$ $\mathbb{P}(41, 13, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16
250	$zx^5 + y^{2n}$ $n = 4; 7; 8, 11; 13, 14$	$t^2 + z^3 + zx + y^2 \subset \mathbb{P}(3, 2, 4, 3)$ $\text{Diff}=(0, 0, \frac{4}{5}, \frac{n-1}{n})$	5,7,11, 15
251	$zx^5 + y^n$ $n = 11; 13; 17; 19; 23; 29$	$t + z^3 + zx + y \subset \mathbb{P}(3, 1, 2, 3)$ $\mathbb{P}(1, 2, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, \frac{n-1}{n})$	11,14,18, 20,26,30
252	$z(x^5 + ay^{2n}) + by^{3n} +$ $+cx^5 y^n, n = 3 \ a + b \neq 0; 7, 9$	$t + z^3 + z(x + ay^2) + by^3 + cxy \subset \mathbb{P}(3, 1, 2, 1)$ $\mathbb{P}(1, 2, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, \frac{n-1}{n})$	6,10
253	$z(x^5 + ay^{4n}) + by^{6n} +$ $+cx^5 y^{2n}, (n = 2, 3) \& a + b \neq 0; 4$	$t^2 + z^3 + z(x + ay^4) + by^6 + cxy^2 \subset \mathbb{P}(3, 2, 4, 1)$ $\text{Diff}=(0, 0, \frac{4}{5}, \frac{n-1}{n})$	5
254	$zx^{2+3i} f_{1-i}(x^3, y^{4n}) +$ $+ay^{2n} x^{3j} f_{2-j}(x^3, y^{4n}), n = 1 \ j = 0; 2$	$t^2 + z^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(15, 10, 4, 3)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	2,3
255	$z(x^5 + ax^2 y^{10}) + bx^3 y^{15} +$ $+cx^6 y^5 + dy^{25}$	$t + z^3 + g(z, x, y^{1/5}) \subset \mathbb{P}(15, 5, 2, 3)$ $\mathbb{P}(5, 2, 3), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{4}{5})$	6
256	$zx^5 + xy^n$ $n = 7; 9; 11; 15; 17;$ $19, 21; 23, 25$	$t + z^3 + zx^5 + xy \subset \mathbb{P}(15, 5, 2, 13)$ $\mathbb{P}(5, 2, 13), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	7,10,11, 16,18,22, 26
257	$zx^5 + xy^{2n}, n = 4, 5; 6,$ $7, 8, 9; 10, 11, 12$	$t^2 + z^3 + zx^5 + xy^2 \subset \mathbb{P}(15, 10, 4, 13)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5,9,13
258	$zx^5 + x^2 y^{2n}$ $n = 4, 5, 6, 7; 8, 9, 10$	$t^2 + z^3 + zx^5 + x^2 y^2 \subset \mathbb{P}(15, 10, 4, 11)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	7,11
259	$zx^5 + x^2 y^n, n = 7; 9;$	$t + z^3 + zx^5 + x^2 y \subset \mathbb{P}(15, 5, 2, 11)$	7,10,14,

	13; 15, 17; 19, 21	$\mathbb{P}(5, 2, 11), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	18, 22
260	$zx^5 + x^3y^n$ $n = 7; 11, 13; 17$	$t + z^3 + zx^5 + x^3y \subset \mathbb{P}(15, 5, 2, 9)$ $\mathbb{P}(5, 2, 9), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	8, 14, 18
261	$zx^5 + x^3y^{2n}$ $n = 4, 5; 7, 8$	$t^2 + z^3 + zx^5 + x^3y^2 \subset \mathbb{P}(15, 10, 4, 9)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5, 9
262	$zx^5 + x^4y^{2n}$ $n = 4, 5, 6$	$t^2 + z^3 + zx^5 + x^4y^2 \subset \mathbb{P}(15, 10, 4, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	7
263	$zx^5 + x^4y^n$ $n = 9; 11, 13$	$t + z^3 + zx^5 + x^4y \subset \mathbb{P}(15, 5, 2, 7)$ $\mathbb{P}(5, 2, 7), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	10, 14
264	$zx^5y + y^{2n}$ $n = 5; 8; 10, 11$	$t^2 + z^3 + zxy + y^{2n} \subset \mathbb{P}(3n, 2n, 4n - 3, 3)$ $\text{Diff}=(0, 0, \frac{4}{5}, 0)$	7, 11, 15
265	$zx^5y + y^n$ $n = 11; 13; 17; 23$	$t + z^3 + zxy + y^n \subset \mathbb{P}(3n, n, 2n - 3, 3)$ $\mathbb{P}(n, 2n - 3, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	14, 16, 22, 30
266	$z(x^5y + ay^{4n}) + bx^5y^{2n+1} + cy^{6n}, n = 2 \mid a \mid + \mid c \mid \neq 0, 3$	$t^2 + z^3 + g(z, x^{1/5}, y) \subset \mathbb{P}(3n, 2n, 4n - 1, 1)$ $\text{Diff}=(0, 0, \frac{4}{5}, 0)$	5
267	$z(x^5y + ay^{2n}) + bx^5y^{n+1} + cy^{3n}, n = 5 \mid a \mid + \mid c \mid \neq 0; 7$	$t + z^3 + g(z, x^{1/5}, y) \subset \mathbb{P}(3n, n, 2n - 1, 1)$ $\mathbb{P}(n, 2n - 1, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	6, 10
268	$zyx^{2+3i}f_{1-i}(x^3, y^5) + ay^4x^{3j}f_{2-j}(x^3, y^5)$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(21, 14, 5, 3)$ $\text{Diff}=\emptyset$	3
269	$zyx^{2+3i}f_{1-i}(x^3, y^7) + ay^5x^{3j}f_{2-j}(x^3, y^7)$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(57, 19, 7, 3)$ $\mathbb{P}(19, 7, 3), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
270	$zx^5y + xy^{2n+1}$ $n = 4; 5, 6, 7; 8, 9$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(15n + 6, 10n + 4, 4n - 1, 13)$ $\text{Diff}=\emptyset$	5, 9, 13
271	$zx^5y + xy^n, n = 10, 12; 14; 16, 18; 20$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(15n - 3, 5n - 1, 2n - 3, 13)$ $\mathbb{P}(5n - 1, 2n - 3, 13), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14, 18, 22, 26
272	$zx^5y + x^2y^{2n}$ $n = 4, 5, 6; 7, 8$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(15n - 3, 10n - 2, 4n - 3, 11)$ $\text{Diff}=\emptyset$	7, 11
273	$zx^5y + x^2y^n$ $n = 9; 11; 13, 15; 17$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(15n - 6, 5n - 2, 2n - 3, 11)$ $\mathbb{P}(5n - 2, 2n - 3, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10, 14, 18, 22
274	$zx^5y + x^3y^{2n+1}$ $n = 3; 5, 6$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(15n + 3, 10n + 2, 4n - 1, 9)$ $\text{Diff}=\emptyset$	5, 9
275	$zx^5y + x^3y^n$ $n = 8; 10; 14$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(15n - 9, 5n - 3, 2n - 3, 9)$ $\mathbb{P}(5n - 3, 2n - 3, 9), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10, 14, 18
276	$zx^5y + x^4y^{2n}$ $n = 4, 5$	$t^2 + z^3 + g(z, x, y) \subset \mathbb{P}(15n - 6, 10n - 4, 4n - 3, 7)$ $\text{Diff}=\emptyset$	7
277	$zx^5y + x^4y^n$ $n = 7, 9; 11$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(15n - 12, 5n - 4, 2n - 3, 7)$ $\mathbb{P}(5n - 4, 2n - 3, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10, 14
278	$zx^7 + y^n$ $n = 11, 13$	$t + z^3 + zx + y \subset \mathbb{P}(3, 1, 2, 3)$ $\mathbb{P}(1, 2, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{6}{7}, \frac{n-1}{n})$	14
279	$zx^7 + xy^{2n}$ $n = 5; 6$	$t^2 + z^3 + zx^7 + xy^2 \subset \mathbb{P}(21, 14, 4, 19)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5, 19
280	$zx^7 + xy^{11}$	$t + z^3 + zx^7 + xy \subset \mathbb{P}(21, 7, 2, 19)$ $\mathbb{P}(7, 2, 19), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{10}{11})$	24
281	$zx^7 + x^2y^n$ $n = 9; 11$	$t + z^3 + zx^7 + x^2y \subset \mathbb{P}(21, 7, 2, 17)$ $\mathbb{P}(7, 2, 17), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	10, 34
282	$zx^7 + x^2y^{10}$	$t^2 + z^3 + zx^7 + x^2y^2 \subset \mathbb{P}(21, 14, 4, 17)$ $\text{Diff}=(0, 0, 0, \frac{4}{5})$	10
283	$zx^7 + x^3y^8$	$t^2 + z^3 + zx^7 + x^3y^2 \subset \mathbb{P}(21, 14, 4, 15)$	5

		Diff=(0, 0, 0, $\frac{3}{4}$)	
284	$zx^7 + x^4y^7$	$t + z^3 + zx^7 + x^4y \subset \mathbb{P}(21, 7, 2, 13)$ $\mathbb{P}(7, 2, 13)$, Diff=($\frac{1}{2}$, 0, 0, $\frac{6}{7}$)	16
285	$zx^7 + x^4y^8$	$t^2 + z^3 + zx^7 + x^4y^2 \subset \mathbb{P}(21, 14, 4, 13)$ Diff=(0, 0, 0, $\frac{3}{4}$)	13
286	$zx^7 + x^5y^7$	$t + z^3 + zx^7 + x^5y \subset \mathbb{P}(21, 7, 2, 11)$ $\mathbb{P}(7, 2, 11)$, Diff=($\frac{1}{2}$, 0, 0, $\frac{6}{7}$)	22
287	$zx^6y + y^n$ $n = 11, 13$	$t^2 + z^3 + zx^2y + y^n \subset \mathbb{P}(3n, 2n, 2n - 3, 6)$ Diff=(0, 0, $\frac{2}{3}$, 0)	9
288	$zx^6y + y^{14}$	$t^2 + z^3 + zxy + y^{14} \subset \mathbb{P}(21, 14, 25, 3)$ Diff=(0, 0, $\frac{5}{6}$, 0)	18
289	$z(x^6y + ay^8) + bx^6y^5 +$ $+cy^{12}, a + c \neq 0$	$t^2 + z^3 + f(z, x^{1/6}, y) \subset \mathbb{P}(6, 4, 7, 1)$ Diff=(0, 0, $\frac{5}{6}$, 0)	6
290	$zx^6y + xy^n$ $n = 10, 11, 12, 13$	$t + z^3 + zx^6y + xy^n \subset \mathbb{P}(18n - 3, 6n - 1, 2n - 3, 16)$ $\mathbb{P}(6n - 1, 2n - 3, 16)$, Diff=($\frac{1}{2}$, 0, 0, 0)	14, 32
291	$zx^6y + x^2y^{2n+1}$ $n = 4, 5$	$t^2 + z^3 + zx^3y + xy^{2n+1} \subset \mathbb{P}(9n + 3, 6n + 2, 4n - 1, 7)$ Diff=(0, 0, $\frac{1}{2}$, 0)	5, 14
292	$zx^6y + x^2y^{10}$	$t^2 + z^3 + zx^6y + x^2y^{10} \subset \mathbb{P}(87, 58, 17, 14)$ Diff= $\emptyset$	7
293	$zx^6y + x^3y^n$ $n = 8, 10$	$t + z^3 + zx^2y + xy^n \subset \mathbb{P}(6n - 3, 2n - 1, 2n - 3, 4)$ $\mathbb{P}(2n - 1, 2n - 3, 4)$, Diff=($\frac{1}{2}$, 0, $\frac{2}{3}$, 0)	10, 24
294	$zx^6y + x^4y^7$	$t^2 + z^3 + zx^6y + x^4y^7 \subset \mathbb{P}(57, 38, 11, 10)$ Diff= $\emptyset$	5
295	$zx^6y + x^4y^8$	$t^2 + z^3 + zx^3y + x^2y^8 \subset \mathbb{P}(33, 22, 13, 5)$ Diff=(0, 0, $\frac{1}{2}$, 0)	10
296	$zx^6y + x^5y^7$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(111, 37, 11, 8)$ $\mathbb{P}(37, 11, 8)$, Diff=($\frac{1}{2}$, 0, 0, 0)	16
297	$zx^5y^2 + y^n$ $n = 11, 17$	$t + z^3 + zxy^2 + y^n \subset \mathbb{P}(3n, n, 2n - 6, 3)$ $\mathbb{P}(n, 2n - 6, 3)$, Diff=($\frac{1}{2}$, 0, $\frac{4}{5}$, 0)	18, 30
298	$zx^5y^2 + y^{14}$	$t^2 + z^3 + zxy^2 + y^{14} \subset \mathbb{P}(21, 14, 22, 3)$ Diff=(0, 0, $\frac{4}{5}$, 0)	11
299	$zx^5y^2 + y^{16}$	$t^2 + z^3 + zxy + y^8 \subset \mathbb{P}(12, 8, 13, 3)$ Diff=(0, 0, $\frac{4}{5}$, $\frac{1}{2}$)	15
300	$z(x^5y^2 + ay^{10}) + bx^5y^7 +$ $+cy^{15}$	$t + z^3 + g(z, x^{1/5}, y) \subset \mathbb{P}(15, 5, 8, 1)$ $\mathbb{P}(5, 8, 1)$, Diff=($\frac{1}{2}$, 0, $\frac{4}{5}$, 0)	10
301	$zx^5y^2 + xy^{4n+2}$ $n = 2, 3$	$t^2 + z^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(15n + 6, 10n + 4, 4n - 1, 13)$ Diff=(0, 0, 0, $\frac{1}{2}$)	9, 13
302	$zx^5y^2 + xy^n$ $n = 11, 13, 15$	$t + z^3 + zx^5y^2 + xy^n \subset \mathbb{P}(15n - 6, 5n - 2, 2n - 6, 13)$ $\mathbb{P}(5n, 2n - 6, 13)$, Diff=($\frac{1}{2}$, 0, 0, 0)	18, 22, 26
303	$zx^5y^2 + xy^{12}$	$t^2 + z^3 + zx^5y^2 + xy^{12} \subset \mathbb{P}(87, 58, 18, 13)$ Diff= $\emptyset$	9
304	$zx^5y^2 + x^2y^n$ $n = 9, 11, 13$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(15n - 12, 5n - 4, 2n - 6, 11)$ $\mathbb{P}(5n - 4, 2n - 6, 11)$, Diff=($\frac{1}{2}$, 0, 0, 0)	14, 18, 22
305	$zx^5y^2 + x^2y^{10}$	$t^2 + z^3 + zx^5y^2 + x^2y^{10} \subset \mathbb{P}(69, 46, 14, 11)$ Diff= $\emptyset$	7
306	$zx^5y^2 + x^2y^{12}$	$t^2 + z^3 + zx^5y + x^2y^6 \subset \mathbb{P}(42, 28, 9, 11)$ Diff=(0, 0, 0, $\frac{1}{2}$)	11
307	$zx^5y^2 + x^3y^8$	$t^2 + z^3 + zx^5y^2 + x^3y^8 \subset \mathbb{P}(51, 34, 10, 9)$	5

		Diff= $\emptyset$	
308	$zx^5y^2 + x^3y^{10}$	$t^2 + z^3 + zx^5y + x^3y^5 \subset \mathbb{P}(33, 22, 7, 9)$ Diff= $(0, 0, 0, \frac{1}{2})$	9
309	$zx^5y^2 + x^3y^{11}$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(147, 49, 16, 9)$ $\mathbb{P}(49, 16, 9)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	18
310	$zx^5y^2 + x^4y^n$ $n = 7; 9$	$t + z^3 + g(z, x, y) \subset \mathbb{P}(15n - 24, 5n - 8, 2n - 6, 7)$ $\mathbb{P}(5n - 8, 2n - 6, 7)$, Diff= $(\frac{1}{2}, 0, 0, 0)$	10, 14
311	$zx^5y^2 + x^4y^8$	$t^2 + z^3 + zx^5y + x^4y^4 \subset \mathbb{P}(24, 16, 5, 7)$ Diff= $(0, 0, 0, \frac{1}{2})$	7

2. Singularity $-t^2 + z^4 + g(z, x, y)$

1	$az^2y^n + x^5 + by^{2n}$ $n = 3; 7, 9$	$t^2 + z^2 + az^2y + x + by^2 \subset \mathbb{P}(1, 1, 2, 1)$ $\mathbb{P}^2$, Diff= $(0, \frac{1}{2}, \frac{4}{5}, \frac{n-1}{n})$	5, 10
2	$x^5 + y^n$ $n = 7; 9; 11; 13; 17, 19$	$t^2 + z^2 + x + y \subset \mathbb{P}(1, 1, 2, 2)$ $\mathbb{P}(1, 1, 2)$, Diff= $(0, \frac{1}{2}, \frac{4}{5}, \frac{n-1}{n})$	7, 10, 11, 15, 20
3	$az^2y^{2n} + bz^3y^{3n} +$ $+cy^{4n} + x^5, n = 2, 3, 4$	$t^2 + z^4 + g(z, x^{1/5}, y^{1/n}) \subset \mathbb{P}(2, 1, 4, 1)$ $\mathbb{P}(2, 1, 1)$, Diff= $(0, 0, \frac{4}{5}, \frac{n-1}{n})$	5
4	$az^2yx^if_{2-i}(x, y^2) +$ $+x^jf_{5-j}(x, y^2), i \leq 1 j \leq 2$	$t^2 + z^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(5, 5, 2, 1)$ Diff= $(0, \frac{1}{2}, 0, 0)$	2
5	$x^if_{5-i}(x, y^3)$ $i = 1, 2, 3$	$t^2 + z^2 + x^if_{5-i}(x, y) \subset \mathbb{P}(5, 5, 2, 2)$ Diff= $(0, \frac{1}{2}, 0, \frac{2}{3})$	4
6	$x^5 + xy^n$ $n = 5; 7; 9; 11; 13, 15$	$t^2 + z^2 + x^5 + xy \subset \mathbb{P}(5, 5, 2, 8)$ Diff= $(0, \frac{1}{2}, 0, \frac{n-1}{n})$	6, 7, 11, 16
7	$x^5 + ax^3y^n + bxy^{2n}$ $n = 3 \ b \neq 0; 5, 7$	$t^2 + z^2 + x^5 + ax^3y + bxy^2 \subset \mathbb{P}(5, 5, 2, 4)$ Diff= $(0, \frac{1}{2}, 0, \frac{n-1}{n})$	3, 8
8	$az^2xy^{2n} + z(bx^3y^n + cy^{5n}) +$ $+x^5 + ex^2y^{4n}, n = 1 \ c \neq 0; 2$ $ a + c + e \neq 0$	$t^2 + z^4 + g(z, x, y^{1/n}) \subset \mathbb{P}(10, 5, 4, 3)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	2, 3
9	$x^5 + x^2y^n$ $n = 5, 7; 11$	$t^2 + z^2 + x^5 + x^2y \subset \mathbb{P}(5, 5, 2, 6)$ Diff= $(0, \frac{1}{2}, 0, \frac{n-1}{n})$	7, 12
10	$x^5 + az^2xy^5 + bx^2y^{10}$	$t^2 + z^2 + x^5 + azxy + bx^2y^2 \subset \mathbb{P}(5, 5, 2, 3)$ Diff= $(0, \frac{1}{2}, 0, \frac{4}{5})$	6
11	$x^5 + zy^n$ $n = 4; 7; 8; 11; 13, 14$	$t^2 + z^4 + x + zy \subset \mathbb{P}(2, 1, 4, 3)$ $\mathbb{P}(2, 1, 3)$, Diff= $(0, 0, \frac{4}{5}, \frac{n-1}{n})$	5, 7, 10, 11, 15
12	$x^5 + zxy^n, n = 4,$ $5, 6; 7; 8, 9, 10$	$t^2 + z^4 + x^5 + zxy \subset \mathbb{P}(10, 5, 4, 11)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	6, 7, 11
13	$x^5 + zx^2y^n$ $n = 4, 5, 6$	$t^2 + z^4 + x^5 + zx^2y \subset \mathbb{P}(10, 5, 4, 7)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	7
14	$\underline{a}z^2x^{i+1}f_{2-i}(x, y) +$ $+ \underline{b}x^jf_{6-j}(x, y), j \leq 1$	$t^2 + z^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(3, 3, 1, 1)$ Diff= $(0, \frac{1}{2}, 0, 0)$	2
15	$az^2x^3 + bx^6 + y^n$ $n = 7; 11$	$t^2 + z^2 + azx + bx^2 + y \subset \mathbb{P}(1, 1, 1, 2)$ $\mathbb{P}^2$, Diff= $(0, \frac{1}{2}, \frac{2}{3}, \frac{n-1}{n})$	7, 12
16	$z^3y^2 + z^2(x^3 + y^4) + zy^2(x^3 +$ $+y^4) + x^6 + x^3y^4 + y^8$	$t^2 + z^4 + g(z, x^{1/3}, y^{1/2}) \subset \mathbb{P}(2, 1, 2, 1)$ Diff= $(0, 0, \frac{2}{3}, \frac{1}{2})$	3
17	$\underline{a}z^2x^{2i+1}f_{1-i}(x^2, y^3) +$ $\underline{b}x^{2j}f_{3-j}(x^2, y^3)$	$t^2 + z^2 + g(z^{1/2}, x, y^{1/3}) \subset \mathbb{P}(3, 3, 1, 2)$ Diff= $(0, \frac{1}{2}, 0, \frac{2}{3})$	4
18	$\underline{a}z^2x^{3i}f_{1-i}(x^3, y^5) +$	$t^2 + z^2 + g(z^{1/2}, x^{1/3}, y^{1/5}) \subset \mathbb{P}(1, 1, 1, 1)$	6

	$bx^{3j}f_{2-j}(x^3, y^5)$	$\text{Diff}=(0, \frac{1}{2}, \frac{2}{3}, \frac{4}{5})$	
19	$az^2x^3 + bx^2y^n + cx^6 + dxy^{2n}, n = 3 \& d \neq 0; n = 4 \& b + d \neq 0$	$t^2 + z^4 + g(z, x, y^{1/n}) \subset \mathbb{P}(6, 3, 2, 5)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	3,5
20	$az^2x^3 + bx^6 + xy^n$ $n = 7; 9$	$t^2 + z^2 + g(z^{1/2}, x, y^{1/n}) \subset \mathbb{P}(3, 3, 1, 5)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{n-1}{n})$	7,10
21	$az^2x^3 + bx^6 + x^2y^n$ $n = 5; 7$	$t^2 + z^2 + g(z^{1/2}, x, y^{1/n}) \subset \mathbb{P}(3, 3, 1, 4)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{n-1}{n})$	5,8
22	$az^2x^3 + zy^n + bx^6$ $n = 5; 7; 8$	$t^2 + z^4 + g(z, x^{1/3}, y^{1/n}) \subset \mathbb{P}(2, 1, 2, 3)$ $\text{Diff}=(0, 0, \frac{2}{3}, \frac{n-1}{n})$	5,7,9
23	$az^2x^3 + zxy^n + cx^6$ $n = 4; 5; 6$	$t^2 + z^4 + g(z, x, y^{1/n}) \subset \mathbb{P}(6, 3, 2, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	4,5,7
24	$x^5y + y^n$ $n = 7; 9; 13; 15$	$t^2 + z^2 + xy + y^n \subset \mathbb{P}(n, n, 2n - 2, 2)$ $\text{Diff}=(0, \frac{1}{2}, \frac{4}{5}, 0)$	8,11,16, 20
25	$az^2y^{2n} + bzy^{3n} + x^5y + dy^{4n}, n = 2, 3$	$t^2 + z^4 + g(z, x^{1/5}, y) \subset \mathbb{P}(2n, n, 4n - 1, 1)$ $\text{Diff}=(0, 0, \frac{4}{5}, 0)$	5
26	$az^2y^n + x^5y + by^{2n}$ $n = 5; 7$	$t^2 + z^2 + azy^n + xy + by^{2n} \subset \mathbb{P}(n, n, 2n - 1, 1)$ $\text{Diff}=(0, \frac{1}{2}, \frac{4}{5}, 0)$	6,10
27	$yx^i f_{5-i}(x, y^2)$ $i \leq 3$	$t^2 + z^2 + g(x, y) \subset \mathbb{P}(11, 11, 4, 2)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	4
28	$x^5y + xy^n$ $n = 6; 8; 10; 12$	$t^2 + z^2 + g(x, y) \subset \mathbb{P}(5n - 1, 5n - 1, 2n - 2, 8)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	7,11,12, 16
29	$x^5y + ax^3y^{n+1} + bxy^{2n+1}$ $n = 3 \ b \neq 0; 5$	$t^2 + z^2 + g(x, y) \subset \mathbb{P}(5n + 2, 5n + 2, 2n, 4)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	4,8
30	$x^5y + x^2y^n$ $n = 5; 9$	$t^2 + z^2 + g(x, y) \subset \mathbb{P}(5n - 2, 5n - 2, 2n - 2, 6)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	7,12
31	$az^2xy^3 + z(bx^3y^2 + cy^7) + x^5y + ex^2y^6, a + c + e \neq 0$	$t^2 + z^4 + g(z, x, y) \subset \mathbb{P}(14, 7, 5, 3)$ $\text{Diff} = \emptyset$	3
32	$az^2xy^4 + x^5y + bx^2y^8$	$t^2 + z^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(19, 19, 7, 3)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	6
33	$zy^n + x^5y$ $n = 5; 8; 10, 11$	$t^2 + z^4 + g(z, x^{1/5}, y) \subset \mathbb{P}(2n, n, 4n - 3, 3)$ $\text{Diff}=(0, 0, \frac{4}{5}, 0)$	6,10,15
34	$zxy^n + x^5y$ $n = 4, 5; 6; 7, 8$	$t^2 + z^4 + g(z, x, y) \subset \mathbb{P}(10n - 2, 5n - 1, 4n - 3, 11)$ $\text{Diff} = \emptyset$	6,7,11
35	$zx^2y^n + x^5y$ $n = 4, 5$	$t^2 + z^4 + g(z, x, y) \subset \mathbb{P}(10n - 4, 5n - 2, 4n - 3, 7)$ $\text{Diff} = \emptyset$	7
36	$x^i y^j f_{7-i-j}(x, y)$ $i \leq 2 \& j \leq 2$	$t^2 + z^2 + g(x, y) \subset \mathbb{P}(7, 7, 2, 2)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	4
37	$z^i y^{2j} (z + y^2)^l f_{4-i-j-l}(z, y^2) + x^7, i \leq 2 \& j \leq 1 \& l \leq 2$	$t^2 + z^4 + g(z, x^{1/7}, y^{1/2}) \subset \mathbb{P}(2, 1, 4, 1)$ $\mathbb{P}(2, 1, 1), \text{Diff}=(0, 0, \frac{6}{7}, \frac{1}{2})$	7
38	$x^7 + y^9$	$t^2 + z^2 + x + y \subset \mathbb{P}(1, 1, 2, 2)$ $\mathbb{P}(1, 1, 2), \text{Diff}=(0, \frac{1}{2}, \frac{6}{7}, \frac{8}{9})$	28
39	$x^7 + xy^7$	$t^2 + z^2 + x^7 + xy \subset \mathbb{P}(7, 7, 2, 12)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{6}{7})$	8
40	$az^2xy^3 + x^7 + bx^2y^6$	$t^2 + z^2 + g(z^{1/2}, x, y^{1/3}) \subset \mathbb{P}(7, 7, 2, 5)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{2}{3})$	10
41	$x^7 + x^3y^5$	$t^2 + z^2 + x^7 + x^3y \subset \mathbb{P}(7, 7, 2, 8)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{4}{5})$	16

42	$zxy^5 + x^7$	$t^2 + z^4 + x^7 + zxy \subset \mathbb{P}(14, 7, 4, 17)$ Diff= $(0, 0, 0, \frac{4}{5})$	10
43	$zx^2y^4 + x^7$	$t^2 + z^4 + x^7 + zx^2y \subset \mathbb{P}(14, 7, 4, 13)$ Diff= $(0, 0, 0, \frac{3}{4})$	13
44	$az^2y^4 + bzy^6 +$ $+x^6y + cy^8$	$t^2 + z^4 + g(z, x^{1/6}, y) \subset \mathbb{P}(4, 2, 7, 1)$ $\mathbb{P}(2, 1, 1)$, Diff= $(0, 0, \frac{5}{6}, 0)$	6
45	$x^6y + ax^3y^5 + by^9$	$t^2 + z^2 + g(x^{1/3}, y) \subset \mathbb{P}(9, 9, 8, 2)$ Diff= $(0, \frac{1}{2}, \frac{2}{3}, 0)$	12
46	$x^6y + xy^n$ $n = 7; 8$	$t^2 + z^2 + x^6y + xy^n \subset \mathbb{P}(6n - 1, 6n - 1, 2n - 2, 10)$ Diff= $(0, \frac{1}{2}, 0, 0)$	8,20
47	$az^2xy^3 + x^6y + bx^2y^6$	$t^2 + z^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(17, 17, 5, 4)$ Diff= $(0, \frac{1}{2}, 0, 0)$	8
48	$zy^7 + x^6y$	$t^2 + z^4 + zy^7 + xy \subset \mathbb{P}(14, 7, 25, 3)$ Diff= $(0, 0, \frac{5}{6}, 0)$	18
49	$zxy^5 + x^6y$	$t^2 + z^4 + zxy^5 + x^6y \subset \mathbb{P}(58, 29, 17, 14)$ Diff = $\emptyset$	7
50	$zx^2y^4 + x^6y$	$t^2 + z^4 + zxy^4 + x^3y \subset \mathbb{P}(22, 11, 13, 5)$ Diff= $(0, 0, \frac{1}{2}, 0)$	10
51	$z^i y^{2j} f_{4-i-j}(z, y^2) +$ $+(zx^5 x^5 y^2)$ $i \leq 2 \& j \leq 2 \& (j \leq 1 j = 0)$	$t^2 + z^4 + g(z, x^{1/5}, y^{1/2}) \subset \mathbb{P}(2, 1, 3, 1)$ Diff= $(0, 0, \frac{4}{5}, \frac{1}{2})$	5
52	$x^5y^2 + y^n$ $n = 9; 11$	$t^2 + z^2 + xy^2 + y^n \subset \mathbb{P}(n, n, 2n - 4, 2)$ Diff= $(0, \frac{1}{2}, \frac{4}{5}, 0)$	15,20
53	$az^2y^5 + x^5y^2 + by^{10}$	$t^2 + z^2 + azy^5 + xy^2 + by^{10} \subset \mathbb{P}(5, 5, 8, 1)$ Diff= $(0, \frac{1}{2}, \frac{4}{5}, 0)$	10
54	$x^5y^2 + xy^n$ $n = 7; 9$	$t^2 + z^2 + x^5y^2 + xy^n \subset \mathbb{P}(5n - 2, 5n - 2, 2n - 4, 8)$ Diff= $(0, \frac{1}{2}, 0, 0)$	11,16
55	$x^5y^2 + ax^3y^5 + bxy^8$	$t^2 + z^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(19, 19, 6, 4)$ Diff= $(0, \frac{1}{2}, 0, 0)$	8
56	$az^2xy^3 + x^5y^2 + bx^2y^6$	$t^2 + z^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(13, 13, 4, 3)$ Diff= $(0, \frac{1}{2}, 0, 0)$	6
57	$x^5y^2 + x^2y^7$	$t^2 + z^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(31, 31, 10, 6)$ Diff= $(0, \frac{1}{2}, 0, 0)$	12
58	$zy^7 + x^5y^2$	$t^2 + z^4 + zy^7 + xy^2 \subset \mathbb{P}(14, 7, 22, 3)$ Diff= $(0, 0, \frac{4}{5}, 0)$	11
59	$zy^8 + x^5y^2$	$t^2 + z^4 + zy^4 + xy \subset \mathbb{P}(8, 4, 13, 3)$ Diff= $(0, 0, \frac{4}{5}, \frac{1}{2})$	15
60	$zxy^5 + x^5y^2$	$t^2 + z^4 + g(z, x, y) \subset \mathbb{P}(46, 23, 14, 11)$ Diff = $\emptyset$	7
61	$zxy^6 + x^5y^2$	$t^2 + z^4 + zxy^3 + x^5y \subset \mathbb{P}(28, 14, 9, 11)$ Diff= $(0, 0, 0, \frac{1}{2})$	11
62	$zx^2y^4 + x^5y^2$	$t^2 + z^4 + zx^2y^2 + x^5y \subset \mathbb{P}(16, 8, 5, 7)$ Diff= $(0, 0, 0, \frac{1}{2})$	7
63	$az^2y^n + zx^4 + by^{2n}$ $n = 3 \ b \neq 0; 5; 7$	$t^2 + z^4 + az^2y + zx + by^2 \subset \mathbb{P}(2, 1, 3, 2)$ Diff= $(0, 0, \frac{3}{4}, \frac{n-1}{n})$	4,5,8
64	$zx^4 + y^n$ $n = 7; 9; 11; 13; 15$	$t^2 + z^4 + zx + y \subset \mathbb{P}(2, 1, 3, 4)$ $\mathbb{P}(2, 1, 3)$, Diff= $(0, 0, \frac{3}{4}, \frac{n-1}{n})$	7,9,12 13,16
65	$az^2yx^{2i}f_{1-i}(x^2, y^3) +$	$t^2 + z^4 + g(z, x^{1/2}, y) \subset \mathbb{P}(4, 2, 3, 1)$	2

	$+zx^{2j}f_{2-j}(x^2, y^3) + by^2x^{2l}.$ $f_{2-l}(x^2, y^3), i=0 j=0 l \leq 1$	Diff=(0, 0, $\frac{1}{2}$, 0)	
66	$az^2y^6 + z(x^4 + by^9) +$ $+cx^4y^3 + dy^{12}, a + b + d \neq 0$	$t^2 + z^4 + g(z, x^{1/4}, y^{1/3}) \subset \mathbb{P}(2, 1, 3, 1)$ Diff=(0, 0, $\frac{3}{4}, \frac{2}{3}$)	4
67	$zx^4 + xy^n, n=5; 6;$ $7; 8, 9; 10; 11, 12$	$t^2 + z^4 + zx^4 + xy \subset \mathbb{P}(8, 4, 3, 13)$ Diff=(0, 0, 0, $\frac{n-1}{n}$)	5,6,7, 9,10,13
68	$az^2xy^n + zx^4 + bx^2y^{2n}$ $n=3 \ b \neq 0, 4$	$t^2 + z^4 + g(z, x, y^{1/n}) \subset \mathbb{P}(8, 4, 3, 5)$ Diff=(0, 0, 0, $\frac{n-1}{n}$)	5
69	$zx^4 + x^2y^n$ $n=7; 9$	$t^2 + z^4 + zx^2 + xy \subset \mathbb{P}(4, 2, 3, 5)$ Diff=(0, 0, $\frac{1}{2}, \frac{n-1}{n}$)	7,10
70	$zx^4 + x^3y^n$ $n=5, 6$	$t^2 + z^4 + zx^4 + x^3y \subset \mathbb{P}(8, 4, 3, 7)$ Diff=(0, 0, 0, $\frac{n-1}{n}$)	7
71	$zx^5 + y^n$ $n=7; 9$	$t^2 + z^4 + zx + y \subset \mathbb{P}(2, 1, 3, 4)$ $\mathbb{P}(2, 1, 3), \text{Diff}=(0, 0, \frac{4}{5}, \frac{n-1}{n})$	7,10
72	$zx^5 + xy^n$ $n=6; 7; 8$	$t^2 + z^4 + zx^5 + xy \subset \mathbb{P}(10, 5, 3, 17)$ Diff=(0, 0, 0, $\frac{n-1}{n}$)	6,7,17
73	$zx^5 + x^2y^5$	$t^2 + z^4 + zx^5 + x^2y \subset \mathbb{P}(10, 5, 3, 14)$ Diff=(0, 0, 0, $\frac{4}{5}$)	7
74	$az^2xy^3 + zx^5 + bx^2y^6$	$t^2 + z^4 + g(z, x, y^{1/3}) \subset \mathbb{P}(10, 5, 3, 7)$ Diff=(0, 0, 0, $\frac{2}{3}$)	7
75	$zx^5 + x^3y^5$	$t^2 + z^4 + zx^5 + x^3y \subset \mathbb{P}(10, 5, 3, 11)$ Diff=(0, 0, 0, $\frac{4}{5}$)	11
76	$zx^4y + y^n$ $n=7; 9; 11$	$t^2 + z^4 + zxy + y^n \subset \mathbb{P}(2n, n, 3n-4, 4)$ Diff=(0, 0, $\frac{3}{4}$, 0)	9,12,16
77	$az^2y^4 + z(x^4y + by^6) +$ $+cx^4y^3 + dy^8, a + b + d \neq 0$	$t^2 + z^4 + g(z, x^{1/4}, y) \subset \mathbb{P}(4, 2, 5, 1)$ Diff=(0, 0, $\frac{3}{4}$, 0)	4
78	$az^2y^5 + zx^4y + by^{10}$	$t^2 + z^4 + g(z, x^{1/4}, y) \subset \mathbb{P}(10, 5, 13, 2)$ Diff=(0, 0, $\frac{3}{4}$, 0)	8
79	$zx^4y + xy^n$ $n=6; 7, 8; 9$	$t^2 + z^4 + g(z, x, y) \subset \mathbb{P}(8n-2, 4n-1, 3n-4, 13)$ Diff = $\emptyset$	7,10,13
80	$zx^4y + x^2y^n$ $n=5; 7$	$t^2 + z^4 + zx^2y + xy^n \subset \mathbb{P}(4n-2, 2n-1, 3n-4, 5)$ Diff=(0, 0, $\frac{1}{2}$, 0)	6,10
81	$az^2xy^3 + zx^4y + bx^2y^6$	$t^2 + z^4 + g(z, x, y) \subset \mathbb{P}(22, 11, 7, 5)$ Diff = $\emptyset$	5
82	$zx^4y + x^3y^5$	$t^2 + z^4 + g(z, x, y) \subset \mathbb{P}(34, 17, 11, 7)$ Diff = $\emptyset$	7

(1) There are two cases. A). $t^2 + z^4 + z^2(ax^3 + by^4) + zy^2(cx^3 + dy^4) + ex^6 + kx^3y^4 + ly^8$. Then the singularity is exceptional if the following condition is satisfied: $|b| + |d| + |l| \neq 0$ & $(e \neq 0 || (a \neq 0 \& |c| + |k| + |l| \neq 0))$.

B). $t^2 + z^3y^2 + z^2x^3 + zy^2(ax^3 + by^4) + cx^6 + dx^3y^4 + ey^8$. The exceptionality condition is $|ab| + |c| + |d| + |e| \neq 0$ & $|a| + |b| + |d| + |e| \neq 0$.

3. Singularity – $t^2 + z^3x + g(z, x, y)$

1	$azx^2y^{2n} + x^5 + by^{6n}$ $n=2 \ b \neq 0; 3, 4$	$t^2 + z^3x + ax^2y^2 + x^5 + by^6 \subset \mathbb{P}(15, 8, 6, 5)$ Diff=(0, 0, 0, $\frac{n-1}{n}$)	4,5
2	$azx^2y^n + x^5 + by^{3n}$ $n=3 \ b \neq 0; 7, 9$	$t + z^3x + ax^2y + x^5 + by^3 \subset \mathbb{P}(15, 4, 3, 5)$ $\mathbb{P}(4, 3, 5), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	4,10
3	$x^5 + y^n$ $n=7; 11; 13; 17; 19;$	$t + zx + x^5 + y \subset \mathbb{P}(5, 4, 1, 5)$ $\mathbb{P}(4, 1, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, \frac{n-1}{n})$	7,12,14, 18,22,24,

	23;29		30
4	$x^5 + y^{2n}$ $n = 4, 7; 8; 11, 13, 14$	$t^2 + zx + x^5 + y^2 \subset \mathbb{P}(5, 8, 2, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{n-1}{n})$	7,9,15
5	$x^i(x+y^2)^j f_{5-i-j}(x, y^2)$ $i \leq 1 \ \& \ j \leq 3$	$t^2 + zx + g(x, y) \subset \mathbb{P}(5, 8, 2, 1)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	3
6 (1)	$z^2 y f_2(x, y^3) + z y^2 f_3(x, y^3) + f_5(x, y^3)$	$t + z^3 x + g(z, x, y) \subset \mathbb{P}(15, 4, 3, 1)$ $\mathbb{P}(4, 3, 1), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	2
7	$x^i f_{5-i}(x, y^4)$ $i \leq 3$	$t^2 + zx + g(x, y^{1/2}) \subset \mathbb{P}(5, 8, 2, 1)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{1}{2})$	3
8	$x^i f_{5-i}(x, y^5)$ $i \leq 3$	$t + zx + g(x, y^{1/5}) \subset \mathbb{P}(5, 4, 1, 1)$ $\mathbb{P}(4, 1, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, \frac{4}{5})$	6
9	$x^5 + x^2 y^{2n}$ $n = 4, 5, 7, 8$	$t^2 + zx + x^5 + x^2 y^2 \subset \mathbb{P}(5, 8, 2, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{n-1}{n})$	9
10	$x^5 + x^2 y^n$ $n = 7; 11; 13, 17$	$t + zx + x^5 + x^2 y \subset \mathbb{P}(5, 4, 1, 3)$ $\mathbb{P}(4, 1, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, \frac{n-1}{n})$	10,12,18
11	$x^5 + z y^{2n}$ $n = 3; 4, 5; 6 - 10$	$t^2 + z^3 x + x^5 + z y^2 \subset \mathbb{P}(15, 8, 6, 11)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	3,5,11
12	$x^5 + z y^n, n = 5; 7;$ $9; 13; 15; 17, 19, 21$	$t + z^3 x + x^5 + z y \subset \mathbb{P}(15, 4, 3, 11)$ $\mathbb{P}(4, 3, 11), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	6,7,10 14,16,22
13	$x^5 + z^2 y^{2n}$ $n = 2, 3; 4, 5, 6$	$t^2 + z^3 x + x^5 + z^2 y^2 \subset \mathbb{P}(15, 8, 6, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	6,7
14	$x^5 + z^2 y^n$ $n = 5; 9, 11, 13$	$t + z^3 x + x^5 + z^2 y \subset \mathbb{P}(15, 4, 3, 7)$ $\mathbb{P}(4, 3, 7), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	6,14
15	$x^i(x+y^n)^j f_{6-i-j}(x, y^n)$ $n = 1 \ \& \ i = 0 \ \& \ j \leq 1,$ $n = 2 \ \& \ i \leq 2 \ \& \ j \leq 4$	$t^2 + zx + g(x, y) \subset \mathbb{P}(3, 5, 1, 1)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{n-1}{n})$	3
16	$x^6 + y^n$ $n = 7; 11; 13; 17$	$t^2 + zx + x^6 + y \subset \mathbb{P}(3, 5, 1, 6)$ $\mathbb{P}(3, 5, 1), \text{Diff}=(0, \frac{2}{3}, 0, \frac{n-1}{n})$	7,12,13 18
17	$x^{3i} f_{2-i}(x^3, y^n)$ $(n = 4; 5) \ \& \ i = 0; (7, 8) \ \& \ i \leq 1$	$t^2 + zx + g(x, y^{1/n}) \subset \mathbb{P}(3, 5, 1, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{n-1}{n})$	4,6,9
18	$a z^2 y x^{2i} f_{1-i}(x^2, y^3) +$ $+ b z y^2 x^{2j+1} f_{1-j}(x^2, y^3) +$ $+ x^{2l} f_{3-l}(x^2, y^3), i = 0 j = 0 l = 0$	$t^2 + z^3 x + g(z, x, y) \subset \mathbb{P}(9, 5, 3, 2)$ $\text{Diff} = \emptyset$	2
19	$x^{2i} f_{3-i}(x^2, y^5)$ $i \leq 1$	$t^2 + zx + g(x, y^{1/5}) \subset \mathbb{P}(3, 5, 1, 2)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{4}{5})$	6
20	$x^6 + x^2 y^n$ $n = 7; 11$	$t^2 + zx + x^6 + x^2 y \subset \mathbb{P}(3, 5, 1, 4)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{n-1}{n})$	7,12
21	$a z^2 y^6 + b z x^3 y^3 + x^6 +$ $+ c x^2 y^9, a + c \neq 0$	$t^2 + z^3 x + g(z, x, y^{1/3}) \subset \mathbb{P}(9, 5, 3, 4)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	4
22	$x^6 + z y^n, n = 5;$ $6, 7; 8; 9, 10; 11, 12$	$t^2 + z^3 x + x^6 + z y \subset \mathbb{P}(9, 5, 3, 13)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5,7,8, 10,13
23	$x^6 + z x^2 y^n$ $n = 5, 6$	$t^2 + z^3 x + x^6 + z x^2 y \subset \mathbb{P}(9, 5, 3, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	7
24	$x^6 + z^2 y^n$ $n = 5; 7$	$t^2 + z^3 x + x^6 + z^2 y \subset \mathbb{P}(9, 5, 3, 8)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5,8
25	$x^5 y + y^n$ $n = 7; 9; 13; 15; 19; 25$	$t + zx + x^5 y + y^n \subset \mathbb{P}(5n, 4n+1, n-1, 5)$ $\mathbb{P}(4n+1, n-1, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	8,10,15 18,22,30
26	$x^5 y + y^{2n}$	$t^2 + zx + x^5 y + y^{2n} \subset \mathbb{P}(5n, 8n+1, 2n-1, 5)$	7,15

	$n = 5, 6; 9, 11, 12$	$\text{Diff}=(0, \frac{2}{3}, 0, 0)$	
27	$ax^2y^{2n+1} + x^5y + by^{6n+2}$ $n = 1 \ b \neq 0; 2, 3$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(15n + 5, 8n + 3, 6n + 1, 5)$ $\text{Diff} = \emptyset$	4,5
28	$ax^2y^n + x^5y + by^{3n-1}$ $n = 6, 8$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(15n - 5, 4n - 1, 3n - 2, 5)$ $\mathbb{P}(4n - 1, 3n - 2, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
29	$zf_4(x, y^2) + z^2y^5 +$ (2) $+yf_5(x, y^2)$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(11, 3, 2, 1)$ $\mathbb{P}(3, 2, 1), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	2
30	$yx^i f_{5-i}(x, y^3)$ $i \leq 3$	$t^2 + zx + g(x, y) \subset \mathbb{P}(8, 13, 3, 1)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	3
31	$yx^i f_{5-i}(x, y^4)$ $i \leq 3$	$t + zx + g(x, y) \subset \mathbb{P}(21, 17, 4, 1)$ $\mathbb{P}(17, 4, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
32	$x^5y + x^2y^n$ $n = 9; 11; 15$	$t + zx + x^5y + x^2y^n \subset \mathbb{P}(5n - 2, 4n - 1, n - 1, 3)$ $\mathbb{P}(4n - 1, n - 1, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,12,18
33	$x^5y + x^2y^{2n}$ $n = 4, 6, 7$	$t^2 + zx + x^5y + x^2y^{2n} \subset \mathbb{P}(5n - 1, 8n - 1, 2n - 1, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
34	$x^5y + zy^{2n+1}$ $n = 2; 3, 4; 5 - 8$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(15n + 8, 8n + 5, 6n + 1, 11)$ $\text{Diff} = \emptyset$	3,5,11
35	$x^5y + zy^n, n = 6;$ 10, 12; 14; 16, 18	$t + z^3x + x^5y + zy^n \subset \mathbb{P}(15n + 1, 4n + 1, 3n - 2, 11)$ $\mathbb{P}(4n + 1, 3n - 2, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	7,14,16, 22
36	$x^5y + z^2y^n$ $n = 3; 7; 9, 11$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(15n + 2, 4n + 1, 3n - 1, 7)$ $\mathbb{P}(4n + 1, 3n - 1, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,8,14
37	$x^5y + z^2y^{2n}$ $n = 2; 3, 4, 5$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(15n + 1, 8n + 1, 6n - 1, 7)$ $\text{Diff} = \emptyset$	6,7
38	$zf_5(x, y) + f_7(x, y)$ (3)	$t + z^3x + g(z, x, y) \subset \mathbb{P}(7, 2, 1, 1)$ $\mathbb{P}(2, 1, 1), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	2
39	$x^3z^i f_{2-i}(z, x^2) + y^{2n}$ $n = 4, 5; 6$	$t^2 + z^3x + g(z, x, y^{1/n}) \subset \mathbb{P}(7, 4, 2, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5,7
40	$x^3z^i f_{2-i}(z, x^2) + y^n$ $n = 9; 11; 13$	$t + z^3x + g(z, x, y^{1/n}) \subset \mathbb{P}(7, 2, 1, 7)$ $\mathbb{P}(2, 1, 7), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	10,12,14
41	$x^3z^i f_{2-i}(z, x^2) + az^2y^4 +$ $+bx^2y^4 + cx^4y^4 + dxy^8$ $ a + d \neq 0$	$t^2 + z^3x + g(z, x, y^{1/2}) \subset \mathbb{P}(7, 4, 2, 3)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	3
42	$x^3z^i f_{2-i}(z, x^2) + az^2y^5 +$ $+bx^2y^5 + cx^4y^5 + dxy^{10}$ $ a + b + d \neq 0$	$t + z^3x + g(z, x, y^{1/5}) \subset \mathbb{P}(7, 2, 1, 3)$ $\mathbb{P}(2, 1, 3), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{4}{5})$	6
43	$x^3z^i f_{2-i}(z, x^2) + azy^{2n} +$ $+bx^2y^{2n}, n = 3 \ a \neq 0; 4$	$t^2 + z^3x + g(z, x, y^{1/n}) \subset \mathbb{P}(7, 4, 2, 5)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	3,5
44	$x^3z^i f_{2-i}(z, x^2) + azy^n +$ $+bx^2y^n, n = 7; 9$	$t + z^3x + g(z, x, y^{1/n}) \subset \mathbb{P}(7, 2, 1, 5)$ $\mathbb{P}(2, 1, 5), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	8,10
45	$x^6y + y^n$ $n = 8, 12, 14$	$t^2 + zx + x^6y + y^n \subset \mathbb{P}(3n, 5n + 1, n - 1, 6)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
46	$x^6y + bx^3y^{n+1} + cy^{2n+1}$ $(c \neq 0 \ \& \ (n = 4, 5)); 7$	$t + zx + g(x, y) \subset \mathbb{P}(6n + 3, 5n + 3, n, 3)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6,18
47	$yx^{2i} f_{3-i}(x^2, y^3)$ $i \leq 1$	$t^2 + zx + g(x, y) \subset \mathbb{P}(10, 17, 3, 2)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	3
48	$yx^i(x + y^2)^j f_{6-i-j}(x, y^2)$ $i \leq 3 \ \& \ j \leq 5$	$t + zx + g(x, y) \subset \mathbb{P}(13, 11, 2, 1)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
49	$x^6y + x^2y^{10}$	$t^2 + zx + x^6y + x^2y^{10} \subset \mathbb{P}(29, 49, 9, 4)$	12

		Diff=(0, $\frac{2}{3}$, 0, 0)	
50	$az^2y^5 + bzx^3y^3 + x^6y + cx^2y^8, a + c \neq 0$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(23, 13, 7, 4)$ Diff = $\emptyset$	4
51	$x^6y + zy^n$ $n = 6, 7, 8, 9; 10, 11$	$t + z^3x + x^6y + zy^n \subset \mathbb{P}(18n + 1, 5n + 1, 3n - 2, 13)$ $\mathbb{P}(5n + 1, 3n - 2, 13)$, Diff=($\frac{1}{2}$, 0, 0, 0)	10, 26
52	$x^6y + zx^2y^n$ $n = 5, 6$	$t + z^3x + x^6y + zx^2y^n \subset \mathbb{P}(18n - 5, 5n - 1, 3n - 2, 7)$ $\mathbb{P}(5n - 1, 3n - 2, 7)$, Diff=($\frac{1}{2}$, 0, 0, 0)	14
53	$x^5y^2 + y^{4n}$ $n = 2; 5$	$t^2 + zx + x^5y + y^{2n} \subset \mathbb{P}(5n, 8n + 1, 2n - 1, 5)$ Diff=(0, $\frac{2}{3}$, 0, $\frac{1}{2}$)	7, 15
54	$x^5y^2 + y^{2n}$ $n = 7; 9$	$t^2 + zx + g(x, y) \subset \mathbb{P}(5n, 8n + 2, 2n - 2, 5)$ Diff=(0, $\frac{2}{3}$, 0, 0)	9, 15
55	$x^5y^2 + y^n$ $n = 9; 11; 15; 21$	$t + zx + g(x, y) \subset \mathbb{P}(5n, 4n + 2, n - 2, 5)$ $\mathbb{P}(4n + 2, n - 2, 5)$ Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	12, 15, 22, 30
56	$azx^2y^4 + x^5y^2 + by^{10}$ $b \neq 0$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(25, 14, 8, 5)$ Diff = $\emptyset$	4
57	$azx^2y^n + x^5y^2 + by^{3n-2}$ $n = 5, 7$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(15n - 10, 4n - 2, 3n - 4, 5)$ $\mathbb{P}(4n - 2, 3n - 4, 5)$, Diff=($\frac{1}{2}$, 0, 0, 0)	10
58	$azx^2y^6 + x^5y^2 + by^{16}$	$t^2 + z^3x + g(z, x, y^{1/2}) \subset \mathbb{P}(20, 11, 7, 5)$ Diff=(0, 0, 0, $\frac{1}{2}$)	5
59	$y^2x^if_{5-i}(x, y^3)$ $i \leq 3$	$t + zx + g(x, y) \subset \mathbb{P}(17, 14, 3, 1)$ $\mathbb{P}(14, 3, 1)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	6
60	$x^5y^2 + x^2y^{10}$	$t^2 + zx + x^5y^2 + x^2y^{10} \subset \mathbb{P}(23, 38, 8, 3)$ Diff=(0, $\frac{2}{3}$, 0, 0)	9
61	$x^5y^2 + x^2y^n$ $n = 7; 9; 13$	$t + zx + x^5y^2 + x^2y^n \subset \mathbb{P}(5n - 4, 4n - 2, n - 2, 3)$ $\mathbb{P}(4n - 2, n - 2, 3)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	10, 12, 18
62	$x^5y^2 + x^2y^{12}$	$t^2 + zx + x^5y + x^2y^6 \subset \mathbb{P}(14, 23, 5, 3)$ Diff=(0, $\frac{2}{3}$, 0, $\frac{1}{2}$)	9
63	$x^5y^2 + zy^{4n+2}$ $n = 1; 2, 3$	$t^2 + z^3x + g(z, x, y^{1/2}) \subset \mathbb{P}(15n + 8, 8n + 5, 6n + 1, 11)$ Diff=(0, 0, 0, $\frac{1}{2}$)	5, 11
64	$x^5y^2 + zy^n$ $n = 7; 9, 11; 13, 15$	$t + z^3x + x^5y^2 + zy^n \subset \mathbb{P}(15n + 2, 4n + 2, 3n - 4, 11)$ $\mathbb{P}(4n + 2, 3n - 4, 11)$, Diff=($\frac{1}{2}$, 0, 0, 0)	10, 16, 22
65	$x^5y^2 + zy^{4n}$ $n = 2; 3$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(30n + 1, 16n + 2, 12n - 4, 11)$ Diff = $\emptyset$	5, 11
66	$x^5y^2 + z^2y^6$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(47, 26, 16, 7)$ Diff = $\emptyset$	7
67	$x^5y^2 + z^2y^{4n}$ $n = 1; 2$	$t^2 + z^3x + g(z, x, y^{1/2}) \subset \mathbb{P}(15n + 1, 8n + 1, 6n - 1, 7)$ Diff=(0, 0, 0, $\frac{1}{2}$)	6, 7
68	$x^5y^2 + z^2y^n$ $n = 5; 7, 9$	$t + z^3x + x^5y^2 + z^2y^n \subset \mathbb{P}(15n + 4, 4n + 2, 3n - 2, 7)$ $\mathbb{P}(4n + 2, 3n - 2, 7)$, Diff=($\frac{1}{2}$, 0, 0, 0)	8, 14
69	$x^i(x + y)^j f_{8-i-j}(x, y)$ $i \leq 1$ & $j \leq 3$	$t^2 + zx + g(x, y) \subset \mathbb{P}(4, 7, 1, 1)$ Diff=(0, $\frac{2}{3}$, 0, 0)	3
70	$x^8 + azx^3y^3 + y^9$	$t^2 + z^3x + x^8 + azx^3y + y^3 \subset \mathbb{P}(12, 7, 3, 8)$ Diff=(0, 0, 0, $\frac{2}{3}$)	4
71	$x^8 + ax^4y^5 + y^{10}$	$t^2 + zx + x^8 + ax^4y + y^2 \subset \mathbb{P}(4, 7, 1, 4)$ Diff=(0, $\frac{2}{3}$, 0, $\frac{4}{5}$)	6
72	$x^8 + y^{11}$	$t^2 + zx + x^8 + y \subset \mathbb{P}(4, 7, 1, 8)$ $\mathbb{P}(4, 7, 1)$, Diff=(0, $\frac{2}{3}$, 0, $\frac{10}{11}$)	12
73	$x^8 + x^2y^i$	$t^2 + zx + x^8 + x^2y \subset \mathbb{P}(4, 7, 1, 6)$	9

		$\text{Diff}=(0, \frac{2}{3}, 0, \frac{6}{7})$	
74	$x^8 + ax^5y^4 + x^2y^8$	$t^2 + zx + x^8 + ax^5y + x^2y^2 \subset \mathbb{P}(4, 7, 1, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{3}{4})$	9
75	$cz^2y^4 + azx^4y^2 + bx^8 + dx^3y^6, c \neq 0$	$t^2 + z^3x + g(z, x, y^{1/2}) \subset \mathbb{P}(12, 7, 3, 5)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	5
76	$x^8 + x^3y^7$	$t^2 + zx + x^8 + x^3y \subset \mathbb{P}(4, 7, 1, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{6}{7})$	15
77	$x^8 + zy^n$ $n = 6, 7; 8$	$t^2 + z^3x + x^8 + zy \subset \mathbb{P}(12, 7, 3, 17)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	7,17
78	$x^8 + zx^2y^5$	$t^2 + z^3x + x^8 + zx^2y \subset \mathbb{P}(12, 7, 3, 11)$ $\text{Diff}=(0, 0, 0, \frac{4}{5})$	11
79	$x'y + y^n$ $n = 9; 11$	$t + zx + x'y + y^n \subset \mathbb{P}(7n, 6n + 1, n - 1, 7)$ $\mathbb{P}(6n + 1, n - 1, 7), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,12
80	$x'y + y^{2n}$ $n = 5; 6$	$t^2 + zx + x'y + y^{2n} \subset \mathbb{P}(7n, 12n + 1, 2n - 1, 7)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12,21
81	$x'y + x^2y^n$ $n = 7; 9$	$t + zx + g(x, y) \subset \mathbb{P}(7n - 2, 6n - 1, n - 1, 5)$ $\mathbb{P}(6n - 1, n - 1, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	16,30
82	$x'y + x^2y^8$	$t^2 + zx + g(x, y) \subset \mathbb{P}(27, 47, 7, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	15
83	$x'y + zy^n$ $n = 6; 8$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(21n + 1, 6n + 1, 3n - 2, 15)$ $\mathbb{P}(6n + 1, 3n - 2, 15), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
84	$x'y + zy^7$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(74, 43, 19, 15)$ $\text{Diff} = \emptyset$	5
85	$x'y + zx^2y^5$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(50, 29, 13, 9)$ $\text{Diff} = \emptyset$	9
86	$x'y + z^2y^4$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(43, 25, 11, 9)$ $\text{Diff} = \emptyset$	9
87	$x'y + z^2y^5$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(107, 31, 14, 9)$ $\mathbb{P}(31, 14, 9), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	18
88	$x^6y^2 + y^n$ $n = 9; 13$	$t^2 + zx + x^6y^2 + y^n \subset \mathbb{P}(3n, 5n + 2, n - 2, 6)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12,18
89	$x^6y^2 + ax^3y^n + by^{2n-2}$ $n = 6, b \neq 0; 7$	$t^2 + zx + g(x, y) \subset \mathbb{P}(3n - 3, 5n - 4, n - 2, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	6,9
90	$y^2x^{2i}f_{3-i}(x^2, y^3)$ $i \leq 1$	$t^2 + zx + g(x, y) \subset \mathbb{P}(11, 19, 3, 2)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	6
91	$x^6y^2 + az^2y^4 + bzx^3y^3 + cx^2y^7, a + c \neq 0$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(19, 11, 5, 4)$ $\text{Diff} = \emptyset$	4
92	$x^6y^2 + x^2y^9$	$t^2 + zx + g(x, y) \subset \mathbb{P}(25, 43, 7, 4)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
93	$x^6y^2 + zy^n$ $n = 6; 7, 8; 9$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(9n + 1, 5n + 2, 3n - 4, 13)$ $\text{Diff} = \emptyset$	7,10,13
94	$x^6y^2 + zx^2y^5$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(40, 23, 11, 7)$ $\text{Diff} = \emptyset$	7
95	$x^6y^2 + z^2y^5$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(47, 27, 13, 8)$ $\text{Diff} = \emptyset$	8
96	$azx^2y^4 + x^5y^3 + by^9$ $b \neq 0$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(45, 13, 6, 5)$ $\mathbb{P}(13, 6, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
97	$azx^2y^5 + x^5y^3 + by^{12}$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(30, 17, 9, 5)$	5

		Diff = $\emptyset$	
98	$ax^2y^6 + x^5y^3 + by^{15}$	$t + z^3x + g(z, x, y^{1/3}) \subset \mathbb{P}(25, 7, 4, 5)$ $\mathbb{P}(7, 4, 5), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	10
99	$x^5y^3 + y^{2n}$ $n = 5; 7, 8$	$t^2 + zx + g(x, y) \subset \mathbb{P}(5n, 8n + 3, 2n - 3, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9,15
100	$x^5y^3 + y^n$ $n = 11; 17$	$t + zx + g(x, y) \subset \mathbb{P}(5n, 4n + 3, n - 3, 5)$ $\mathbb{P}(4n + 3, n - 3, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	22,30
101	$yx^i(x + y^2)^j f_{6-i-j}(x, y^2)$ $i \leq 3 \ \& \ j \leq 5$	$t + zx + g(x, y) \subset \mathbb{P}(13, 11, 2, 1)$ $\mathbb{P}(11, 2, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
102	$x^5y^3 + x^2y^n$ $n = 7; 11$	$t + zx + g(x, y) \subset \mathbb{P}(5n - 6, 4n - 3, n - 3, 3)$ $\mathbb{P}(4n - 3, n - 3, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,18
103	$x^5y^3 + x^2y^{2n}$ $n = 4, 5$	$t^2 + zx + g(x, y) \subset \mathbb{P}(5n - 3, 8n - 3, 2n - 3, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
104	$x^5y^3 + zy^{3n}$ $n = 2; 4$	$t + z^3x + x^5y + zy^n \subset \mathbb{P}(15n + 1, 4n + 1, 3n - 2, 11)$ $\mathbb{P}(4n + 1, 3n - 2, 11), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	10,22
105	$x^5y^3 + zy^{2n+1}$ $n = 3; 5$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(15n + 9, 8n + 7, 6n - 3, 11)$ $\text{Diff} = \emptyset$	5,11
106	$x^5y^3 + zy^n$ $n = 8; 10$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(15n + 3, 4n + 3, 3n - 6, 11)$ $\mathbb{P}(4n + 3, 3n - 6, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16
107	$x^5y^3 + zy^9$	$t^2 + z^3x + x^5y + zy^3 \subset \mathbb{P}(23, 13, 7, 11)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	11
108	$x^5y^3 + z^2y^4$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(33, 19, 9, 7)$ $\text{Diff} = \emptyset$	7
109	$x^5y^3 + z^2y^n$ $n = 5; 7$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(15n + 6, 4n + 3, 3n - 3, 7)$ $\mathbb{P}(4n + 3, 3n - 3, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,14
110	$x^5y^3 + z^2y^6$	$t^2 + z^3x + x^5y + z^2y^2 \subset \mathbb{P}(16, 9, 5, 7)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	7
111	$x^i(x + y)^j y^k f_{9-i-j-k}(x, y)$ $i \leq 2 \ \& \ j \leq 4 \ \& \ k \leq 4$	$t + zx + g(x, y) \subset \mathbb{P}(9, 8, 1, 1)$ $\mathbb{P}(8, 1, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
112	$x^9 + y^{10}$	$t^2 + zx + x^9 + y^2 \subset \mathbb{P}(9, 16, 2, 9)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{4}{5})$	15
113	$x^9 + x^2y^8$	$t^2 + zx + x^9 + x^2y^2 \subset \mathbb{P}(9, 16, 2, 7)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{3}{4})$	21
114	$x^9 + zy^7$	$t + z^3x + x^9 + zy \subset \mathbb{P}(27, 8, 3, 19)$ $\mathbb{P}(8, 3, 19), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{6}{7})$	14
115	$x^9 + x^2y^5$	$t + z^3x + x^9 + zx^2y \subset \mathbb{P}(27, 8, 3, 13)$ $\mathbb{P}(8, 3, 13), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{4}{5})$	26
116	$x^9 + z^2y^4$	$t^2 + z^3x + x^9 + z^2y^2 \subset \mathbb{P}(27, 16, 6, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
117	$x^8y + y^{10}$	$t^2 + zx + g(x, y) \subset \mathbb{P}(40, 71, 9, 8)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
118	$x^8y + x^2y^8$	$t^2 + zx + x^8y + x^2y^8 \subset \mathbb{P}(31, 55, 7, 6)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	18
119	$x^8y + x^3y^7$	$t + zx + x^8y + x^3y^7 \subset \mathbb{P}(53, 47, 6, 5)$ $\mathbb{P}(47, 6, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	30
120	$x^8y + zy^7$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(169, 50, 19, 17)$ $\mathbb{P}(50, 19, 17), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	24
121	$x^8y + zx^2y^5$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(115, 34, 13, 11)$	22

		$\mathbb{P}(34, 13, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	
122	$x^8y + z^2y^4$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(49, 29, 11, 10)$ $\text{Diff} = \emptyset$	10
123	$x^7y^2 + y^{10}$	$t^2 + zx + g(x, y) \subset \mathbb{P}(35, 62, 8, 7)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
124	$x^7y^2 + y^{11}$	$t + zx + g(x, y) \subset \mathbb{P}(77, 68, 9, 7)$ $\mathbb{P}(68, 9, 7), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	42
125	$x^7y^2 + x^2y^8$	$t^2 + zx + x^7y + x^2y^4 \subset \mathbb{P}(13, 23, 3, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{1}{2})$	15
126	$x^7y^2 + zy^7$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(149, 44, 17, 15)$ $\mathbb{P}(44, 17, 15), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
127	$x^7y^2 + zx^2y^5$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(95, 28, 11, 9)$ $\mathbb{P}(28, 11, 9), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	18
128	$x^7y^2 + z^2y^4$	$t^2 + z^3x + x^7y + z^2y^2 \subset \mathbb{P}(22, 13, 5, 9)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	9
129	$x^6y^3 + y^{10}$	$t^2 + zx + x^6y^3 + y^{10} \subset \mathbb{P}(30, 53, 7, 6)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
130	$x^6y^3 + y^{11}$	$t + zx + x^6y^3 + y^{11} \subset \mathbb{P}(33, 29, 4, 3)$ $\mathbb{P}(29, 4, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18
131	$x^6y^3 + x^2y^8$	$t^2 + zx + x^6y^3 + x^2y^8 \subset \mathbb{P}(21, 37, 5, 4)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
132	$x^6y^3 + zy^n$ $n = 7; 8$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(18n + 3, 5n + 3, 3n - 6, 13)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10, 26
133	$x^6y^3 + zx^2y^5$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(75, 22, 9, 7)$ $\mathbb{P}(22, 9, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
134	$x^6y^3 + z^2y^4$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(39, 23, 9, 8)$ $\text{Diff} = \emptyset$	8
135	$x^5y^4 + y^{10}$	$t^2 + zx + g(x, y) \subset \mathbb{P}(25, 44, 6, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
136	$x^5y^4 + y^{11}$	$t + z^3x + g(x, y) \subset \mathbb{P}(55, 16, 7, 5)$ $\mathbb{P}(16, 7, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
137	$x^5y^4 + y^{12}$	$t^2 + zx + x^5y^2 + y^6 \subset \mathbb{P}(15, 26, 4, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{1}{2})$	15
138	$x^5y^4 + y^{13}$	$t + zx + g(x, y) \subset \mathbb{P}(65, 56, 9, 5)$ $\mathbb{P}(56, 9, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	30
139	$x^5y^4 + x^2y^9$	$t + zx + g(x, y) \subset \mathbb{P}(37, 32, 5, 3)$ $\mathbb{P}(32, 5, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18
140	$x^5y^4 + zy^n$ $n = 7; 9$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(15n + 4, 4n + 4, 3n - 8, 11)$ $\mathbb{P}(4n + 4, 3n - 8, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16, 22
141	$x^5y^4 + zy^8$	$t^2 + z^3x + x^5y^2 + zy^4 \subset \mathbb{P}(31, 18, 8, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
142	$ax^6 + bx^5y^4 + z^2y^4$	$t^2 + z^3x + g(z, x, y^{1/2}) \subset \mathbb{P}(17, 10, 4, 7)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	7
143	$x^5y^4 + z^2y^5$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(83, 24, 11, 7)$ $\mathbb{P}(24, 11, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
144	$zx^4 + y^{2n}$ $n = 4, 5; 6, 7; 8, 9, 10$	$t^2 + z^3x + zx^4 + y^2 \subset \mathbb{P}(11, 6, 4, 11)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5, 7, 11
145	$zx^4 + y^n, n = 7;$	$t + z^3x + zx^4 + y \subset \mathbb{P}(11, 3, 2, 11)$	7, 10, 14,

	9; 13; 15; 17; 19, 21	$\mathbb{P}(3, 2, 11), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	16,18,22
146	$zx^4 + x^2y^{2n}$ $n = 4, 5, 6$	$t^2 + z^3x + zx^4 + x^2y^2 \subset \mathbb{P}(11, 6, 4, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	7
147	$zx^4 + x^2y^n$ $n = 9; 11, 13$	$t + z^3x + zx^4 + x^2y \subset \mathbb{P}(11, 3, 2, 7)$ $\mathbb{P}(3, 2, 7), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	10,14
148	$zx^4 + az^2y^n + bx^3y^n$ $n = 3 \text{ } ab \neq 0; 7, 9$	$t + z^3x + g(z, x, y^{1/n}) \subset \mathbb{P}(11, 3, 2, 5)$ $\mathbb{P}(3, 2, 5), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	4,10
149	$zx^4 + az^2y^{2n} + bx^3y^{2n}$ $n = 2 \text{ } ab \neq 0, 3 \text{ } a \neq 0; 4$	$t^2 + z^3x + g(z, x, y^{1/n}) \subset \mathbb{P}(11, 6, 4, 5)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	4,5
150	$zx^4y + y^{2n}$ $n = 4; 5, 6; 7, 8$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(11n, 6n + 1, 4n - 3, 11)$ $\text{Diff} = \emptyset$	5,7,11
151	$zx^4y + y^n$ $n = 9; 11; 13; 15; 17$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(11n, 3n + 1, 2n - 3, 11)$ $\mathbb{P}(3n + 1, 2n - 3, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,14,16, 18,22
152	$zx^4y + x^2y^{2n}$ $n = 4, 5$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(11n - 3, 6n - 1, 4n - 3, 7)$ $\text{Diff} = \emptyset$	7
153	$zx^4y + x^2y^n$ $n = 7; 9; 11$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(11n - 6, 3n - 1, 2n - 3, 7)$ $\mathbb{P}(3n - 1, 2n - 3, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,10,14
154	$zx^4y + x^3y'$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(34, 19, 11, 5)$ $\text{Diff} = \emptyset$	5
155	$zx^4y + x^3y^8$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(79, 22, 13, 5)$ $\mathbb{P}(22, 13, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
156	$zx^6 + y^n$ $n = 9; 11$	$t + z^3x + zx^6 + y \subset \mathbb{P}(17, 5, 2, 17)$ $\mathbb{P}(5, 2, 17), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	10,34
157	$zx^6 + y^{10}$	$t^2 + z^3x + zx^6 + y^2 \subset \mathbb{P}(17, 10, 4, 17)$ $\text{Diff}=(0, 0, 0, \frac{4}{5})$	10
158	$zx^6 + x^2y'$	$t + z^3x + zx^6 + x^2y \subset \mathbb{P}(17, 5, 2, 13)$ $\mathbb{P}(5, 2, 13), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{6}{7})$	14
159	$zx^6 + x^2y^8$	$t^2 + z^3x + zx^6 + x^2y^2 \subset \mathbb{P}(17, 10, 4, 13)$ $\text{Diff}=(0, 0, 0, \frac{3}{4})$	13
160	$zx^6 + x^3y^7$	$t + z^3x + zx^6 + x^3y \subset \mathbb{P}(17, 5, 2, 11)$ $\mathbb{P}(5, 2, 11), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{6}{7})$	22
161	$zx^5y + y^n$ $n = 9; 10; 11$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(7n, 4n + 1, 2n - 3, 14)$ $\text{Diff} = \emptyset$	5,7,14
162	$zx^5y + x^2y^n$ $n = 7; 8$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(7n - 3, 4n - 1, 2n - 3, 10)$ $\text{Diff} = \emptyset$	5,10
163	$zx^5y + z^2y^4 + ax^4y^5$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(29, 17, 7, 6)$ $\text{Diff} = \emptyset$	6
164	$zx^4y^2 + y^n$ $n = 9; 11; 13$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(11n, 3n + 2, 2n - 6, 11)$ $\mathbb{P}(3n + 2, 2n - 6, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14,16,22
165	$zx^4y^2 + y^{10}$	$t^2 + z^3x + g(z, x, y) \subset \mathbb{P}(55, 32, 14, 11)$ $\text{Diff} = \emptyset$	7
166	$zx^4y^2 + y^{12}$	$t^2 + z^3x + zx^4y + y^6 \subset \mathbb{P}(33, 19, 9, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
167	$zx^4y^2 + x^2y^n$ $n = 7; 9$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(11n - 12, 3n - 2, 2n - 6, 7)$ $\mathbb{P}(3n - 2, 2n - 6, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,14
168	$zx^4y^2 + x^2y^8$	$t^2 + z^3x + zx^4y + x^2y^4 \subset \mathbb{P}(19, 11, 5, 7)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	7
169	$zx^4y^2 + az^2y^5 + bx^3y^7$	$t + z^3x + g(z, x, y) \subset \mathbb{P}(59, 17, 8, 5)$	10

$$\boxed{\boxed{\mathbb{P}(17, 8, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)}\boxed{}}$$

(1) See example 3.22.

(2) The singularity is $t^2 + z^3x + \underline{a}zx^i(x+y^2)^j f_{4-i-j}(x, y^2) + bz^2y^5 + \underline{c}x^k y(x+y^2)^l f_{5-k-l}(x, y^2)$. The exceptionality condition is $b \neq 0 \parallel (b = 0 \& (i \leq 1 \parallel k \leq 2) \& (j \leq 2 \parallel l \leq 4))$.

(3) The singularity is $t^2 + z^3x + \underline{a}zx^i y^j (x+y)^k f_{5-i-j-k}(x, y) + \underline{b}x^l y^m (x+y)^n f_{7-l-m-n}(x, y) + cz^2y^3$. The common exceptionality condition for these two cases is $(c \neq 0) \parallel ((i = 0 \parallel l = 0) \& (k \leq 2 \parallel n \leq 2))$. The first case is $j \leq 1$ and the coefficient $\underline{a}$ is absent (see the comments to the tables). The second case is $m \leq 2$ and the coefficient $\underline{b}$ is absent.

4. Singularity $-t^2 + z^3y + g(z, x, y)$

1	$x^7 + y^{2n}$ $n = 4; 6; 9; 10; 12, 13$	$t^2 + zy + x + y^{2n} \subset \mathbb{P}(n, 2n-1, 2n, 1)$ $\mathbb{P}(n, 2n-1, 1), \text{Diff}=(0, \frac{2}{3}, \frac{6}{7}, 0)$	7,9,15, 21
2	$x^7 + y^n$ $n = 9; 11; 15; 17; 23; 27$	$t + zy + x + y^n \subset \mathbb{P}(n, n-1, n, 1)$ $\mathbb{P}(n-1, n, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, \frac{6}{7}, 0)$	14,16,22 28,36,42
3	$ay^{2n}z^i f_{2-i}(z, y^{2n-1}) + x^7$ $n = 2, 3, 4$	$t^2 + z^3y + g(z, x^{1/7}, y) \subset \mathbb{P}(3n-1, 2n-1, 6n-2, 1)$ $\mathbb{P}(3n-1, 2n-1, 1), \text{Diff}=(0, 0, \frac{6}{7}, 0)$	7
4	$ay^n z^i f_{2-i}(z, y^{n-1}) + x^7$ $n = 5; 7, 9$	$t + z^3y + g(z, x^{1/7}, y) \subset \mathbb{P}(3n-2, n-1, 3n-2, 1)$ $\mathbb{P}(n-1, 3n-2, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{6}{7}, 0)$	8,14
5	$x^i f_{7-i}(x, y^2)$ $i \leq 4$	$t^2 + zy + g(x, y) \subset \mathbb{P}(7, 13, 2, 1)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	3
6	$x^i f_{7-i}(x, y^3)$ $i \leq 5$	$t + zy + g(x, y) \subset \mathbb{P}(21, 20, 3, 1)$ $\mathbb{P}(20, 3, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
7	$x^7 + xy^n$ $n = 7; 11; 13; 17; 19; 23$	$t + zy + x^7 + xy^n \subset \mathbb{P}(7n, 7n-6, n, 6)$ $\mathbb{P}(7n-6, n, 6), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,16,22 30,36
8	$x^{3i+1} f_{2-i}(x^3, y^{2n})$ $n = 2 \ i = 0; (4, 5) \ i \leq 1$	$t^2 + zy + g(x, y) \subset \mathbb{P}(7n, 14n-3, 2n, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	3,9
9	$x^{3i+1} f_{2-i}(x^3, y^n)$ $n = 5, 7; 11$	$t + zy + g(x, y) \subset \mathbb{P}(7n, 7n-3, n, 3)$ $\mathbb{P}(7n-3, n, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12,18
10	$x^{2i+1} f_{3-i}(x^2, y^n)$ $n = 3 \ i \leq 1; 7 \ i \leq 2$	$t + zy + g(x, y) \subset \mathbb{P}(7n, 7n-2, n, 2)$ $\mathbb{P}(7n-2, n, 2), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6,12
11	$azy^2 x^{2j} f_{2-j}(x^2, y^5) +$ $x^{2i+1} f_{3-i}(x^2, y^5)$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(35, 11, 5, 2)$ $\mathbb{P}(11, 5, 2), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
12	$x^7 + x^2 y^{2n}$ $n = 3, 6; 8, 9$	$t^2 + zy + g(x, y) \subset \mathbb{P}(7n, 14n-5, 2n, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9,15
13	$x^7 + x^2 y^n$ $n = 7; 9; 13; 19$	$t + zy + g(x, y) \subset \mathbb{P}(7n, 7n-5, n, 5)$ $\mathbb{P}(7n-5, n, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,16,18 30
14	$azx^3 y^{2n+1} + x^7 + bx^2 y^{6n+2}$ $n = 1 \ b \neq 0; 2$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(21n+7, 14n+3, 6n+2, 5)$ $\text{Diff} = \emptyset$	4,5
15	$azx^3 y^n + x^7 + bx^2 y^{3n-1}$ $n = 4, 6$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(21n-7, 7n-4, 3n-1, 5)$ $\mathbb{P}(7n-4, 3n-1, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
16	$x^7 + x^3 y^n$ $n = 5; 9, 11; 15$	$t + zy + g(x, y) \subset \mathbb{P}(7n, 7n-4, n, 4)$ $\mathbb{P}(7n-4, n, 4), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,18,24
17	$azx^2 y^{2n-1} + x^7 +$ $+bx^3 y^{3n-2}, n = 3, 5$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(21n-14, 7n-6, 3n-2, 4)$ $\mathbb{P}(7n-6, 3n-2, 4), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8

18	$x^7 + zy^n$ $n = 6; 8, 10; 14; 16, 18$	$t + z^3y + x + zy^n \subset \mathbb{P}(3n - 1, n - 1, 3n - 1, 2)$ $\mathbb{P}(n - 1, 3n - 1, 2), \text{Diff}=(\frac{1}{2}, 0, \frac{6}{7}, 0)$	8,14,22 28
19	$x^7 + zxy^{2n+1}$ $n = 2, 3; 4, 5, 6$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(21n + 7, 14n + 1, 6n + 2, 11)$ $\text{Diff} = \emptyset$	5,11
20	$x^7 + zxy^n$ $n = 6; 8, 10; 12, 14$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(21n - 7, 7n - 6, 3n - 1, 11)$ $\mathbb{P}(7n - 6, 3n - 1, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,16,22
21	$x^7 + zx^2y^n$ $n = 4, 6; 8, 10$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(21n - 7, 7n - 5, 3n - 1, 8)$ $\mathbb{P}(7n - 5, 3n - 1, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,16
22	$x^8 + y^n$ $n = 9; 11; 15$	$t^2 + zy + x^2 + y^n \subset \mathbb{P}(n, 2n - 2, n, 2)$ $\text{Diff}=(0, \frac{2}{3}, \frac{3}{4}, 0)$	13,16,24
23	$zy^2(bx^4 + cy^5) + x^8 + dx^4y^5 +$ $ey^{10}, c + d + e \neq 0$	$t^2 + z^3y + g(z, x^{1/4}, y) \subset \mathbb{P}(5, 3, 5, 1)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	4
24	$x^{2i}f_{4-i}(x^2, y^3)$ $i \leq 2$	$t^2 + zy + g(x^{1/2}, y) \subset \mathbb{P}(6, 11, 3, 1)$ $\text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	6
25	$y^5z^if_{2-i}(z, y^4) + x^8$	$t^2 + z^3y + g(z, x^{1/4}, y) \subset \mathbb{P}(13, 8, 13, 2)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	8
26	$x^8 + ax^4y^7 + by^{14}$	$t^2 + zy + x^2 + axy^7 + by^{14} \subset \mathbb{P}(7, 13, 7, 1)$ $\text{Diff}=(0, \frac{2}{3}, \frac{3}{4}, 0)$	12
27	$azx^3y^n + x^8 + bxy^{3n-1}$ $n = 3 \text{ } b \neq 0; 4$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(12n - 4, 8n - 5, 3n - 1, 7)$ $\text{Diff} = \emptyset$	4,7
28	$x^8 + xy^n$ $n = 9; 10; 12; 13$	$t^2 + zy + x^8 + xy^n \subset \mathbb{P}(4n, 8n - 7, n, 7)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	13,15,18 21
29	$x^8 + x^2y^n$ $n = 7; 11$	$t^2 + zy + x^4 + xy^n \subset \mathbb{P}(2n, 4n - 3, n, 3)$ $\text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	10,18
30	$x^{3i+2}f_{2-i}(x^3, y^n)$ $n = 4 \text{ } i = 0; 5 \text{ } i \leq 1$	$t^2 + zy + g(x, y) \subset \mathbb{P}(4n, 8n - 3, n, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	6,9
31	$x^8 + x^3y^n$ $n = 6; 8; 9$	$t^2 + zy + g(x, y) \subset \mathbb{P}(4n, 8n - 5, n, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9,12,15
32	$bzx^2y^5 + x^8 + cx^3y^7$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(28, 17, 7, 5)$ $\text{Diff} = \emptyset$	5
33	$x^8 + zy^n$ $n = 6; 8; 10$	$t^2 + z^3y + x^2 + zy^n \subset \mathbb{P}(3n - 1, 2n - 2, 3n - 1, 4)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	10,13,16
34	$x^8 + zxy^n$ $n = 5; 6, 7; 8$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(12n - 4, 8n - 7, 3n - 1, 13)$ $\text{Diff} = \emptyset$	7,10,13
35	$x^8 + zx^2y^n$ $n = 4; 6$	$t^2 + z^3y + x^4 + zxy^n \subset \mathbb{P}(6n - 2, 4n - 3, 3n - 1, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	7,10
36	$\underline{a}z^2x^3 + bzy^2x^{2i}f_{2-i}(x^2, y^3) +$ $\underline{c}x^{2j+1}yf_{3-j}(x^2, y^3)$ $i \leq 1 j \leq 1$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(23, 7, 3, 2)$ $\mathbb{P}(7, 3, 2), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
37	$y^4z^if_{2-i}(z, y^3) + x^9$	$t^2 + z^3y + g(z, x^{1/9}, y) \subset \mathbb{P}(5, 3, 10, 1)$ $\mathbb{P}(5, 3, 1), \text{Diff}=(0, 0, \frac{8}{9}, 0)$	9
38	$x^9 + ax^3y^4 + by^{11}$	$t + z^3y + g(z, x^{1/3}, y) \subset \mathbb{P}(33, 10, 11, 3)$ $\mathbb{P}(10, 11, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	18
39	$x^9 + xy^9$	$t + zy + g(x, y) \subset \mathbb{P}(81, 73, 9, 8)$ $\mathbb{P}(73, 9, 8), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	30
40	$x^9 + ax^5y^5 + bxy^{10}$	$t + zy + g(x, y) \subset \mathbb{P}(45, 41, 5, 4)$ $\mathbb{P}(41, 5, 4), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	24
41	$x^9 + x^2y^8$	$t^2 + zy + g(x, y) \subset \mathbb{P}(36, 65, 8, 7)$	12

		Diff=(0, $\frac{2}{3}$, 0, 0)	
42	$x^9 + x^2y^9$	$t + zy + g(x, y) \subset \mathbb{P}(81, 74, 9, 7)$ $\mathbb{P}(74, 9, 7)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	42
43	$bx^2y^5 + x^9 + cx^3y^7$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(63, 19, 7, 6)$ $\mathbb{P}(19, 7, 6)$, Diff=($\frac{1}{2}$, 0, 0, 0)	12
44	$x^9 + x^4y^6$	$t^2 + zy + g(x, y) \subset \mathbb{P}(27, 49, 6, 5)$ Diff=(0, $\frac{2}{3}$, 0, 0)	15
45	$x^9 + zy^8$	$t + z^3y + x + zy^8 \subset \mathbb{P}(23, 7, 23, 2)$ $\mathbb{P}(7, 23, 2)$, Diff=($\frac{1}{2}$, 0, $\frac{8}{9}$, 0)	36
46	$x^9 + zxy^6$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(153, 46, 17, 15)$ $\mathbb{P}(46, 17, 15)$, Diff=($\frac{1}{2}$, 0, 0, 0)	10
47	$ay^4z^i f_{2-i}(z, y^3) + x^7y^2$	$t^2 + z^3y + g(z, x^{1/7}, y) \subset \mathbb{P}(5, 3, 8, 1)$ Diff=(0, 0, $\frac{6}{7}$, 0)	7
48	$ay^5z^i f_{2-i}(z, y^4) + x^7y^2$	$t + z^3y + g(z, x^{1/7}, y) \subset \mathbb{P}(13, 4, 11, 1)$ $\mathbb{P}(4, 11, 1)$, Diff=($\frac{1}{2}$, 0, $\frac{6}{7}$, 0)	14
49	$x^7y^2 + y^n$ $n = 11; 15$	$t + zy + xy^2 + y^n \subset \mathbb{P}(n, n-1, n-2, 1)$ $\mathbb{P}(n-1, n-2, 1)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, $\frac{6}{7}$, 0)	28, 42
50	$x^7y^2 + y^{2n}$ $n = 6; 7$	$t^2 + zy + xy^2 + y^{2n} \subset \mathbb{P}(n, 2n-1, 2n-2, 1)$ Diff=(0, $\frac{2}{3}$, $\frac{6}{7}$, 0)	15, 21
51	$x^7y^2 + xy^n$ $n = 9; 13$	$t + zy + x^7y^2 + xy^n \subset \mathbb{P}(7n-2, 7n-8, n-2, 6)$ $\mathbb{P}(7n-8, n-2, 6)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	22, 36
52	$x^7y^2 + ax^4y^6 + bxy^{10}$	$t^2 + zy + g(x, y) \subset \mathbb{P}(17, 31, 4, 3)$ Diff=(0, $\frac{2}{3}$, 0, 0)	9
53	$y^2x^{2i+1}f_{3-i}(x^2, y^3)$ $i \leq 2$	$t + zy + g(x, y) \subset \mathbb{P}(25, 23, 3, 2)$ $\mathbb{P}(23, 3, 2)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	12
54	$x^7y^2 + ax^4y^7 + bxy^{12}$	$t + zy + g(x, y) \subset \mathbb{P}(41, 38, 5, 3)$ $\mathbb{P}(38, 5, 3)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	18
55	$x^7y^2 + x^2y^{2n}$ $n = 4; 5$	$t^2 + zy + g(x, y) \subset \mathbb{P}(7n-2, 14n-9, 2n-2, 5)$ Diff=(0, $\frac{2}{3}$, 0, 0)	9, 15
56	$azx^3y^4 + x^7y^2 + bx^2y^9$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(59, 18, 7, 5)$ $\mathbb{P}(18, 7, 5)$, Diff=($\frac{1}{2}$, 0, 0, 0)	10
57	$x^7y^2 + x^2y^{11}$	$t + zy + g(x, y) \subset \mathbb{P}(73, 68, 9, 5)$ $\mathbb{P}(68, 9, 5)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	30
58	$bx^6 + cx^2y^5 + dx^7y^2 + ex^3y^7, b + d \neq 0$ $ c + e \neq 0$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(43, 13, 5, 4)$ $\mathbb{P}(13, 5, 4)$, Diff=($\frac{1}{2}$, 0, 0, 0)	8
59	$x^7y^2 + x^3y^9$	$t + zy + g(x, y) \subset \mathbb{P}(57, 53, 7, 4)$ $\mathbb{P}(53, 7, 4)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	24
60	$x^7y^2 + zy^n$ $n = 8; 10$	$t + z^3y + xy^2 + zy^n \subset \mathbb{P}(3n-1, n-1, 3n-5, 2)$ $\mathbb{P}(n-1, 3n-5, 2)$, Diff=($\frac{1}{2}$, 0, $\frac{6}{7}$, 0)	22, 28
61	$x^7y^2 + zxy^n$ $n = 6; 8$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(21n-13, 7n-8, 3n-5, 11)$ $\mathbb{P}(7n-8, 3n-5, 11)$, Diff=($\frac{1}{2}$, 0, 0, 0)	16, 22
62	$x^7y^2 + zxy^7$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(67, 41, 16, 11)$ Diff = $\emptyset$	11
63	$x^7y^2 + zx^2y^6$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(107, 33, 13, 8)$ $\mathbb{P}(33, 13, 8)$, Diff=($\frac{1}{2}$, 0, 0, 0)	16
64	$zx^5 + y^{2n}$ $n = 4; 7; 9, 10$	$t^2 + z^3y + zx + y^{2n} \subset \mathbb{P}(3n, 2n-1, 4n+1, 3)$ Diff=(0, 0, $\frac{4}{5}$, 0)	5, 11, 15

65	$zx^5 + y^n$ $n = 9; 11; 15; 21$	$t + z^3y + zx + y^n \subset \mathbb{P}(3n, n-1, 2n+1, 3)$ $\mathbb{P}(n-1, 2n+1, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	12,15,20 30
66	$ay^{2n}z^if_{2-i}(z, y^{2n-1}) + zx^5$ $+dx^5y^{2n-1}, n=2 \ a \neq 0, 3$	$t^2 + z^3y + g(z, x^{1/5}, y) \subset \mathbb{P}(3n-1, 2n-1, 4n-1, 1)$ $\text{Diff}=(0, 0, \frac{4}{5}, 0)$	5
67	$ay^n z^i f_{2-i}(z, y^{n-1}) + zx^5 +$ $+cx^5y^{n-1}, n=5 \ a \neq 0; 7$	$t + z^3y + g(z, x^{1/5}, y) \subset \mathbb{P}(3n-2, n-1, 2n-1, 1)$ $\mathbb{P}(n-1, 2n-1, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	6,10
68	$zx^5 + bx^2y^5 +$ $+ex^6y^2 + cx^3y^7 + dy^{12}$ $ b + c + d \neq 0$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(18, 11, 5, 3)$ $\text{Diff} = \emptyset$	3
69	$zx^5 + bx^2y^7 +$ $+ex^6y^3 + cx^3y^{10} + dy^{17}$ $ b + c + d \neq 0$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(51, 16, 7, 3)$ $\mathbb{P}(16, 7, 3) \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
70	$zx^5 + xy^{2n+1}$ $n = 3; 4, 5, 6; 7, 8$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(15n+8, 10n+1, 4n+3, 13)$ $\text{Diff} = \emptyset$	5,9,13
71	$zx^5 + xy^n$ $n = 8; 10; 12; 14, 16; 18$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(15n+1, 5n-4, 2n+1, 13)$ $\mathbb{P}(5n-4, 2n+1, 13), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	11,14,16 22,26
72	$zx^5 + x^2y^{2n}$ $n = 3, 4, 5; 6, 7$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(15n+1, 10n-3, 4n+1, 11)$ $\text{Diff} = \emptyset$	7,11
73	$zx^5 + x^2y^n$ $n = 7; 9; 11, 13; 15$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(15n+2, 5n-3, 2n+1, 11)$ $\mathbb{P}(5n-3, 2n+1, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	12,14,18 22
74	$zx^5 + x^3y^{2n+1}$ $n = 2; 4, 5$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(15n+9, 10n+3, 4n+3, 9)$ $\text{Diff} = \emptyset$	8,9
75	$zx^5 + x^3y^n$ $n = 6; 8; 12$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(15n+3, 5n-2, 2n+1, 9)$ $\mathbb{P}(5n-2, 2n+1, 9), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,14,18
76	$zx^5 + x^4y^{2n}$ $n = 3, 4$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(15n+2, 10n-1, 4n+1, 7)$ $\text{Diff} = \emptyset$	7
77	$zx^5 + x^4y^n$ $n = 5, 7; 9$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(15n+4, 5n-1, 2n+1, 7)$ $\mathbb{P}(5n-1, 2n+1, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,14
78	$zx^6 + y^n$ $n = 9, 11$	$t^2 + z^3y + zx^2 + y^n \subset \mathbb{P}(3n, 2n-2, 2n+1, 6)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	9
79	$ay^4z^if_{2-i}(z, y^3) + zx^6 +$ $+cx^6y^3$	$t^2 + z^3y + g(z, x^{1/6}, y) \subset \mathbb{P}(5, 3, 7, 1)$ $\text{Diff}=(0, 0, \frac{5}{6}, 0)$	6
80	$zx^6 + y^{12}$	$t^2 + z^3y + g(z, x^{1/6}, y) \subset \mathbb{P}(18, 11, 25, 3)$ $\text{Diff}=(0, 0, \frac{5}{6}, 0)$	18
81	$zx^6 + xy^n$ $n = 8, 9, 10; 11$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(18n+1, 6n-5, 2n+1, 16)$ $\mathbb{P}(6n-5, 2n+1, 16), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14,32
82	$zx^6 + x^2y^{2n+1}$ $n = 3; 4$	$t^2 + z^3y + g(z, x^{1/2}, y) \subset \mathbb{P}(9n+5, 6n+1, 4n+3, 7)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	5,14
83	$zx^6 + x^2y^8$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(73, 44, 17, 14)$ $\text{Diff} = \emptyset$	7
84	$zx^6 + x^3y^n$ $n = 6; 8$	$t + z^3y + zx^2 + xy^n \subset \mathbb{P}(6n+1, 2n-1, 2n+1, 4)$ $\mathbb{P}(2n-1, 2n+1, 4), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	10,24
85	$zx^6 + x^4y^5$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(47, 28, 11, 10)$ $\text{Diff} = \emptyset$	5
86	$zx^6 + x^4y^6$	$t^2 + z^3y + zx^3 + x^2y^6 \subset \mathbb{P}(28, 17, 13, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	10
87	$zx^6 + x^5y^5$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(95, 29, 11, 8)$ $\mathbb{P}(29, 11, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16

88	$z^2x^3 + y^{2n}$ $n = 4; 6; 7$	$t^2 + z^3y + z^2x + y^{2n} \subset \mathbb{P}(3n, 2n-1, 2n+2, 3)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	5,7,9
89	$z^2x^3 + y^n$ $n = 9; 11; 15$	$t + z^3y + z^2x + y^n \subset \mathbb{P}(3n, n-1, n+2, 3)$ $\mathbb{P}(n-1, n+2, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	10,12,18
90	$z^2x^3 + zy^3(bx^3 + cy^4) + dx^6y^2$ $+ex^3y^6 + ly^{10}, c + e + l \neq 0$	$t^2 + z^3y + g(z, x^{1/3}, y) \subset \mathbb{P}(5, 3, 4, 1)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	3
91	$z^2x^3 + zy^4(bx^3 + cy^5) +$ $dx^6y^3 + ex^3y^8 + ly^{13}$	$t + z^3y + g(z, x^{1/3}, y) \subset \mathbb{P}(13, 4, 5, 1)$ $\mathbb{P}(4, 5, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	6
92	$z^2x^3 + xy^n$ $n = 7; 9; 11; 13$	$t + z^3y + z^2x^3 + xy^n \subset \mathbb{P}(9n+2, 3n-2, n+2, 8)$ $\mathbb{P}(3n-2, n+2, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,10,14 16
93	$z^2x^3 + azx^2y^n + bx^5y^{n-1} +$ $+cxy^{2n}, n = 4 \mid a \mid + \mid c \mid \neq 0; 6$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(9n+1, 3n-1, n+1, 4)$ $\mathbb{P}(3n-1, n+1, 4), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,8
94	$z^2x^3 + x^2y^{2n}$ $n = 3, 4; 5$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(9n+2, 6n-1, 2n+2, 7)$ $\text{Diff} = \emptyset$	5,7
95	$z^2x^3 + x^2y^n$ $n = 7; 9; 11$	$t + z^3y + z^2x^3 + x^2y^n \subset \mathbb{P}(9n+4, 3n-1, n+2, 7)$ $\mathbb{P}(3n-1, n+2, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,12,14
96	$z^2x^3 + azy^n + bx^3y^{n-1}$ $n = 6; 10$	$t + z^3y + z^2x + zy^n + xy^{n-1} \subset \mathbb{P}(3n-1, n-1, n+1, 2)$ $\mathbb{P}(n-1, n+1, 2), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	8,12
97	$z^2x^3 + azxy^{2n+1} + bx^4y^{2n}$ $n = 2 \mid a \neq 0; 3$	$t^2 + z^3y + g(z, x, y) \subset \mathbb{P}(9n+4, 6n+1, 2n+2, 5)$ $\text{Diff} = \emptyset$	3,5
98	$z^2x^3 + azxy^{n+1} + bx^4y^n$ $n = 5; 7$	$t + z^3y + g(z, x, y) \subset \mathbb{P}(9n+8, 3n+1, n+2, 5)$ $\mathbb{P}(3n+1, n+2, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,10

5. Singularity $-t^2 + z^5 + g(z, x, y)$

1	$x^i y^j f_{6-i-j}(x, y)$ $i \leq 2 \& j \leq 2$	$t^2 + z + g(z, x, y) \subset \mathbb{P}(3, 6, 1, 1)$ $\mathbb{P}(3, 1, 1), \text{Diff}=(0, \frac{4}{5}, 0, 0)$	5
2	$x^6 + y^7$	$t^2 + z + x^2 + y \subset \mathbb{P}(1, 2, 1, 2)$ $\mathbb{P}(1, 1, 2), \text{Diff}=(0, \frac{4}{5}, \frac{2}{3}, \frac{6}{7})$	15
3	$x^6 + xy^6$	$t^2 + z + x^6 + xy \subset \mathbb{P}(3, 6, 1, 5)$ $\mathbb{P}(3, 1, 5), \text{Diff}=(0, \frac{4}{5}, 0, \frac{5}{6})$	25
4	$x^6 + zy^5$	$t^2 + z^5 + x^2 + zy \subset \mathbb{P}(5, 2, 5, 8)$ $\text{Diff}=(0, 0, \frac{2}{3}, \frac{4}{5})$	6
5	$x^6 + zxy^4$	$t^2 + z^5 + x^6 + zxy \subset \mathbb{P}(15, 6, 5, 19)$ $\text{Diff}=(0, 0, 0, \frac{3}{4})$	8
6	$x^6 + zx^2y^3$	$t^2 + z^5 + x^6 + zx^2y \subset \mathbb{P}(15, 6, 5, 14)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	7
7	$ax^6 + bzx^3y^2 + z^2y^4$	$t^2 + z^5 + ax^2 + bzxxy + z^2y^2 \subset \mathbb{P}(5, 2, 5, 3)$ $\text{Diff}=(0, 0, \frac{2}{3}, \frac{1}{2})$	9
8	$x^6 + z^2xy^3$	$t^2 + z^5 + x^6 + z^2xy \subset \mathbb{P}(15, 6, 5, 13)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	13
9	$az^2xy^3 + bz^3y^2 + cx^5y +$ $dy^7, a + d \neq 0$	$t + z^5 + g(z, x, y) \subset \mathbb{P}(35, 7, 6, 5)$ $\mathbb{P}(7, 6, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
10	$x^5y + xy^6$	$t + z + x^5y + xy^6 \subset \mathbb{P}(29, 29, 5, 4)$ $\mathbb{P}(29, 5, 4), \text{Diff}=(\frac{1}{2}, \frac{4}{5}, 0, 0)$	30
11	$x^5y + x^2y^5$	$t + z + x^5y + x^2y^5 \subset \mathbb{P}(23, 23, 4, 3)$ $\mathbb{P}(23, 4, 3), \text{Diff}=(\frac{1}{2}, \frac{4}{5}, 0, 0)$	30
12	$x^5y + zy^5$	$t + z^5 + xy + zy^5 \subset \mathbb{P}(25, 5, 21, 4)$	30

		$\mathbb{P}(5, 21, 4), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	
13	$x^5y + az^6y^3 + bz^3y^3$	$t + z^5 + xy + az^6y^3 + bz^3y^3 \subset \mathbb{P}(15, 3, 13, 2)$ $\mathbb{P}(3, 13, 2), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	20
14	$x^5y + zxy^4$	$t + z^5 + x^5y + zxy^4 \subset \mathbb{P}(95, 19, 16, 15)$ $\mathbb{P}(19, 16, 15), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
15	$x^5y + zx^2y^3$	$t + z^5 + x^5y + zx^2y^3 \subset \mathbb{P}(65, 13, 11, 10)$ $\mathbb{P}(13, 11, 10), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
16	$x^5y + z^2y^4$	$t^2 + z^5 + xy + z^2y^4 \subset \mathbb{P}(10, 4, 17, 3)$ $\text{Diff}=(0, 0, \frac{4}{5}, 0)$	10
17	$x^4y^2 + y^n$ $n = 7; 9$	$t^2 + z + x^2y^2 + y^n \subset \mathbb{P}(n, 2n, n - 2, 2)$ $\mathbb{P}(n, n - 2, 2), \text{Diff}=(0, \frac{4}{5}, \frac{1}{2}, 0)$	15, 20
18	$x^4y^2 + ax^2y^5 + by^8$	$t^2 + z + x^2y^2 + axy^5 + by^8 \subset \mathbb{P}(4, 8, 3, 1)$ $\mathbb{P}(4, 3, 1), \text{Diff}=(0, \frac{4}{5}, \frac{1}{2}, 0)$	10
19	$x^4y^2 + xy^n$ $n = 6; 7$	$t^2 + z + x^4y^2 + xy^n \subset \mathbb{P}(2n - 1, 4n - 2, n - 2, 3)$ $\mathbb{P}(2n - 1, n - 2, 3), \text{Diff}=(0, \frac{4}{5}, 0, 0)$	10, 15
20	$x^4y^2 + zy^n$ $n = 5; 7$	$t^2 + z^5 + x^2y^2 + zy^n \subset \mathbb{P}(5n, 2n, 5n - 8, 8)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	11, 16
21	$x^4y^2 + zy^6$	$t^2 + z^5 + x^2y^2 + zy^6 \subset \mathbb{P}(15, 6, 11, 4)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	8
22	$x^4y^2 + zxy^n$ $n = 4; 5$	$t^2 + z^5 + g(z, x, y) \subset \mathbb{P}(10n - 5, 4n - 2, 5n - 8, 11)$ $\text{Diff} = \emptyset$	6, 11
23	$x^4y^2 + az^2y^4 + bz^2y^3$	$t^2 + z^5 + g(z, x^{1/2}, y) \subset \mathbb{P}(10, 4, 7, 3)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	6
24	$x^4y^2 + z^2y^5$	$t^2 + z^5 + x^2y^2 + z^2y^5 \subset \mathbb{P}(25, 10, 19, 6)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	12
25	$x^4y^2 + z^2xy^3$	$t^2 + z^5 + g(z, x, y) \subset \mathbb{P}(25, 10, 9, 7)$ $\text{Diff} = \emptyset$	7
26	$zx^5 + y^7$	$t + z^5 + zx + y \subset \mathbb{P}(5, 1, 4, 5)$ $\mathbb{P}(1, 4, 5), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, \frac{6}{7})$	50
27	$zx^5 + z^2y^4$	$t^2 + z^5 + zx + z^2y^2 \subset \mathbb{P}(5, 2, 8, 3)$ $\text{Diff}=(0, 0, \frac{4}{5}, \frac{1}{2})$	15
28	$zx^5 + z^2xy^3$	$t + z^5 + zx^5 + z^2xy \subset \mathbb{P}(25, 5, 4, 11)$ $\mathbb{P}(5, 4, 11), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	22
29	$zx^4y + y^7$	$t^2 + z^5 + zx^2y + y^7 \subset \mathbb{P}(35, 14, 23, 10)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	20
30	$zx^4y + xy^6$	$t + z^5 + zx^4y + xy^6 \subset \mathbb{P}(115, 23, 19, 16)$ $\mathbb{P}(23, 19, 16), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	32
31	$zx^4y + z^2y^4$	$t^2 + z^5 + zxy + z^2y^4 \subset \mathbb{P}(10, 4, 13, 3)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	12
32	$zx^4y + z^2xy^3$	$t + z^5 + zx^4y + z^2xy^3 \subset \mathbb{P}(55, 11, 9, 8)$ $\mathbb{P}(11, 9, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16
33	$zx^3y^2 + y^8$	$t^2 + z^5 + zxy + y^4 \subset \mathbb{P}(10, 4, 11, 5)$ $\text{Diff}=(0, 0, \frac{2}{3}, \frac{1}{2})$	15
34	$zx^3y^2 + y^9$	$t + z^5 + zxy^2 + y^9 \subset \mathbb{P}(45, 9, 26, 5)$ $\mathbb{P}(9, 26, 5), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	30
35	$zx^3y^2 + xy^6$	$t^2 + z^5 + zx^3y + xy^3 \subset \mathbb{P}(20, 8, 7, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
36	$zx^3y^2 + xy^7$	$t + z^5 + zx^3y^2 + xy^7 \subset \mathbb{P}(95, 19, 18, 11)$	22

		$\mathbb{P}(19, 18, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	
37	$zx^3y^2 + x^2y^5$	$t + z^5 + zx^3y^2 + x^2y^5 \subset \mathbb{P}(55, 11, 10, 7)$ $\mathbb{P}(11, 10, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
38	$zx^3y^2 + z^2y^5$	$t + z^5 + zx^3y^2 + z^2y^5 \subset \mathbb{P}(25, 5, 14, 3)$ $\mathbb{P}(5, 14, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	18

6. Singularity $-t^2 + z^4x + g(z, x, y)$

1	$x^i y^j f_{6-i-j}(x, y)$ $i \leq 1, j \leq 2$	$t^2 + zx + g(z, x, y) \subset \mathbb{P}(3, 5, 1, 1)$ $\text{Diff}=(0, \frac{3}{4}, 0, 0)$	4
2	$x^6 + y^7$	$t^2 + zx + x^6 + y \subset \mathbb{P}(3, 5, 1, 6)$ $\mathbb{P}(3, 5, 1), \text{Diff}=(0, \frac{3}{4}, 0, \frac{6}{7})$	8
3	$x^6 + x^2y^5$	$t^2 + zx + x^6 + x^2y \subset \mathbb{P}(3, 5, 1, 4)$ $\text{Diff}=(0, \frac{3}{4}, 0, \frac{4}{5})$	8
4	$x^6 + zy^n$ $n = 5; 6$	$t^2 + z^4x + x^6 + zy \subset \mathbb{P}(12, 5, 4, 19)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5, 19
5	$ax^6 + bzx^3y^2 + z^2y^4$	$t^2 + z^4x + ax^6 + bzx^3y + z^2y^2 \subset \mathbb{P}(12, 5, 4, 7)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	7
6	$x^5y + y^7$	$t^2 + z^2x + x^5y + y^7 \subset \mathbb{P}(35, 29, 12, 10)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	8
7	$x^5y + y^8$	$t^2 + zx + x^5y + y^8 \subset \mathbb{P}(20, 33, 7, 5)$ $\text{Diff}=(0, \frac{3}{4}, 0, 0)$	20
8	$x^5y + x^2y^5$	$t^2 + z^2x + x^5y + x^2y^5 \subset \mathbb{P}(23, 19, 8, 6)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	12
9	$x^5y + zy^n$ $n = 5; 6$	$t + z^4x + x^5y + zy^n \subset \mathbb{P}(20n + 1, 4n + 1, 4n - 3, 16)$ $\mathbb{P}(4n + 1, 4n - 3, 16), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6, 14
10	$x^5y + z^2y^4$	$t^2 + z^4x + x^5y + z^2y^4 \subset \mathbb{P}(41, 17, 14, 12)$ $\text{Diff} = \emptyset$	6
11	$x^5y + z^3y^3$	$t + z^4x + x^5y + z^3y^3 \subset \mathbb{P}(63, 13, 11, 8)$ $\mathbb{P}(13, 11, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16
12	$x^4y^2 + y^n$ $n = 7; 9$	$t^2 + zx + x^4y^2 + y^n \subset \mathbb{P}(2n, 3n + 2, n - 2, 4)$ $\text{Diff}=(0, \frac{3}{4}, 0, 0)$	12, 16
13	$x^4y^2 + y^8$	$t^2 + zx + x^4y^2 + y^8 \subset \mathbb{P}(8, 13, 3, 2)$ $\text{Diff}=(0, \frac{3}{4}, 0, 0)$	8
14	$x^4y^2 + zy^n$ $n = 5, 6; 7$	$t^2 + z^4x + g(z, x, y) \subset \mathbb{P}(8n + 1, 3n + 2, 4n - 6, 13)$ $\text{Diff} = \emptyset$	9, 13
15	$x^4y^2 + ax^2y^3 + z^2y^4$	$t^2 + z^4x + g(z, x, y) \subset \mathbb{P}(17, 7, 6, 5)$ $\text{Diff} = \emptyset$	5
16	$x^4y^2 + z^2y^5$	$t^2 + z^2x + x^4y^2 + zy^5 \subset \mathbb{P}(21, 17, 8, 5)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	10
17	$x^4y^2 + z^3y^3$	$t^2 + z^4x + g(z, x, y) \subset \mathbb{P}(27, 11, 10, 7)$ $\text{Diff} = \emptyset$	7
18	$zx^5 + y^7$	$t + z^4x + zx^5 + y \subset \mathbb{P}(19, 4, 3, 19)$ $\mathbb{P}(4, 3, 19), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{6}{7})$	14
19	$zx^5 + x^2y^5$	$t + z^4x + zx^5 + x^2y \subset \mathbb{P}(19, 4, 3, 13)$ $\mathbb{P}(4, 3, 13), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{4}{5})$	26
20	$zx^5 + z^2y^4$	$t^2 + z^4x + zx^5 + z^2y^2 \subset \mathbb{P}(19, 8, 6, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11

21	$zx^4y + y^7$	$t + z^4x + zx^4y + y^7 \subset \mathbb{P}(105, 22, 17, 15)$ $\mathbb{P}(22, 17, 15), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
22	$zx^4y + x^2y^5$	$t + z^4x + zx^4y + x^2y^5 \subset \mathbb{P}(67, 14, 11, 9)$ $\mathbb{P}(14, 11, 9), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	18
23	$zx^4y + z^2y^4$	$t^2 + z^4x + g(z, x, y) \subset \mathbb{P}(31, 13, 10, 9)$ $\text{Diff} = \emptyset$	9
24	$zx^3y^2 + y^n$ $n = 7; 9$	$t + z^4x + zx^3y^2 + y^n \subset \mathbb{P}(11n, 2n + 2, 3n - 8, 11)$ $\mathbb{P}(2n + 2, 3n - 8, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16, 22
25	$zx^3y^2 + y^8$	$t^2 + z^4x + zx^3y + y^4 \subset \mathbb{P}(22, 9, 8, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
26	$zx^3y^2 + x^2y^5$	$t + z^4x + zx^3y^2 + x^2y^5 \subset \mathbb{P}(39, 8, 7, 5)$ $\mathbb{P}(8, 7, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
27	$zx^3y^2 + z^2y^5$	$t + z^4x + zx^3y^2 + z^2y^5 \subset \mathbb{P}(59, 12, 11, 7)$ $\mathbb{P}(12, 11, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14

7. Singularity $-t^2 + z^4y + g(z, x, y)$

1	$x^6 + azx^3y^2 + bz^2y^4 + cy^7, b + c \neq 0$	$t^2 + z^4y + g(z, x^{1/3}, y) \subset \mathbb{P}(7, 3, 7, 2)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	6
2	$x^6 + ax^3y^4 + by^8$	$t^2 + zy + x^2 + axy^4 + by^8 \subset \mathbb{P}(4, 7, 4, 1)$ $\text{Diff}=(0, \frac{3}{4}, \frac{2}{3}, 0)$	12
3	$x^6 + xy^n$ $n = 6; 7$	$t^2 + zy + x^6 + xy^n \subset \mathbb{P}(3n, 6n - 5, n, 5)$ $\text{Diff}=(0, \frac{3}{4}, 0, 0)$	8, 20
4	$x^6 + azxy^3 + bx^2y^5$	$t^2 + z^2y + g(z, x, y) \subset \mathbb{P}(15, 13, 5, 4)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	8
5	$x^6 + zy^n$ $n = 5; 6$	$t^2 + z^4y + x^2 + zy^n \subset \mathbb{P}(4n - 1, 2n - 2, 4n - 1, 6)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	6, 9
6	$x^6 + zxy^n$ $n = 4; 5$	$t^2 + z^4y + g(z, x, y) \subset \mathbb{P}(12n - 3, 6n - 5, 4n - 1, 14)$ $\text{Diff} = \emptyset$	5, 14
7	$x^6 + zx^2y^3$	$t^2 + z^4y + g(z, x, y) \subset \mathbb{P}(33, 14, 11, 10)$ $\text{Diff} = \emptyset$	5
8	$az^3x^2 + bz^2xy^3 + czx^3y^2 + dzy^6 + ex^5y + kx^2y^5 + lz^4y$ can. & $ e + c \neq 0$	$t + z^4y + g(z, x, y) \subset \mathbb{P}(23, 5, 4, 3)$ $\mathbb{P}(5, 4, 3), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
9	$zx^5 + az^2y^4 + by^7$	$t^2 + z^4y + zx + az^2y^4 + by^7 \subset \mathbb{P}(7, 3, 11, 2)$ $\text{Diff}=(0, 0, \frac{4}{5}, 0)$	10
10	$zx^5 + xy^6$	$t + z^4y + zx^5 + xy^6 \subset \mathbb{P}(121, 26, 19, 17)$ $\mathbb{P}(26, 19, 17), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	24
11	$zx^5 + x^2y^5$	$t + z^4y + zx^5 + x^2y^5 \subset \mathbb{P}(51, 11, 8, 7)$ $\mathbb{P}(11, 8, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16
12	$zx^5 + x^3y^4$	$t + z^4y + zx^5 + x^3y^4 \subset \mathbb{P}(83, 18, 13, 11)$ $\mathbb{P}(18, 13, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	22

8. Singularity $-t^2 + z^3x^2 + g(z, x, y)$

1	$x^i y^j f_{6-i-j}(x, y)$ $i \leq 1, j \leq 3$	$t^2 + zx^2 + g(z, x, y) \subset \mathbb{P}(3, 4, 1, 1)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	3
2	$x^6 + y^7$	$t^2 + zx^2 + x^6 + y \subset \mathbb{P}(3, 4, 1, 6)$	7

		$\mathbb{P}(3, 4, 1), \text{Diff}=(0, \frac{2}{3}, 0, \frac{6}{7})$	
3	$ax^6 + bx^3y^4 + y^8$	$t^2 + zx^2 + ax^6 + bx^3y + y^2 \subset \mathbb{P}(3, 4, 1, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{3}{4})$	9
4	$z^2y^4 + bx^3y^2 + cx^6 + dxy^6$ $ b + c \neq 0$	$t^2 + z^3x^2 + g(z, x, y^{1/2}) \subset \mathbb{P}(9, 4, 3, 5)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	5
5	$x^6 + xy^7$	$t^2 + zx^2 + x^6 + xy \subset \mathbb{P}(3, 4, 1, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{6}{7})$	15
6	$x^6 + zy^5$	$t^2 + z^3x^2 + x^6 + zy \subset \mathbb{P}(9, 4, 3, 14)$ $\text{Diff}=(0, 0, 0, \frac{4}{5})$	5
7	$x^6 + zy^6$	$t^2 + z^3x^2 + x^6 + zy^2 \subset \mathbb{P}(9, 4, 3, 7)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	7
8	$x^6 + zxy^n$ $n = 4; 5$	$t^2 + z^3x^2 + x^6 + zxy \subset \mathbb{P}(9, 4, 3, 11)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	4,11
9	$x^5y + y^n$ $n = 7; 9$	$t + zx^2 + x^5y + y^n \subset \mathbb{P}(5n, 3n + 2, n - 1, 5)$ $\mathbb{P}(3n + 2, n - 1, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	15,30
10	$x^5y + y^8$	$t^2 + zx^2 + x^5y + y^8 \subset \mathbb{P}(20, 26, 7, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	13
11	$x^5y + xy^6$	$t + zx^2 + x^5y + xy^6 \subset \mathbb{P}(29, 19, 5, 4)$ $\mathbb{P}(19, 5, 4), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
12	$az^3y^3 + bx^5y + cx^3y^4 + dxy^7$ $ b + c \neq 0, a + d \neq 0$	$t + zx^2 + g(z, x, y) \subset \mathbb{P}(17, 11, 3, 2)$ $\mathbb{P}(11, 3, 2), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12
13	$x^5y + zy^n$ $n = 5; 7$	$t + z^3x^2 + x^5y + zy^n \subset \mathbb{P}(15n + 2, 3n + 2, 3n - 2, 12)$ $\mathbb{P}(3n + 2, 3n - 2, 12), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	11,24
14	$x^5y + zxy^4$	$t + z^3x^2 + x^5y + zxy^4 \subset \mathbb{P}(59, 13, 10, 9)$ $\mathbb{P}(13, 10, 9), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
15	$x^5y + zxy^5$	$t^2 + z^3x^2 + x^5y + zxy^5 \subset \mathbb{P}(37, 16, 13, 9)$ $\text{Diff} = \emptyset$	9
16	$x^5y + z^2y^4$	$t^2 + z^3x^2 + x^5y + z^2y^4 \subset \mathbb{P}(32, 14, 11, 9)$ $\text{Diff} = \emptyset$	7
17	$x^5y + z^2y^5$	$t + z^3x^2 + x^5y + z^2y^5 \subset \mathbb{P}(79, 17, 14, 9)$ $\mathbb{P}(17, 14, 9), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	18
18	$az^4y + bz^2y^4 + czx^2y^3 +$ $+x^4y^2 + dy^7$	$t^2 + z^3x^2 + g(z, x^{1/2}, y) \subset \mathbb{P}(7, 3, 5, 2)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	4
19	$az^3y^3 + x^4y^2 + bx^2y^5 +$ $+cy^8, a + c \neq 0$	$t^2 + zx + g(z^{1/3}, x^{1/2}, y) \subset \mathbb{P}(4, 5, 3, 1)$ $\text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	6
20	$x^4y^2 + y^9$	$t^2 + zx + x^2y^2 + y^9 \subset \mathbb{P}(9, 11, 7, 2)$ $\text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	12
21	$x^4y^2 + xy^n$ $n = 6; 7$	$t^2 + zx^2 + g(z, x, y) \subset \mathbb{P}(2n - 1, 2n + 2, n - 2, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	6,9
22	$x^4y^2 + zy^n$ $n = 5; 7$	$t^2 + z^3x + x^2y^2 + zy^n \subset \mathbb{P}(3n + 1, n + 2, 3n - 4, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	7,10
23	$az^2xy^3 + bzy^6 + x^4y^2$	$t^2 + z^3x^2 + g(z, x, y) \subset \mathbb{P}(19, 8, 7, 5)$ $\text{Diff} = \emptyset$	5
24	$x^4y^2 + zxy^n$ $n = 4; 5$	$t^2 + z^3x^2 + g(z, x, y) \subset \mathbb{P}(6n - 1, 2n + 2, 3n - 4, 7)$ $\text{Diff} = \emptyset$	4,7
25	$x^4y^2 + z^2y^5$	$t^2 + z^3x + x^2y^2 + z^2y^5 \subset \mathbb{P}(17, 7, 13, 4)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	8
26	$x^3y^3 + y^n$	$t + zx^2 + x^3y^3 + y^n \subset \mathbb{P}(3n, n + 6, n - 3, 3)$	10,18

	$n = 7; 11$	$\mathbb{P}(n+6, n-3, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	
27	$x^3y^3 + y^{2n}$ $n = 4; 5$	$t^2 + zx^2 + x^3y^3 + y^{2n} \subset \mathbb{P}(3n, 2n+6, 2n-3, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	7,9
28	$az^3y^5 + x^3y^3 + bxy^8$	$t + zx^2 + g(z, x, y) \subset \mathbb{P}(21, 11, 5, 2)$ $\mathbb{P}(11, 5, 2), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12
29	$x^3y^3 + zy^n$ $n = 5; 7$	$t + z^3x^2 + x^3y^3 + zy^n \subset \mathbb{P}(9n+6, n+6, 3n-6, 8)$ $\mathbb{P}(n+6, 3n-6, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,10
30	$az^5y + bz^2xy^4 + czy^8 + x^3y^3$	$t + z^3x^2 + g(z, x, y) \subset \mathbb{P}(39, 7, 9, 4)$ $\mathbb{P}(7, 9, 4), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8
31	$x^3y^3 + zy^9$	$t + z^3x^2 + x^3y + zy^3 \subset \mathbb{P}(29, 5, 7, 8)$ $\mathbb{P}(5, 7, 8), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	16
32	$az^4y + bzxxy^4 + x^3y^3$	$t + z^3x^2 + g(z, x, y) \subset \mathbb{P}(33, 7, 6, 5)$ $\mathbb{P}(7, 6, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
33	$az^4y^2 + bzxxy^5 + x^3y^3$	$t^2 + z^3x^2 + g(z, x, y) \subset \mathbb{P}(21, 8, 9, 5)$ $\text{Diff} = \emptyset$	5
34	$az^4y^3 + bzxxy^6 + x^3y^3$	$t + z^3x^2 + g(z, x, y^{1/3}) \subset \mathbb{P}(17, 3, 4, 5)$ $\mathbb{P}(3, 4, 5), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	10
35	$z^2y^4 + x^3y^3$	$t^2 + z^3x^2 + g(z, x, y) \subset \mathbb{P}(24, 10, 9, 7)$ $\text{Diff} = \emptyset$	5
36	$x^3y^3 + z^2y^n$ $n = 5; 7$	$t + z^3x^2 + g(z, x, y) \subset \mathbb{P}(9n+12, n+6, 3n-3, 7)$ $\mathbb{P}(n+6, 3n-3, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,14
37	$x^3y^3 + z^2y^6$	$t^2 + z^3x^2 + x^3y + z^2y^2 \subset \mathbb{P}(11, 4, 5, 7)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	7
38	$x^iy^j f_{7-i-j}(x, y)$ $i \leq 1 \ j \leq 4$	$t + zx^2 + g(z, x, y) \subset \mathbb{P}(7, 5, 1, 1)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
39	$x^7 + y^8$	$t^2 + zx^2 + x^7 + y^2 \subset \mathbb{P}(7, 10, 2, 7)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{3}{4})$	12
40	$x^7 + xy^7$	$t + zx^2 + x^7 + xy \subset \mathbb{P}(7, 5, 1, 6)$ $\mathbb{P}(5, 1, 6), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, \frac{6}{7})$	36
41	$az^2xy^3 + bzy^6 + x^7$	$t + z^3x^2 + g(z, x, y^{1/3}) \subset \mathbb{P}(21, 5, 3, 8)$ $\mathbb{P}(5, 3, 8), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	16
42	$x^7 + zxy^5$	$t + z^3x^2 + x^7 + zxy \subset \mathbb{P}(21, 5, 3, 13)$ $\mathbb{P}(5, 3, 13), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{4}{5})$	26
43	$x^7 + z^2y^4$	$t^2 + z^3x^2 + x^7 + z^2y^2 \subset \mathbb{P}(21, 10, 6, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
44	$x^6y + y^8$	$t^2 + zx + x^3y + y^8 \subset \mathbb{P}(12, 17, 7, 3)$ $\text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	18
45	$x^6y + xy^7$	$t + zx^2 + x^6y + xy^7 \subset \mathbb{P}(41, 29, 6, 5)$ $\mathbb{P}(29, 6, 5), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	30
46	$az^2xy^3 + bzy^6 + x^6y$	$t + z^3x^2 + g(z, x, y) \subset \mathbb{P}(55, 13, 8, 7)$ $\mathbb{P}(13, 8, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
47	$x^6y + zxy^5$	$t + z^3x^2 + x^6y + zxy^5 \subset \mathbb{P}(89, 21, 13, 11)$ $\mathbb{P}(21, 13, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	22
48	$x^6y + z^2y^4$	$t^2 + z^3x + x^3y + z^2y^4 \subset \mathbb{P}(19, 9, 11, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	10
49	$x^5y^2 + y^8$	$t^2 + zx^2 + x^5y + y^4 \subset \mathbb{P}(10, 14, 3, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, \frac{1}{2})$	15
50	$x^5y^2 + xy^7$	$t + zx^2 + x^5y^2 + xy^7 \subset \mathbb{P}(33, 23, 5, 4)$	24

		$\mathbb{P}(23, 5, 4), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	
51	$x^5y^2 + az^2xy^3 + bzy^6$	$t + z^3x^2 + g(z, x, y) \subset \mathbb{P}(47, 11, 7, 6)$ $\mathbb{P}(11, 7, 6), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	12
52	$x^5y^2 + zxy^5$	$t + z^3x^2 + x^5y^2 + zxy^5 \subset \mathbb{P}(73, 17, 11, 9)$ $\mathbb{P}(17, 11, 9), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	18
53	$x^5y^2 + z^2y^4$	$t^2 + z^3x^2 + x^5y + z^2y^2 \subset \mathbb{P}(17, 8, 5, 9)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	9
54	$x^4y^3 + y^8$	$t^2 + zx + x^2y^3 + y^8 \subset \mathbb{P}(8, 11, 5, 2)$ $\text{Diff}=(0, \frac{2}{3}, \frac{1}{2}, 0)$	12
55	$x^4y^3 + xy^7$	$t + zx^2 + x^4y^3 + xy^7 \subset \mathbb{P}(25, 17, 4, 3)$ $\mathbb{P}(17, 4, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18
56	$az^2xy^3 + bzx^5 + czy^6 + dx^4y^3$ $ b + d \neq 0, a + c \neq 0$	$t + z^3x^2 + g(z, x, y^{1/3}) \subset \mathbb{P}(13, 3, 2, 5)$ $\mathbb{P}(3, 2, 5), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	10
57	$x^4y^3 + zxy^5$	$t + z^3x^2 + g(z, x, y) \subset \mathbb{P}(57, 13, 9, 7)$ $\mathbb{P}(13, 9, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
58	$x^4y^3 + z^2y^4$	$t^2 + z^3x^2 + x^2y^3 + z^2y^4 \subset \mathbb{P}(15, 7, 9, 4)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	8
59	$x^3y^4 + y^9$	$t + zx^2 + x^3y^4 + y^9 \subset \mathbb{P}(27, 17, 5, 3)$ $\mathbb{P}(17, 5, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18
60	$x^3y^4 + zy^7$	$t + z^3x^2 + x^3y^4 + zy^7 \subset \mathbb{P}(71, 15, 13, 8)$ $\mathbb{P}(15, 13, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16
61	$az^4y + bzxxy^5 + x^3y^4$	$t + z^3x^2 + g(z, x, y) \subset \mathbb{P}(41, 9, 7, 5)$ $\mathbb{P}(9, 7, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
62	$x^3y^4 + z^2y^5$	$t + z^3x^2 + x^3y^4 + z^2y^5 \subset \mathbb{P}(61, 13, 11, 7)$ $\mathbb{P}(13, 11, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
63	$zx^5 + y^7$	$t + z^3x^2 + zx^5 + y \subset \mathbb{P}(13, 3, 2, 13)$ $\mathbb{P}(3, 2, 13), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{6}{7})$	14
64	$zx^5 + y^8$	$t^2 + z^3x^2 + zx^5 + y^2 \subset \mathbb{P}(13, 6, 4, 13)$ $\text{Diff}=(0, 0, 0, \frac{3}{4})$	13
65	$zx^5 + xy^6$	$t^2 + z^3x^2 + zx^5 + xy^2 \subset \mathbb{P}(13, 6, 4, 11)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	6
66	$zx^5 + xy^7$	$t + z^3x^2 + zx^5 + xy \subset \mathbb{P}(13, 3, 2, 11)$ $\mathbb{P}(3, 2, 11), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{6}{7})$	22
67	$z^2y^4 + azx^5 + bx^3y^4$	$t^2 + z^3x^2 + g(z, x, y^{1/2}) \subset \mathbb{P}(13, 6, 4, 7)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	7
68	$zx^4y + y^7$	$t^2 + z^3x^2 + g(z, x, y) \subset \mathbb{P}(35, 16, 11, 10)$ $\text{Diff} = \emptyset$	5
69	$zx^4y + y^8$	$t^2 + z^3x + zx^2y + y^8 \subset \mathbb{P}(20, 9, 13, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	10
70	$zx^4y + xy^n$ $n = 6; 7$	$t + z^3x^2 + zx^4y + xy^n \subset \mathbb{P}(10n - 3, 2n + 1, 2n - 3, 8)$ $\mathbb{P}(2n + 1, 2n - 3, 8), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6, 16
71	$z^2y^4 + zx^4y + ax^2y^5$	$t^2 + z^3x + g(z, x^{1/2}, y) \subset \mathbb{P}(11, 5, 7, 3)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	6
72	$az^2xy^3 + bzx^4y + czy^6 + dx^3y^4$ $ b + d \neq 0 \& a + c \neq 0$	$t + z^3x^2 + g(z, x, y) \subset \mathbb{P}(31, 7, 5, 4)$ $\mathbb{P}(7, 5, 4), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8
73	$zx^3y^2 + y^n$ $n = 7; 9$	$t + z^3x^2 + zx^3y^2 + y^n \subset \mathbb{P}(7n, n + 4, 2n - 6, 7)$ $\mathbb{P}(n + 4, 2n - 6, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10, 14
74	$zx^3y^2 + y^8$	$t^2 + z^3x^2 + zx^3y + y^4 \subset \mathbb{P}(14, 6, 5, 7)$	7

		Diff=(0, 0, 0, $\frac{1}{2}$)	
75	$az^2y^5 + zx^3y^2 + bxy^7$	$t + z^3x^2 + g(z, x, y) \subset \mathbb{P}(43, 9, 8, 5)$ $\mathbb{P}(9, 8, 5)$, Diff=($\frac{1}{2}$, 0, 0, 0)	10

9. Singularity $-t^2 + z^3y^2 + g(z, x, y)$

1	$x^7 + y^n$ $n = 9; 13$	$t + zy^2 + x + y^n \subset \mathbb{P}(n, n-2, n, 1)$ $\mathbb{P}(n-2, n, 1)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, $\frac{6}{7}$, 0)	28,42
2	$x^7 + y^{10}$	$t^2 + zy^2 + x + y^{10} \subset \mathbb{P}(5, 8, 10, 1)$ $\mathbb{P}(5, 8, 1)$, Diff=(0, $\frac{2}{3}$, $\frac{6}{7}$, 0)	15
3	$x^7 + ay^5z^if_{2-i}(z, y^3)$ $i \leq 1$	$t + z^3y^2 + g(z, x^{1/7}, y) \subset \mathbb{P}(11, 3, 11, 1)$ $\mathbb{P}(3, 11, 1)$, Diff=($\frac{1}{2}$, 0, $\frac{6}{7}$, 0)	14
4	$x^7 + y^{12}$	$t^2 + zy + x + y^6 \subset \mathbb{P}(3, 5, 6, 1)$ $\mathbb{P}(3, 5, 1)$, Diff=(0, $\frac{2}{3}$, $\frac{6}{7}$, $\frac{1}{2}$)	21
5	$x^7 + xy^n$ $n = 7; 11$	$t + zy^2 + x^7 + xy^n \subset \mathbb{P}(7n, 7n-12, n, 6)$ $\mathbb{P}(7n-12, n, 6)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	22,36
6	$x^{3i+1}f_{2-i}(x^3, y^4)$ $i \leq 1$	$t^2 + zy + g(x, y^{1/2}) \subset \mathbb{P}(7, 11, 2, 3)$ Diff=(0, $\frac{2}{3}$, 0, $\frac{1}{2}$)	9
7	$x^{2i+1}f_{3-i}(x^2, y^3)$ $i \leq 2$	$t + zy^2 + g(x, y) \subset \mathbb{P}(21, 17, 3, 2)$ $\mathbb{P}(17, 3, 2)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	12
8	$x^{3i+1}f_{2-i}(x^3, y^5)$ $i \leq 1$	$t + zy^2 + g(x, y) \subset \mathbb{P}(35, 29, 5, 3)$ $\mathbb{P}(29, 5, 3)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	18
9	$x^7 + x^2y^6$	$t^2 + zy^2 + x^7 + x^2y^6 \subset \mathbb{P}(21, 32, 6, 5)$ Diff=(0, $\frac{2}{3}$, 0, 0)	9
10	$azx^3y^3 + x^7 + bx^2y^7$	$t + z^3y^2 + g(z, x, y) \subset \mathbb{P}(49, 13, 7, 5)$ $\mathbb{P}(13, 7, 5)$, Diff=($\frac{1}{2}$, 0, 0, 0)	10
11	$x^7 + x^2y^8$	$t^2 + zy + x^7 + x^2y^4 \subset \mathbb{P}(14, 23, 4, 5)$ Diff=(0, $\frac{2}{3}$, 0, $\frac{1}{2}$)	12
12	$x^7 + x^2y^9$	$t + zy^2 + x^7 + x^2y^9 \subset \mathbb{P}(63, 53, 9, 5)$ $\mathbb{P}(53, 9, 5)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	30
13	$bzx^2y^4 + x^7 + cx^3y^5$	$t + z^3y^2 + g(z, x, y) \subset \mathbb{P}(35, 9, 5, 4)$ $\mathbb{P}(9, 5, 4)$, Diff=($\frac{1}{2}$, 0, 0, 0)	8
14	$x^7 + x^3y^7$	$t + zy^2 + x^7 + x^3y^7 \subset \mathbb{P}(49, 41, 7, 4)$ $\mathbb{P}(41, 7, 4)$, Diff=($\frac{1}{2}$, $\frac{2}{3}$, 0, 0)	24
15	$x^7 + zy^n$ $n = 7; 9$	$t + z^3y^2 + x + zy^n \subset \mathbb{P}(3n-2, n-2, 3n-2, 2)$ $\mathbb{P}(n-2, 3n-2, 2)$, Diff=($\frac{1}{2}$, 0, $\frac{6}{7}$, 0)	22,28
16	$x^7 + zxy^n$ $n = 5; 7$	$t + z^3y^2 + x^7 + zxy^n \subset \mathbb{P}(21n-14, 7n-12, 3n-2, 11)$ $\mathbb{P}(7n-12, 3n-2, 11)$, Diff=($\frac{1}{2}$, 0, 0, 0)	16,22
17	$x^7 + zxy^6$	$t^2 + z^3y + x^7 + zxy^3 \subset \mathbb{P}(28, 15, 8, 11)$ Diff=(0, 0, 0, $\frac{1}{2}$)	11
18	$x^7 + zx^2y^5$	$t + z^3y^2 + x^7 + zx^2y^5 \subset \mathbb{P}(91, 25, 13, 8)$ $\mathbb{P}(25, 13, 8)$, Diff=($\frac{1}{2}$, 0, 0, 0)	16
19	$zx^5 + y^n$ $n = 7; 13$	$t + z^3y^2 + zx + y^n \subset \mathbb{P}(3n, n-2, 2n+2, 3)$ $\mathbb{P}(n-2, 2n+2, 3)$, Diff=($\frac{1}{2}$, 0, $\frac{4}{5}$, 0)	15,30
20	$zx^5 + czx^2y^4 +$ $+ex^6y + dx^3y^5 + ly^9$ $ c + d + l \neq 0$	$t + z^3y^2 + g(z, x, y) \subset \mathbb{P}(27, 7, 4, 3)$ $\mathbb{P}(7, 4, 3)$, Diff=($\frac{1}{2}$, 0, 0, 0)	6

21	$zx^5 + y^{10}$	$t^2 + z^3y^2 + zx + y^{10} \subset \mathbb{P}(15, 8, 22, 3)$ Diff = $(0, 0, \frac{4}{5}, 0)$	11
22	$by^5z^if_{3-i}(z, y^3) + zx^5 + ax^5y^3, i \leq 2 \parallel a \neq 0$	$t + z^3y^2 + g(z, x^{1/5}, y) \subset \mathbb{P}(11, 3, 8, 1)$ $\mathbb{P}(3, 8, 1), \text{Diff} = (\frac{1}{2}, 0, \frac{4}{5}, 0)$	10
23	$zx^5 + y^{12}$	$t^2 + z^3y^2 + zx + y^6 \subset \mathbb{P}(9, 5, 13, 3)$ Diff = $(0, 0, \frac{4}{5}, \frac{1}{2})$	15
24	$zx^5 + xy^{4n+2}$ $n = 1; 2$	$t^2 + z^3y + g(z, x, y^{1/2}) \subset \mathbb{P}(15n + 8, 10n + 1, 4n + 3, 13)$ Diff = $(0, 0, 0, \frac{1}{2})$	9, 13
25	$zx^5 + xy^n$ $n = 7; 9; 11$	$t + z^3y^2 + zx^5 + xy^n \subset \mathbb{P}(15n + 2, 5n - 8, 2n + 2, 13)$ $\mathbb{P}(5n - 8, 2n + 2, 13), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	16, 22, 26
26	$zx^5 + xy^8$	$t^2 + z^3y^2 + g(z, x, y) \subset \mathbb{P}(61, 32, 18, 13)$ Diff = $\emptyset$	9
27	$zx^5 + x^2y^n$ $n = 5; 7; 9$	$t + z^3y^2 + zx^5 + x^2y^n \subset \mathbb{P}(15n + 4, 5n - 6, 2n + 2, 11)$ $\mathbb{P}(5n - 6, 2n + 2, 11), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	12, 18, 22
28	$zx^5 + x^2y^6$	$t^2 + z^3y^2 + g(z, x, y) \subset \mathbb{P}(47, 24, 14, 11)$ Diff = $\emptyset$	7
29	$zx^5 + x^2y^8$	$t^2 + z^3y + zx^5 + x^2y^4 \subset \mathbb{P}(31, 17, 9, 11)$ Diff = $(0, 0, 0, \frac{1}{2})$	11
30	$zx^5 + x^3y^4$	$t^2 + z^3y^2 + g(z, x, y) \subset \mathbb{P}(33, 16, 10, 9)$ Diff = $\emptyset$	5
31	$zx^5 + x^3y^6$	$t^2 + z^3y + zx^5 + x^3y^3 \subset \mathbb{P}(24, 13, 7, 9)$ Diff = $(0, 0, 0, \frac{1}{2})$	9
32	$zx^5 + x^3y^7$	$t + z^3y^2 + zx^5 + x^3y^7 \subset \mathbb{P}(111, 31, 16, 9)$ $\mathbb{P}(31, 16, 9), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	18
33	$zx^5 + x^4y^n$ $n = 3; 5$	$t + z^3y^2 + zx^5 + x^4y^n \subset \mathbb{P}(15n + 8, 5n - 2, 2n + 2, 7)$ $\mathbb{P}(5n - 2, 2n + 2, 7), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	10, 14
34	$zx^5 + x^4y^4$	$t^2 + z^3y + zx^5 + x^4y^2 \subset \mathbb{P}(17, 9, 5, 7)$ Diff = $(0, 0, 0, \frac{1}{2})$	7

10. Singularity – $t^2 + z^3xy + g(z, x, y)$

1	$x^5 + y^{2n}$ $n = 3; 6; 8, 9$	$t^2 + zxy + g(x, y) \subset \mathbb{P}(5n, 8n - 5, 2n, 5)$ Diff = $(0, \frac{2}{3}, 0, 0)$	7, 9, 15
2	$x^5 + y^n$ $n = 7; 9; 13; 19$	$t + zxy + x^5 + y^n \subset \mathbb{P}(5n, 4n - 5, n, 5)$ $\mathbb{P}(4n - 5, n, 5), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	10, 14, 22 30
3	$ax^2y^{2n+1} + x^5 + by^{6n+2}$ $n = 1 \ b \neq 0; 2$	$t^2 + z^3xy + g(z, x, y) \subset \mathbb{P}(15n + 5, 8n + 1, 6n + 2, 5)$ Diff = $\emptyset$	4, 5
4	$x^i(x + y^2)^j f_{5-i-j}(x, y^2)$ $i \leq 2 \& j \leq 4$	$t^2 + zxy + g(x, y) \subset \mathbb{P}(5, 7, 2, 1)$ Diff = $(0, \frac{2}{3}, 0, 0)$	3
5	$ax^2y^{2n} + x^5 + by^{6n-1}$ $n = 2; 3$	$t + z^3xy + g(z, x, y) \subset \mathbb{P}(30n - 5, 8n - 3, 6n - 1, 5)$ $\mathbb{P}(8n - 3, 6n - 1, 5), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	10
6	$x^if_{5-i}(x, y^3)$ $i \leq 3$	$t + zxy + g(x, y) \subset \mathbb{P}(15, 11, 3, 1)$ $\mathbb{P}(11, 3, 1), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
7	$x^5 + x^2y^{2n}$ $n = 4, 5$	$t^2 + zxy + g(x, y) \subset \mathbb{P}(5n, 8n - 3, 2n, 3)$ Diff = $(0, \frac{2}{3}, 0, 0)$	9
8	$x^5 + x^2y^n$ $n = 5, 7; 11$	$t + zxy + g(x, y) \subset \mathbb{P}(5n, 4n - 3, n, 3)$ $\mathbb{P}(4n - 3, n, 3), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	10, 18
9	$x^5 + zy^{2n+1}$ $n = 2; 3; 4, 5, 6$	$t^2 + z^3xy + g(z, x, y) \subset \mathbb{P}(15n + 5, 8n - 1, 6n + 2, 11)$ Diff = $\emptyset$	3, 5, 11

10	$x^5 + zy^n$ $n = 6; 8, 10; 12, 14$	$t + z^3xy + g(z, x, y) \subset \mathbb{P}(15n - 5, 4n - 5, 3n - 1, 11)$ $\mathbb{P}(4n - 5, 3n - 1, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,14,22
11	$x^5 + z^2y^{2n}$ $n = 2; 3, 4$	$t^2 + z^3xy + g(z, x, y) \subset \mathbb{P}(15n - 5, 8n - 5, 6n - 2, 7)$ $\text{Diff} = \emptyset$	6,7
12	$x^5 + z^2y^n$ $n = 5; 7, 9$	$t + z^3xy + g(z, x, y) \subset \mathbb{P}(15n - 10, 4n - 5, 3n - 2, 7)$ $\mathbb{P}(4n - 5, 3n - 2, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,14
13	$x^i y^j (x + y)^l f_{6-i-j-l}(x, y)$ $i \leq 1 \& j \leq 1 \& l \leq 3$	$t^2 + zxy + g(x, y) \subset \mathbb{P}(3, 4, 1, 1)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	3
14	$x^6 + y^n$ $n = 7; 11$	$t^2 + zxy + g(x, y) \subset \mathbb{P}(3n, 5n - 6, n, 6)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	10,18
15	$x^{3i} f_{2-i}(x^3, y^n)$ $n = 4 \ i = 0; 5$	$t^2 + zxy + g(x, y) \subset \mathbb{P}(3n, 5n - 3, n, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	6,9
16	$x^{2i} f_{3-i}(x^2, y^3)$ $i \leq 1$	$t^2 + zxy + g(x, y) \subset \mathbb{P}(9, 13, 3, 2)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	6
17	$az^2y^4 + bzx^3y^2 + x^6 +$ $+cx^2y^5, a + c \neq 0$	$t^2 + z^3xy + g(z, x, y) \subset \mathbb{P}(15, 7, 5, 4)$ $\text{Diff} = \emptyset$	4
18	$x^6 + x^2y^7$	$t^2 + zxy + g(x, y) \subset \mathbb{P}(21, 31, 7, 4)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
19	$x^6 + zy^n$ $n = 5; 6; 7; 8$	$t^2 + z^3xy + g(z, x, y) \subset \mathbb{P}(9n - 3, 5n - 6, 3n - 1, 13)$ $\text{Diff} = \emptyset$	7,8,10 13
20	$x^6 + zx^2y^4$	$t^2 + z^3xy + g(z, x, y) \subset \mathbb{P}(33, 16, 11, 7)$ $\text{Diff} = \emptyset$	7
21	$x^6 + z^2y^5$	$t^2 + z^3xy + g(z, x, y) \subset \mathbb{P}(39, 19, 13, 8)$ $\text{Diff} = \emptyset$	8
22	$x^i y^j (x + y)^l f_{7-i-j-l}(x, y)$ $i \leq 2 \& j \leq 2 \& l \leq 4$	$t + zxy + g(x, y) \subset \mathbb{P}(7, 5, 1, 1)$ $\mathbb{P}(5, 1, 1), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
23	$x^7 + y^8$	$t^2 + zxy + x^7 + y^8 \subset \mathbb{P}(28, 41, 8, 7)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
24	$x^7 + y^9$	$t + zxy + x^7 + y^9 \subset \mathbb{P}(63, 47, 9, 7)$ $\mathbb{P}(47, 9, 7), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	42
25	$x^7 + x^2y^6$	$t^2 + zxy + x^7 + x^2y^6 \subset \mathbb{P}(21, 31, 6, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	15
26	$x^7 + zy^6$	$t + z^3xy + x^7 + zy^6 \subset \mathbb{P}(119, 29, 17, 15)$ $\mathbb{P}(29, 17, 15), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
27	$x^7 + zx^2y^4$	$t + z^3xy + x^7 + zx^2y^4 \subset \mathbb{P}(77, 19, 11, 9)$ $\mathbb{P}(19, 11, 9), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	18
28	$x^7 + z^2y^4$	$t^2 + z^3xy + x^7 + z^2y^4 \subset \mathbb{P}(35, 17, 10, 9)$ $\text{Diff} = \emptyset$	9
29	$x^5y^2 + y^{2n}$ $n = 4; 5$	$t^2 + zxy + x^5y^2 + y^{2n} \subset \mathbb{P}(5n, 8n - 3, 2n - 2, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9,15
30	$axx^2y^4 + x^5y^2 + by^9$	$t + z^3xy + g(z, x, y) \subset \mathbb{P}(45, 11, 7, 5)$ $\mathbb{P}(11, 7, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
31	$x^5y^2 + y^{11}$	$t + zxy + x^5y^2 + y^{11} \subset \mathbb{P}(55, 41, 9, 5)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	30
32	$x^5y^2 + x^2y^6$	$t^2 + zxy + x^5y^2 + x^2y^6 \subset \mathbb{P}(13, 19, 4, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
33	$x^5y^2 + x^2y^7$	$t + zxy + x^5y^2 + x^2y^7 \subset \mathbb{P}(31, 23, 5, 3)$ $\mathbb{P}(23, 5, 3), \text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18

34	$x^5y^2 + zy^n$ $n = 6; 8$	$t + z^3xy + g(z, x, y) \subset \mathbb{P}(15n - 3, 4n - 3, 3n - 5, 11)$ $\mathbb{P}(4n - 3, 3n - 5, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16,22
35	$x^5y^2 + zy'$	$t^2 + z^3xy + x^5y^2 + zy' \subset \mathbb{P}(51, 25, 16, 11)$ $\text{Diff} = \emptyset$	11
36	$x^5y^2 + z^2y^4$	$t^2 + z^3xy + x^5y^2 + z^2y^4 \subset \mathbb{P}(27, 13, 8, 7)$ $\text{Diff} = \emptyset$	7
37	$x^5y^2 + z^2y^5$	$t + z^3xy + x^5y^2 + z^2y^5 \subset \mathbb{P}(69, 17, 11, 7)$ $\mathbb{P}(17, 11, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
38	$zx^5 + y^n$ $n = 7; 8; 9$	$t^2 + z^3xy + zx^5 + y^n \subset \mathbb{P}(7n, 4n - 5, 2n + 1, 14)$ $\text{Diff} = \emptyset$	5,7,14
39	$zx^5 + x^2y^n$ $n = 5; 6$	$t^2 + z^3xy + zx^5 + x^2y^n \subset \mathbb{P}(7n + 1, 4n - 3, 2n + 1, 10)$ $\text{Diff} = \emptyset$	5,6
40	$z^2y^4 + zx^5 + bx^4y^3$	$t^2 + z^3xy + g(z, x, y) \subset \mathbb{P}(23, 11, 7, 6)$ $\text{Diff} = \emptyset$	6

11. Singularity –

$$t^2 + f(z, x) + g(z, x, y) = t^2 + z^i x^j f_{5-i-j}(z, x) + g(z, x, y), (i \leq j)$$

1	y^{2n} $n = 3 \ j = 1; 4 \ j \leq 2$	$t^2 + f(z, x) + y^2 \subset \mathbb{P}(5, 2, 2, 5)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	3,5
2	y^n $(j \leq 2) \ \& \ (n = 7; 9)$	$t + f(z, x) + y \subset \mathbb{P}(5, 1, 1, 5)$ $\mathbb{P}(1, 1, 5), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	8,10
3	$zy^n, n = 5 \& j = 1;$ $n = 7 \ \& \ i \leq 2 \ \& \ j \leq 3$	$t + f(z, x) + zy \subset \mathbb{P}(5, 1, 1, 4)$ $\mathbb{P}(1, 1, 4), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	6,8
4	$y^3 f_3'(z, x) + zy^6$	$t + f(z, x) + g(z, x, y^{1/3}) \subset \mathbb{P}(5, 1, 1, 2)$ $\mathbb{P}(1, 1, 2), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	6
(1)	$zxy^4 \ \ z^2y^4$ $j \leq 1 \ \ (i \leq 1 \& j \leq 2)$	$t^2 + f(z, x) + g(z, x, y^{1/2}) \subset \mathbb{P}(5, 2, 2, 3)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	3
6	$zxy^5 \ \ z^2y^5$ $j \leq 2 \ \ (i \leq 1 \& j \leq 2)$	$t + f(z, x) + g(z, x, y^{1/5}) \subset \mathbb{P}(5, 1, 1, 3)$ $\mathbb{P}(1, 1, 3), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{4}{5})$	6

(1) The singularity is $t^2 + z^i x^j f_{5-i-j}(z, x) + ay^3 z^k x^l f_{3-k-l}(z, x) + bzy^6$. Assume that $f_5 \in \mathcal{M}_1$ after each quasihomogeneous coordinate change. The canonical singularity of this kind is exceptional, except the cases: $(i = 3 \& k = 1)$; $(i = 2 \& k \geq 1)$; $(j = 3 \& l = 1)$.

12. Singularity – $t^2 + g(z, x, y)$, where $g_5 \in \mathcal{M}_1$

1	$z^5 + zx^3y + y^{2n}$ $n = 3; 6; 8, 9$	$t^2 + z^5 + zxy + y^{2n} \subset \mathbb{P}(5n, 2n, 8n - 5, 5)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	7,9,15
2	$z^5 + zx^3y + y^n$ $n = 7; 9; 13; 19$	$t + z^5 + zxy + y^n \subset \mathbb{P}(5n, n, 4n - 5, 5)$ $\mathbb{P}(n, 4n - 5, 5), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	10,14,22 30
3	$z^5 + zx^3y + az^2xy^{2n+1} + by^{6n+2}, n = 1 \ b \neq 0; 2$	$t^2 + g(z, x, y) \subset \mathbb{P}(15n + 5, 6n + 2, 8n + 1, 5)$ $\text{Diff} = \emptyset$	4,5
4	$z^5 + zx^3y + y^4 z^i f_{3-i}(z, y^2) + ax^3y^3, i \leq 2 a \neq 0$	$t^2 + g(z, x^{1/3}, y) \subset \mathbb{P}(5, 2, 7, 1)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	3
5	$z^5 + zx^3y + az^2xy^{n+1} + by^{3n+2}, n = 3, 5$	$t + g(z, x, y) \subset \mathbb{P}(15n + 10, 3n + 2, 4n + 1, 5)$ $\mathbb{P}(3n + 2, 4n + 1, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
6	$z^5 + zx^3y + by^6 z^i f_{3-i}(z, y^3) +$	$t + g(z, x^{1/3}, y) \subset \mathbb{P}(15, 3, 11, 1)$	6

	ax^3y^4	$\mathbb{P}(3, 11, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	
7	$z^5 + zx^3y + xy^{2n+1}$ $n = 2; 3; 4, 5, 6$	$t^2 + g(z, x, y) \subset \mathbb{P}(15n + 5, 6n + 2, 8n - 1, 11)$ $\text{Diff} = \emptyset$	3,5,11
8	$z^5 + zx^3y + xy^n$ $n = 6; 8, 10; 12, 14$	$t + g(z, x, y) \subset \mathbb{P}(15n - 5, 3n - 1, 4n - 5, 11)$ $\mathbb{P}(3n - 1, 4n - 5, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,14,22
9	$z^5 + zx^3y + x^2y^{2n}$ $n = 2; 3, 4$	$t^2 + g(z, x, y) \subset \mathbb{P}(15n - 5, 6n - 2, 8n - 5, 7)$ $\text{Diff} = \emptyset$	6,7
10	$z^5 + zx^3y + x^2y^n$ $n = 5; 7, 9$	$t + g(z, x, y) \subset \mathbb{P}(15n - 10, 3n - 2, 4n - 5, 7)$ $\mathbb{P}(3n - 2, 4n - 5, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,14
11	$z^5 + zx^3y + z^2y^{2n}$ $n = 4, 5$	$t^2 + z^5 + zxy + z^2y^{2n} \subset \mathbb{P}(5n, 2n, 8n - 3, 3)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	9
12	$z^5 + zx^3y + z^2y^n$ $n = 5, 7; 11$	$t + z^5 + zxy + z^2y^n \subset \mathbb{P}(5n, n, 4n - 3, 3)$ $\mathbb{P}(n, 4n - 3, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	10,18
13	$z^5 + x^4y + y^{2n}$ $n = 3, 4; 6; 7$	$t^2 + z + xy + y^{2n} \subset \mathbb{P}(n, 2n, 2n - 1, 1)$ $\mathbb{P}(n, 2n - 1, 1), \text{Diff}=(0, \frac{4}{5}, \frac{3}{4}, 0)$	5,12,20
14	$z^5 + x^4y + ax^2y^{n+1} + by^{2n+1}, n = 3, 5$	$t^2 + g(z^{1/5}, x, y) \subset \mathbb{P}(2n + 1, 4n + 2, n, 2)$ $\mathbb{P}(2n + 1, n, 2), \text{Diff}=(0, \frac{4}{5}, 0, 0)$	5
15	$z^5 + x^4y + ay^n x^i f_{3-i}(x, y^{n-1})$ $n = 3 \ i \leq 2; 4$	$t + g(z^{1/5}, x, y) \subset \mathbb{P}(4n - 3, 4n - 3, n - 1, 1)$ $\mathbb{P}(4n - 3, n - 1, 1), \text{Diff}=(\frac{1}{2}, \frac{4}{5}, 0, 0)$	6,10
16	$z^5 + x^4y + ay^4 x^i f_{3-i}(z, y^2)$	$t^2 + g(z, x^{1/4}, y) \subset \mathbb{P}(5, 2, 9, 1)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	4
17	$z^5 + x^4y + xy^n$ $n = 5; 6, 8; 11$	$t + z + x^4y + xy^n \subset \mathbb{P}(4n - 1, 4n - 1, n - 1, 3)$ $\mathbb{P}(4n - 1, n - 1, 3), \text{Diff}=(\frac{1}{2}, \frac{4}{5}, 0, 0)$	6,10,30
18	$z^5 + x^4y + az^3xy^2 + bz^2y^7$ $czx^2y^4 + dxy^9, b + c + d \neq 0$	$t + g(z, x, y) \subset \mathbb{P}(35, 7, 8, 3)$ $\mathbb{P}(7, 8, 3), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
19	$z^5 + x^4y + zy^n$ $n = 5; 7, 9; 11$	$t^2 + z^5 + x^2y + zy^n \subset \mathbb{P}(5n, 2n, 5n - 4, 8)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	3,12,16
20	$z^5 + x^4y + az^3y^n + bzy^{2n}, n = 3 \ b \neq 0; 5$	$t^2 + g(z, x^{1/2}, y) \subset \mathbb{P}(5n, 2n, 5n - 2, 4)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	4,8
21	$z^5 + x^4y + zxy^n$ $n = 4, 5; 6, 7; 8$	$t + g(z, x, y) \subset \mathbb{P}(20n - 5, 4n - 1, 5n - 4, 11)$ $\mathbb{P}(4n - 1, 5n - 4, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,18,22
22	$z^5 + x^4y + az^2y^5 + bx^2y^3$	$t^2 + g(z, x, y) \subset \mathbb{P}(25, 10, 11, 6)$ $\text{Diff} = \emptyset$	3
23	$z^5 + x^4y + z^2y^{2n}$ $n = 2; 3$	$t^2 + z^5 + xy + z^2y^{2n} \subset \mathbb{P}(5n, 2n, 10n - 3, 3)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	10,12
24	$z^5 + x^4y + z^2xy^n$ $n = 3, 4; 5$	$t + g(z, x, y) \subset \mathbb{P}(20n - 5, 4n - 1, 5n - 3, 7)$ $\mathbb{P}(4n - 1, 5n - 3, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,14
25	$z^4x + zx^3y + y^{2n}$ $n = 3, 4; 5; 6, 7$	$t^2 + g(z, x, y) \subset \mathbb{P}(11n, 4n + 1, 6n - 4, 11)$ $\text{Diff} = \emptyset$	5,7,11
26	$z^4x + zx^3y + y^n$ $n = 7; 9; 11; 13; 15$	$t + g(z, x, y) \subset \mathbb{P}(11n, 2n + 1, 3n - 4, 11)$ $\mathbb{P}(2n + 1, 3n - 4, 11), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,12,16 18,22
27	$z^4x + zx^3y + az^3y^{2n-1} + bx^2y^{2n}, n = 2 \ ab \neq 0; 3$	$t^2 + g(z, x, y) \subset \mathbb{P}(11n - 4, 4n - 1, 6n - 4, 5)$ $\text{Diff} = \emptyset$	4,5
28	$z^4x + zx^3y + az^3y^{n-1} + bx^2y^n, n = 5 \ b \neq 0; 7$	$t + g(z, x, y) \subset \mathbb{P}(11n - 8, 2n - 1, 3n - 4, 5)$ $\mathbb{P}(2n - 1, 3n - 4, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,10
29	$z^4x + zx^3y + z^2y^{2n}$ $n = 3, 4$	$t^2 + g(z, x, y) \subset \mathbb{P}(11n + 1, 4n + 1, 6n - 2, 7)$ $\text{Diff} = \emptyset$	7
30	$z^4x + zx^3y + z^2y^n$	$t + g(z, x, y) \subset \mathbb{P}(11n + 2, 2n + 1, 3n - 2, 7)$	8,10,14

	$n = 5; 7; 9$	$\mathbb{P}(2n+1, 3n-2, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	
31	$z^4x + x^4y + y^n$ $n = 6, 8; 10; 12$	$t^2 + zx + x^4y + y^n \subset \mathbb{P}(2n, 3n+1, n-1, 4)$ $\text{Diff}=(0, \frac{3}{4}, 0, 0)$	5,12,16
32	$z^4x + x^4y + ax^2y^{n+1} + by^{2n+1}, n = 3 \ b \neq 0; 5$	$t^2 + g(z^{1/4}, x, y) \subset \mathbb{P}(2n+1, 3n+2, n, 2)$ $\text{Diff}=(0, \frac{3}{4}, 0, 0)$	4,8
33	$z^4x + x^4y + by^3x^i f_{3-i}(x, y^2) + az^4y^2, i \leq 2 \ a \neq 0$	$t^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(9, 7, 4, 2)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	4
34	$z^4x + x^4y + zy^n$ $n = 5; 6, 7, 8; 9; 10$	$t + g(z, x, y) \subset \mathbb{P}(16n+1, 3n+1, 4n-3, 13)$ $\mathbb{P}(3n+1, 4n-3, 13), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,10,22 26
35	$z^4x + x^4y + az^2y^{2n-1} + bzx^2y^n, n = 3 \ a \neq 0; 4$	$t + g(z, x, y) \subset \mathbb{P}(16n-7, 3n-1, 4n-3, 5)$ $\mathbb{P}(3n-1, 4n-3, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,10
36	$z^4x + x^4y + z^2y^n$ $n = 4, 6$	$t^2 + g(z, x, y) \subset \mathbb{P}(8n+1, 3n+1, 4n-2, 10)$ $\text{Diff} = \emptyset$	5
37	$z^4x + x^4y + z^3y^n$ $n = 4; 5$	$t + g(z, x, y) \subset \mathbb{P}(16n+3, 3n+1, 4n-1, 7)$ $\mathbb{P}(3n+1, 4n-1, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,14
38	$z^4y + z^2x^3 + y^{10}$	$t^2 + z^2y + zx + y^{10} \subset \mathbb{P}(10, 9, 11, 2)$ $\text{Diff}=(0, \frac{1}{2}, \frac{2}{3}, 0)$	12
39	$z^4y + z^2x^3 + az^2y^4 + bx^3y^3 + cy^7, ab + c \neq 0$	$t^2 + g(z, x^{1/3}, y) \subset \mathbb{P}(7, 3, 8, 2)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	3
40	$z^4y + z^2x^3 + az^2xy^3 + bx^4y^2 + cx^2y^5 + dy^8, ab + c + d \neq 0$	$t^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(8, 7, 3, 2)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	4
41	$z^4y + z^2x^3 + cy^5z^i f_{2-i}(z, y^2) + azx^3y^2 + bx^3y^4$ $ a + b + c \neq 0 \& i \leq 1$	$t + g(z, x^{1/3}, y) \subset \mathbb{P}(9, 2, 5, 1)$ $\mathbb{P}(2, 5, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	6
42	$z^4y + z^2x^3 + xy^n$ $n = 5; 7$	$t^2 + g(z, x, y) \subset \mathbb{P}(6n+1, 3n-2, 2n+2, 10)$ $\text{Diff} = \emptyset$	3,5
43	$z^4y + z^2x^3 + azx^2y^n + bxy^{2n}, n = 3; 4$	$t + g(z, x, y) \subset \mathbb{P}(12n+1, 3n-1, 2n+1, 5)$ $\mathbb{P}(3n-1, 2n+1, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,10
44	$z^4y + z^2x^3 + x^2y^n$ $n = 4; 6$	$t^2 + g(z^{1/2}, x, y) \subset \mathbb{P}(3n+1, 3n-1, n+1, 4)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	6,8
45	$z^4y + z^2x^3 + zy^8$	$t + z^4y + z^2x + zy^8 \subset \mathbb{P}(31, 7, 17, 3)$ $\mathbb{P}(7, 17, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	18
46	$z^4y + z^2x^3 + zxy^n$ $n = 5; 6$	$t + g(z, x, y) \subset \mathbb{P}(12n-1, 3n-2, 2n+1, 7)$ $\mathbb{P}(3n-2, 2n+1, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	12,14
47	$z^4y + zx^4 + y^n$ $n = 6, 8; 12; 16$	$t^2 + z^4y + zx + y^n \subset \mathbb{P}(2n, n-1, 3n+1, 4)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	5,12,16
48	$z^4y + zx^4 + az^2y^{n+1} + by^{2n+1}, n = 3 \ b \neq 0; 5$	$t^2 + g(z, x^{1/4}, y) \subset \mathbb{P}(2n+1, n, 3n+2, 2)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	4,8
49	$z^4y + zx^4 + by^5z^i f_{2-i}(z, y^2) + ax^4y^2, i \leq 2$	$t^2 + g(z, x^{1/2}, y) \subset \mathbb{P}(9, 4, 7, 2)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	4
50	$z^4y + zx^4 + xy^n$ $n = 5; 6, 7, 8; 9; 10$	$t + g(z, x, y) \subset \mathbb{P}(16n+1, 4n-3, 3n+1, 13)$ $\mathbb{P}(4n-3, 3n+1, 13), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,10,22 26
51	$z^4y + zx^4 + x^2y^n$ $n = 4, 6$	$t^2 + g(z, x, y) \subset \mathbb{P}(8n+1, 4n-2, 3n+1, 10)$ $\text{Diff} = \emptyset$	5
52	$z^4y + zx^4 + x^2y^{2n+1}$ $n = 2; 3$	$t + g(z, x, y) \subset \mathbb{P}(16n+9, 4n+1, 3n+2, 5)$ $\mathbb{P}(4n+1, 3n+2, 5), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,10
53	$z^4y + zx^4 + x^3y^n$ $n = 4; 5$	$t + g(z, x, y) \subset \mathbb{P}(16n+3, 4n-1, 3n+1, 7)$ $\mathbb{P}(4n-1, 3n+1, 7), \text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,14

54	$z^3x^2 + zx^3y + y^{2n}$ $n = 4; 5$	$t^2 + g(z, x, y) \subset \mathbb{P}(7n, 2n + 2, 4n - 3, 7)$ $\text{Diff} = \emptyset$	5,7
55	$z^3x^2 + zx^3y + y^n$ $n = 7; 9; 11$	$t + g(z, x, y) \subset \mathbb{P}(7n, n + 2, 2n - 3, 7)$ $\mathbb{P}(n + 2, 2n - 3, 7), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	8,10,14
56	$z^3x^2 + zx^3y + az^2y^{2n} +$ $bxy^{2n+1}, n = 2 \text{ } ab \neq 0; 3$	$t^2 + g(z, x, y) \subset \mathbb{P}(7n + 2, 2n + 2, 4n - 1, 5)$ $\text{Diff} = \emptyset$	3,5
57	$z^3x^2 + zx^3y + az^2y^{n-1} +$ $bxy^n, n = 6 \text{ } a \neq 0; 8$	$t + g(z, x, y) \subset \mathbb{P}(7n - 3, n + 1, 2n - 3, 5)$ $\mathbb{P}(n + 1, 2n - 3, 5), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	6,10
58	$z^3x^2 + zx^3y + az^4y^2 + bz^2xy^3 +$ $czy^6 + dx^2y^4, c + d \neq 0 \text{ \& }$ $ ad + b + c \neq 0$	$t^2 + g(z, x, y) \subset \mathbb{P}(11, 4, 5, 3)$ $\text{Diff} = \emptyset$	3
59	$z^3x^2 + zx^3y + az^4y^3 + bz^2xy^4 +$ $czy^8 + dx^2y^5, ad + b + c \neq 0$	$t + g(z, x, y) \subset \mathbb{P}(29, 5, 7, 3)$ $\mathbb{P}(5, 7, 3), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	6
60	$z^3x^2 + x^3y^2 + y^{10}$	$t^2 + zx^2 + x^3y^2 + y^{10} \subset \mathbb{P}(15, 14, 8, 3)$ $\text{Diff} = (0, \frac{2}{3}, 0, 0)$	7
61	$z^3x^2 + x^3y^2 + y^n$ $n = 7; 9; 13$	$t + zx^2 + x^3y^2 + y^n \subset \mathbb{P}(3n, n + 4, n - 2, 3)$ $\mathbb{P}(n + 4, n - 2, 3), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	9,12,18
62	$z^3x^2 + x^3y^2 + az^6y + bz^3y^6 +$ $cz^3xy^3 + dx^2y^5 + exy^8 + ly^{11}$ $ ad + b + c + e + l \neq 0$	$t + g(z^{1/3}, x, y) \subset \mathbb{P}(11, 5, 3, 1)$ $\mathbb{P}(5, 3, 1), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
63	$z^3x^2 + x^3y^2 + y^{12}$	$t^2 + zx^2 + x^3y + y^6 \subset \mathbb{P}(9, 8, 5, 3)$ $\text{Diff} = (0, \frac{2}{3}, 0, \frac{1}{2})$	9
64	$z^3x^2 + x^3y^2 + az^3y^{n-2} +$ $bxy^n, n = 5 \text{ } a \neq 0; 9$	$t + g(z^{1/3}, x, y) \subset \mathbb{P}(3n - 2, n + 2, n - 2, 2)$ $\mathbb{P}(n + 2, n - 2, 2), \text{Diff} = (\frac{1}{2}, \frac{2}{3}, 0, 0)$	6,12
65	$z^3x^2 + x^3y^2 + az^5y^2 + bz^4xy +$ $(1) \text{ } cz^3y^5 + dz^2xy^4 + ezx^2y^3 +$ $lzy^8 + nxy^7$	$t + g(z, x, y) \subset \mathbb{P}(19, 3, 5, 2)$ $\mathbb{P}(3, 5, 2), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	4
66	$z^3x^2 + x^3y^2 + zy^n$ $n = 5; 7; 9; 11$	$t + g(z, x, y) \subset \mathbb{P}(9n + 4, n + 4, 3n - 4, 8)$ $\mathbb{P}(n + 4, 3n - 4, 8), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	7,10,14 16
67	$z^3x^2 + x^3y^2 + az^2xy^n +$ $bzy^{2n}, n = 3; 5$	$t + g(z, x, y) \subset \mathbb{P}(9n + 2, n + 2, 3n - 2, 4)$ $\mathbb{P}(n + 2, 3n - 2, 4), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	6,8
68	$z^3x^2 + x^3y^2 + az^4y^2 + bzxxy^4$ $ab \neq 0$	$t^2 + g(z, x, y) \subset \mathbb{P}(17, 6, 8, 5)$ $\text{Diff} = \emptyset$	3
69	$z^3x^2 + x^3y^2 + az^4y^{n-2} +$ $bzxxy^n, n = 5; 7$	$t + g(z, x, y) \subset \mathbb{P}(9n - 2, n + 2, 3n - 4, 5)$ $\mathbb{P}(n + 2, 3n - 4, 5), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	8,10
70	$z^3x^2 + x^3y^2 + az^4y^4 +$ $bzxxy^6$	$t^2 + g(z, x, y^{1/2}) \subset \mathbb{P}(13, 4, 7, 5)$ $\text{Diff} = (0, 0, 0, \frac{1}{2})$	5
71	$z^3x^2 + x^3y^2 + z^2y^{4n}$ $n = 1; 2$	$t^2 + g(z, x, y^{1/2}) \subset \mathbb{P}(9n + 2, 2n + 2, 6n - 1, 7)$ $\text{Diff} = (0, 0, 0, \frac{1}{2})$	6,7
72	$z^3x^2 + x^3y^2 + z^2y^n$ $n = 5; 7; 9$	$t + g(z, x, y) \subset \mathbb{P}(9n + 8, n + 4, 3n - 2, 7)$ $\mathbb{P}(n + 4, 3n - 2, 7), \text{Diff} = (\frac{1}{2}, 0, 0, 0)$	8,12,14
73	$z^3x^2 + x^3y^2 + z^2y^6$	$t^2 + g(z, x, y) \subset \mathbb{P}(31, 10, 16, 7)$ $\text{Diff} = \emptyset$	5

(1) The exceptionality condition is $|c| + |d| + |l| + |n| + |e(|a| + |b|)| \neq 0$.

13. Singularity $-t^3 + g(t, z, x, y)$

1	$z^2x^2 + x^4y^j f_{5-i-j}(x, y)$ $2i + j \leq 3$	$t + zx^2 + x^4y^j f_{5-i-j}(x, y) \subset \mathbb{P}(5, 3, 1, 1)$ $\mathbb{P}(3, 1, 1), \text{Diff} = (\frac{2}{3}, \frac{1}{2}, 0, 0)$	6
2	$z^2x^2 + x^5 + zy^4$	$t + z^2x^2 + x^5 + zy \subset \mathbb{P}(10, 3, 2, 7)$	21

		$\mathbb{P}(3, 2, 7), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{3}{4})$	
3	$z^2x^2 + x^4y + zy^4$	$t + z^2x + x^2y + zy^4 \subset \mathbb{P}(17, 5, 7, 3)$ $\mathbb{P}(5, 7, 3), \text{Diff}=(\frac{2}{3}, 0, \frac{1}{2}, 0)$	18
4	$z^2x^2 + x^3y^2 + zy^4$	$t + z^2x^2 + x^3y + zy^2 \subset \mathbb{P}(14, 4, 3, 5)$ $\mathbb{P}(4, 3, 5), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{1}{2})$	15
5	$z^2xy + x^iy^jf_{5-i-j}(x, y)$ $i, j \leq 1$	$t + zx^2 + x^iy^jf_{5-i-j}(x, y) \subset \mathbb{P}(5, 3, 1, 1)$ $\mathbb{P}(3, 1, 1), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, 0)$	6
6	$z^2xy + x^5 + aty^4 + by^6$	$t^3 + zxy + aty^4 + by^6 \subset \mathbb{P}(10, 19, 6, 5)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	10
7	$z^2xy + x^5 + y^7$	$t + zxy + x^5 + y^7 \subset \mathbb{P}(35, 23, 7, 5)$ $\mathbb{P}(23, 7, 5), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, 0)$	30
8	$z^2xy + x^5 + x^2y^4$	$t + zxy + x^5 + x^2y^4 \subset \mathbb{P}(20, 13, 4, 3)$ $\mathbb{P}(13, 4, 3), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, 0)$	18
9	$z^2xy + x^5 + zy^4$	$t + g(z, x, y) \subset \mathbb{P}(35, 11, 7, 6)$ $\mathbb{P}(11, 7, 6), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	9
10	$z^2xy + x^5 + txy^3$	$t^3 + zxy + x^5 + txy^3 \subset \mathbb{P}(15, 29, 9, 7)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	14
11	$z^ix^jf_{4-i-j}(z, x) + y^5$ $i = j = 1$	$t + z^ix^jf_{4-i-j}(z, x) + y \subset \mathbb{P}(4, 1, 1, 4)$ $\mathbb{P}(4, 1, 1), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{4}{5})$	6
12	$z^ix^jf_{4-i-j}(z, x) + xy^4$ $i \leq 2 \text{ \& } j \leq 1$	$t + z^ix^jf_{4-i-j}(z, x) + xy \subset \mathbb{P}(4, 1, 1, 3)$ $\mathbb{P}(1, 1, 3), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{3}{4})$	9
13	$z^ix^jf_{4-i-j}(z, x) + ty^3$ $i, j \leq 1$	$t^3 + g(z, x, y^{1/3}) \subset \mathbb{P}(4, 3, 3, 8)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	4
14	$z^ix^jf_{4-i-j}(z, x) + txy^2$ $i \leq 2 \text{ \& } j \leq 1$	$t^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(4, 3, 3, 5)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	5
15	$z^4 + zx^2y + y^n$ $n = 5; 7; 11$	$t + z^4 + zxy + y^n \subset \mathbb{P}(4n, n, 3n - 4, 4)$ $\mathbb{P}(n, 3n - 4, 4), \text{Diff}=(\frac{2}{3}, 0, \frac{1}{2}, 0)$	9,15,24
16	$z^4 + zx^2y + aty^4 + btz^2y +$ $cz^2y^3 + dy^6, a + d \neq 0$	$t^3 + g(z, x^{1/2}, y) \subset \mathbb{P}(4, 3, 7, 2)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	4
17	$z^4 + zx^2y + ay^4z^if_{2-i}(z, y^2)$	$t + g(z, x^{1/2}, y) \subset \mathbb{P}(8, 2, 5, 1)$ $\mathbb{P}(2, 5, 1), \text{Diff}=(\frac{2}{3}, 0, \frac{1}{2}, 0)$	6
18	$z^4 + zx^2y + aty^6 + by^9$	$t^3 + g(z, x^{1/2}, y) \subset \mathbb{P}(12, 9, 23, 4)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	8
19	$z^4 + zx^2y + az^2y^5 + by^{10}$	$t + g(z, x^{1/2}, y) \subset \mathbb{P}(20, 5, 13, 2)$ $\mathbb{P}(5, 13, 2), \text{Diff}=(\frac{2}{3}, 0, \frac{1}{2}, 0)$	12
20	$z^4 + zx^2y + xy^n$ $n = 4; 6; 7$	$t + g(z, x, y) \subset \mathbb{P}(8n - 4, 2n - 1, 3n - 4, 5)$ $\mathbb{P}(2n - 1, 3n - 4, 5), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	6,15
21	$z^4 + zx^2y + atzy^3 + bxy^5$	$t^3 + g(z, x, y) \subset \mathbb{P}(12, 9, 11, 5)$ $\text{Diff} = \emptyset$	5
22	$z^4 + zx^2y + ty^n$ $n = 3; 5; 7$	$t^3 + g(z, x^{1/2}, y) \subset \mathbb{P}(4n, 3n, 9n - 8, 8)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	7,10,16
23	$z^4 + zx^2y + txy^n$ $n = 2; 3$	$t^3 + g(z, x, y) \subset \mathbb{P}(8n - 4, 6n - 3, 9n - 8, 7)$ $\text{Diff} = \emptyset$	5,7
24	$z^4 + zx^2y + tzy^4$	$t^3 + z^4 + zxy + tzy^4 \subset \mathbb{P}(16, 12, 31, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	10
25	$z^4 + x^3y + atxy^2 + by^5$	$t^3 + z + x^3y + atxy^2 + by^5 \subset \mathbb{P}(5, 15, 4, 3)$ $\mathbb{P}(5, 4, 3), \text{Diff}=(0, \frac{3}{4}, 0, 0)$	4
26	$z^4 + x^3y + atz^2y + bty^4 +$	$t^3 + g(z^{1/2}, x^{1/3}, y) \subset \mathbb{P}(2, 3, 5, 1)$	6

	$cz^2y^3 + dy^6, b + c + d \neq 0$	$\text{Diff}=(0, \frac{1}{2}, \frac{2}{3}, 0)$	
27	$z^4 + x^3y + ay^3x^i f_{2-i}(x, y^2)$	$t + g(z^{1/4}, x, y) \subset \mathbb{P}(7, 7, 2, 1)$ $\mathbb{P}(7, 2, 1), \text{Diff}=(\frac{2}{3}, \frac{3}{4}, 0, 0)$	12
28	$z^4 + x^3y + xy^4$	$t + z + x^3y + xy^4 \subset \mathbb{P}(11, 11, 3, 2)$ $\mathbb{P}(11, 3, 2), \text{Diff}=(\frac{2}{3}, \frac{3}{4}, 0, 0)$	12
29	$z^4 + x^3y + zy^n$ $n = 4; 5$	$t^3 + g(z, x, y) \subset \mathbb{P}(4n, 3n, 4n - 3, 9)$ $\text{Diff} = \emptyset$	6,9
30	$z^4 + x^3y + zxy^3$	$t + z^4 + x^3y + zxy^3 \subset \mathbb{P}(32, 8, 9, 5)$ $\mathbb{P}(8, 9, 5), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	12
31	$z^4 + x^3y + ty^n$ $n = 3; 5$	$t^3 + z + xy + ty^n \subset \mathbb{P}(n, 3n, 3n - 2, 2)$ $\mathbb{P}(n, 3n - 2, 2), \text{Diff}=(0, \frac{3}{4}, \frac{2}{3}, 0)$	4,24
32	$z^4 + x^3y + tzy^3$	$t^3 + z^4 + xy + tzy^3 \subset \mathbb{P}(12, 9, 31, 5)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	15
33	$z^4 + tx^3 + y^5$	$t^3 + z + tx + y \subset \mathbb{P}(1, 3, 2, 3)$ $\mathbb{P}(1, 2, 3), \text{Diff}=(0, \frac{3}{4}, \frac{2}{3}, \frac{4}{5})$	36
34	$z^4 + tx^3 + tzy^2$	$t^3 + z^4 + tx + tzy \subset \mathbb{P}(4, 3, 8, 5)$ $\text{Diff}=(0, 0, \frac{2}{3}, \frac{1}{2})$	15
35	$z^4 + tx^2y + y^5$	$t^3 + z^2 + tx^2y + y^5 \subset \mathbb{P}(10, 15, 7, 6)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	12
36	$z^4 + tx^2y + xy^4$	$t^3 + z + tx^2y + xy^4 \subset \mathbb{P}(7, 21, 5, 4)$ $\mathbb{P}(7, 5, 4), \text{Diff}=(0, \frac{3}{4}, 0, 0)$	16
37	$z^4 + tx^2y + zy^4$	$t^3 + z^4 + txy + zy^4 \subset \mathbb{P}(16, 12, 23, 9)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	18
38	$z^4 + tx^2y + tzy^2$	$t^3 + z^4 + txy + tzy^2 \subset \mathbb{P}(8, 6, 11, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	10
39	$z^3x + zx^2y + y^n$ $n = 5; 7; 8$	$t + g(z, x, y) \subset \mathbb{P}(5n, n + 1, 2n - 3, 5)$ $\mathbb{P}(n + 1, 2n - 3, 5), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	9,12,15
40	$z^3x + zx^2y + aty^4 + by^6$	$t^3 + g(z, x, y) \subset \mathbb{P}(10, 7, 9, 5)$ $\text{Diff} = \emptyset$	5
41	$z^3x + zx^2y + az^2y^4 + bxy^5$	$t + g(z, x, y) \subset \mathbb{P}(22, 5, 7, 3)$ $\mathbb{P}(5, 7, 3), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	9
42	$z^3x + zx^2y + tzy^3$	$t^3 + g(z, x, y) \subset \mathbb{P}(16, 11, 15, 7)$ $\text{Diff} = \emptyset$	7
43	$z^3x + x^3y + atxy^2 + y^5$	$t^3 + g(z^{1/3}, x, y) \subset \mathbb{P}(5, 11, 4, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	6
44	$z^3x + x^3y + aty^4 + by^6$	$t^3 + g(z^{1/3}, x, y) \subset \mathbb{P}(6, 13, 5, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
45	$z^3x + x^3y + zy^n$ $n = 4; 5$	$t + g(z, x, y) \subset \mathbb{P}(9n + 1, 2n + 1, 3n - 2, 7)$ $\mathbb{P}(2n + 1, 3n - 2, 7), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	16,21
46	$z^3x + x^3y + z^2y^3$	$t + g(z, x, y) \subset \mathbb{P}(29, 7, 8, 5)$ $\mathbb{P}(7, 8, 5), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	12
47	$z^3x + tx^3 + y^5$	$t^3 + zx + tx^3 + y \subset \mathbb{P}(3, 7, 2, 9)$ $\mathbb{P}(3, 7, 2), \text{Diff}=(0, \frac{2}{3}, 0, \frac{4}{5})$	15
48	$z^3x + tx^3 + tzy^2$	$t^3 + z^3x + tx^3 + tzy \subset \mathbb{P}(9, 7, 6, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
49	$z^3x + tx^2y + y^5$	$t^3 + zx + tx^2y + y^5 \subset \mathbb{P}(10, 23, 7, 6)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
50	$z^3y + az^2x^2 + btx^2y +$	$t^3 + g(z, x^{1/2}, y) \subset \mathbb{P}(5, 4, 7, 3)$	6

	$ctzy^2 + dy^5, ab \neq 0 \&$ $(d \neq 0 \parallel c \neq 0)$	$\text{Diff}=(0, 0, \frac{1}{2}, 0)$	
51	$z^3y + z^2x^2 + xy^4$	$t + z^3y + z^2x^2 + xy^4 \subset \mathbb{P}(26, 7, 6, 5)$ $\mathbb{P}(7, 6, 5), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	9
52	$z^3y + z^2x^2 + azy^4 + bx^2y^3$	$t + z^3y + z^2x + azy^4 + bxy^3 \subset \mathbb{P}(11, 3, 5, 2)$ $\mathbb{P}(3, 5, 2), \text{Diff}=(\frac{2}{3}, 0, \frac{1}{2}, 0)$	12
53	$z^3y + z^2x^2 + txy^2$	$t^3 + g(z, x, y) \subset \mathbb{P}(14, 11, 10, 9)$ $\text{Diff} = \emptyset$	5
54	$z^3y + tx^3 + xy^4$	$t^3 + zy + tx^3 + xy^4 \subset \mathbb{P}(12, 29, 8, 7)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
55	$tz^3 + z^2x^2 + xy^4$	$t^3 + tz^3 + z^2x^2 + xy \subset \mathbb{P}(6, 4, 5, 13)$ $\text{Diff}=(0, 0, 0, \frac{3}{4})$	13
56	$atz^3 + z^2x^2 + txy^2 +$ $bzy^4, a + b \neq 0$	$t^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(6, 4, 5, 7)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	7
57	$tz^3 + zx^2y + y^{2n+1}$ $n = 2; 3$	$t^3 + g(z, x, y) \subset \mathbb{P}(6n + 3, 4n + 2, 7n - 1, 9)$ $\text{Diff} = \emptyset$	5,9
58	$tz^3 + zx^2y + aty^4 +$ $bz^3y^2 + cy^6, a + c \neq 0$	$t^3 + g(z, x^{1/2}, y) \subset \mathbb{P}(6, 4, 11, 3)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	6
59	$tz^3 + zx^2y + y^8$	$t^3 + tz^3 + zxy + y^8 \subset \mathbb{P}(24, 16, 47, 9)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	18
60	$tz^3 + zx^2y + xy^4$	$t^3 + g(z, x, y) \subset \mathbb{P}(21, 14, 19, 11)$ $\text{Diff} = \emptyset$	7
61	$tz^3 + zx^2y + z^2y^4$	$t^3 + tz^3 + zxy + z^2y^4 \subset \mathbb{P}(12, 8, 23, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	10
61	$z^2x^2 + tx^2y + zy^4$	$t^3 + z^2x + txy + zy^4 \subset \mathbb{P}(9, 7, 13, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	10
62	$tz^2x + x^2y^2 + y^5$	$t^3 + tzx + x^2y^2 + y^5 \subset \mathbb{P}(10, 11, 9, 6)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	11
63	$tz^2x + x^2y^2 + zy^4$	$t^3 + tz^2x + x^2y + zy^2 \subset \mathbb{P}(9, 5, 8, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	10
64	$tz^2x + x^2y^2 + z^2y^3$	$t^3 + tzx + x^2y^2 + zy^3 \subset \mathbb{P}(8, 9, 7, 5)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	9
65	$z^2x + x^4 + y^n$ $n = 7; 11; 13; 17; 19, 23$	$t + zx + x^4 + y \subset \mathbb{P}(4, 3, 1, 4)$ $\mathbb{P}(4, 3, 1), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, \frac{n-1}{n})$	7,12,15 18,24
66	$z^2x + x^4 + ty^n(cx^2 + dy^{3n}) +$ $ax^2y^{3n} + by^{6n}, n = 1 \& b + d \neq 0; 3$	$t^3 + g(z^{1/2}, x, y^{1/n}) \subset \mathbb{P}(4, 9, 3, 2)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{n-1}{n})$	3,4
67	$z^2x + x^i(x + y^2)^j f_{4-i}(x, y^{2n}) + azy^{5n}$ $n = 1 \& (a \neq 0 \parallel (i = 0 \& j \leq 3)), 2$	$t + g(z, x, y^{1/n}) \subset \mathbb{P}(8, 3, 2, 1)$ $\mathbb{P}(3, 2, 1), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{n-1}{n})$	3
68	$z^2x + x^4 + aty^{2n} + by^{3n}$ $n = 3; 5, 7$	$t^3 + zx + x^4 + aty^2 + by^3 \subset \mathbb{P}(4, 9, 3, 4)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{n-1}{n})$	7,8
69	$z^2x + x^4 + ax^2y^n + by^{2n}$ $n = 5 \& b \neq 0; 7; 11$	$t + zx + x^4 + ax^2y + by^2 \subset \mathbb{P}(4, 3, 1, 2)$ $\mathbb{P}(3, 1, 2), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, \frac{n-1}{n})$	6,12
70	$z^2x + aty^2x^i f_{2-i}(x, y^3) +$ $x^k f_{4-k}(x, y^3), i \leq 1 \parallel k \leq 1$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(4, 9, 3, 1)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	2
71	$z^2x + x^4 + atxy^{2n} + bzy^{3n}$ $(n = 1; 2 \& b \neq 0); 3, 4$	$t^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(8, 9, 6, 5)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	2,4,5
72	$z^2x + x^4 + zy^n$ $n = 4; 7; 8; 11, 13, 14$	$t + z^2x + x^4 + zy \subset \mathbb{P}(8, 3, 2, 5)$ $\mathbb{P}(3, 2, 5), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{n-1}{n})$	6,7,9 15

73	$z^2x + x^4 + ty^n$ $n = 5, 7; 9; 11, 13, 15$	$t^3 + zx + x^4 + ty \subset \mathbb{P}(4, 9, 3, 8)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{n-1}{n})$	7,10,16
74	$z^2x + x^4 + txy^n$ $n = 7, 9$	$t^3 + zx + x^4 + txy \subset \mathbb{P}(4, 9, 3, 5)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{n-1}{n})$	10
75	$z^2x + x^4 + tzy^n$ $n = 2, 3; 4, 5, 6$	$t^3 + z^2x + x^4 + tzy \subset \mathbb{P}(8, 9, 6, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	6,7
76	$z^2x + x^i(x + y^n)^j y^{nk} f_{5-i-j-k}(x, y^n) +$ $(0) \quad azy^{3n}, n = 1, 2$	$t + g(z, x, y^{1/n}) \subset \mathbb{P}(5, 2, 1, 1)$ $\mathbb{P}(2, 1, 1), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{n-1}{n})$	3
77	$z^2x + x^5 + aty^{2n} + by^{3n}$ $n = 3, 4$	$t^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(5, 6, 3, 5)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5
78	$z^2x + x^5 + y^n$ $n = 7; 8; 11; 13, 14$	$t + z^2x + x^5 + y \subset \mathbb{P}(5, 2, 1, 5)$ $\mathbb{P}(2, 1, 5), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{n-1}{n})$	7,9,12 15
79	$z^2x + x^5 + azy^n + bx^2y^n$ $(n = 4; 5) \& a \neq 0; 7; 8$	$t + z^2x + x^5 + azy + bx^2y \subset \mathbb{P}(5, 2, 1, 3)$ $\mathbb{P}(2, 1, 3), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{n-1}{n})$	4,6,9
80	$z^2x + x^5 + txy^n$ $n = 5, 6$	$t^3 + z^2x + x^5 + txy \subset \mathbb{P}(5, 6, 3, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	7
81	$z^2x + x^4y + aty^{2n} + by^{3n}$ $n = 4, 6$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(4n, 9n + 1, 3n - 1, 4)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	8
82	$z^2x + x^4y + ax^2y^n + by^{2n-1}$ $(n = 4; 6) \& b \neq 0; 10$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(4n - 2, 3n - 1, n - 1, 2)$ $\mathbb{P}(3n - 1, n - 1, 2), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, 0)$	4,6,12
83	$z^2x + x^4y + y^n$ $n = 8; 10; 14; 16; 20$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(4n, 3n + 1, n - 1, 4)$ $\mathbb{P}(3n + 1, n - 1, 4), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, 0)$	9,12,16 18,24
84	$z^2x + tf_3(x, y^2) + yf_4(x, y^2)$ (1)	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(3, 7, 2, 1)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	2
84'	$z^2x + azy^8 + x^i y f_{4-i}(x, y^3)$ $a \neq 0 i \leq 2$	$t + g(z, x, y) \subset \mathbb{P}(13, 5, 3, 1)$ $\mathbb{P}(5, 3, 1), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	3
85	$z^2x + ty^3(ax^2 + by^7) + x^{2i} y f_{2-i}(x^2, y^7)$ $b \neq 0 i \leq 1$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(10, 23, 7, 2)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	4
86	$z^2x + x^i y f_{4-i}(x, y^4)$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(17, 13, 4, 1)$ $\mathbb{P}(13, 4, 1), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, 0)$	6
87	$z^2x + x^4y + atxy^{2n+1} + bzy^{3n+1}$ $n = 1 \ b \neq 0; 2, 3$	$t^3 + g(z, x, y) \subset \mathbb{P}(8n + 3, 9n + 4, 6n + 1, 5)$ $\text{Diff} = \emptyset$	4,5
88	$z^2x + x^4y + zy^n$ $n = 5; 6; 9, 11, 12$	$t + g(z, x, y) \subset \mathbb{P}(8n + 1, 3n + 1, 2n - 1, 5)$ $\mathbb{P}(3n + 1, 2n - 1, 5), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	6,7,15
89	$z^2x + x^4y + ty^n$ $n = 5; 7; 9; 11, 13$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(4n, 9n + 2, 3n - 2, 8)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	7,8,10 16
90	$z^2x + x^4y + txy^n$ $n = 6, 8$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(4n - 1, 9n - 1, 3n - 2, 5)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	10
91	$z^2x + x^4y + tzy^n$ $n = 2; 3, 4, 5$	$t^3 + g(z, x, y) \subset \mathbb{P}(8n + 1, 9n + 2, 6n - 1, 7)$ $\text{Diff} = \emptyset$	5,7
92	$z^2x + \underline{a}tx^i y^j f_{4-i-j}(x, y) + \underline{b}x^k y^l \cdot$ $\cdot f_{6-k-l}(x, y), i = 0 k = 0$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(2, 5, 1, 1)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	2
93	$z^2x + ax^6 + btx^4 + y^n$ $n = 7; 11$	$t^3 + zx + ax^6 + btx^4 + y \subset \mathbb{P}(2, 5, 1, 6)$ $\mathbb{P}(2, 5, 1), \text{Diff}=(0, \frac{1}{2}, 0, \frac{n-1}{n})$	7,12
94	$z^2x + tx(\underline{a}x^3 + by^4) + ctzy^2 + \underline{d}x^{3i} \cdot$ $\cdot f_{2-i}(x^3, y^4), c \neq 0 i = 0$	$t^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(4, 5, 2, 3)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	3
95	$z^2x + t(\underline{a}x^4 + bx^2y^3 + cy^6) + \underline{d}x^{2i} y^{3j} \cdot$ $\cdot f_{3-i-j}(x^2, y^3), j \leq 1 \& (c \neq 0 i = 0)$	$t^3 + g(z^{1/2}, x, y^{1/3}) \subset \mathbb{P}(2, 5, 1, 2)$ $\text{Diff}=(0, \frac{1}{2}, 0, \frac{2}{3})$	4

96	$z^2x + tx(\underline{a}x^3 + by^5) + \underline{c}x^{3i}f_{2-i}(x^3, y^5)$ $b \neq 0 i = 0$	$t^3 + g(z^{1/2}, x, y^{1/5}) \subset \mathbb{P}(2, 5, 1, 3)$ Diff= $(0, \frac{1}{2}, 0, \frac{4}{5})$	6
97	$z^2x + \underline{a}tx^4 + cty^n + \underline{b}x^6 + dx^2y^n$ $n = 5 \& c \neq 0; n = 7$	$t^3 + g(z^{1/2}, x, y^{1/n}) \subset \mathbb{P}(2, 5, 1, 4)$ Diff= $(0, \frac{1}{2}, 0, \frac{n-1}{n})$	6,8
98	$z^2x + ax^6 + btx^4 + zy^n$ $n = 4, 5; 6$	$t^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(4, 5, 2, 7)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	5,7
99	$z^2x + x^5y + y^n$ $n = 7, 8; 10; 12$	$t + zx + x^5y + y^n \subset \mathbb{P}(5n, 4n + 1, n - 1, 5)$ $\mathbb{P}(4n + 1, n - 1, 5)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	9,12,30
100	$z^2x + x^5y + aty^{2n} + by^{3n}$ $n = 3, 4$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(5n, 12n + 1, 3n - 1, 5)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10
101	$z^2x + x^i y^{2j+1} f_{5-i-j}(x, y^2)$ $i \leq 2 \& j \leq 1$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(11, 9, 2, 1)$ $\mathbb{P}(9, 2, 1)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	6
102	$z^2x + zy^2(ax^2 + by^3) + x^{2i+1}y \cdot$ $f_{2-i}(x^2, y^3), b \neq 0 i = 0$	$t + g(z, x, y) \subset \mathbb{P}(17, 7, 3, 2)$ $\mathbb{P}(7, 3, 2)$, Diff= $(\frac{2}{3}, 0, 0, 0)$	3
103	$z^2x + x^5y + x^2y^8$	$t + zx + x^5y + x^2y^8 \subset \mathbb{P}(38, 31, 7, 3)$ $\mathbb{P}(31, 7, 3)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	18
104	$z^2x + x^5y + zy^n$ $n = 4, 6, 7$	$t + g(z, x, y) \subset \mathbb{P}(10n + 1, 4n + 1, 2n - 1, 6)$ $\mathbb{P}(4n + 1, 2n - 1, 6)$, Diff= $(\frac{2}{3}, 0, 0, 0)$	9
105	$z^2x + x^5y + ty^n$ $n = 5, 7$	$t^3 + g(z, x, y) \subset \mathbb{P}(5n, 6n + 1, 3n - 2, 10)$ Diff = $\emptyset$	5
106	$z^2x + x^5y + txy^n$ $n = 5, 6$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(5n - 1, 12n - 1, 3n - 2, 7)$ Diff= $(0, \frac{1}{2}, 0, 0)$	14
107	$z^2x + x^4y^2 + y^n$ $n = 7; 11; 13; 17$	$t + zx + x^4y^2 + y^n \subset \mathbb{P}(4n, 3n + 2, n - 2, 4)$ $\mathbb{P}(3n + 2, n - 2, 4)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	9,15,18 24
108	$z^2x + x^{2i}y^2f_{2-i}(x^2, y^n)$ $n = 3 \ i = 0; 7$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(4n + 4, 3n + 4, n, 2)$ $\mathbb{P}(3n + 4, n, 2)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	6,12
109	$z^2x + x^4y^2 + aty^{2n} + by^{3n}$ $n = 3, 5$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(4n, 9n + 2, 3n - 2, 4)$ Diff= $(0, \frac{1}{2}, 0, 0)$	8
110	$z^2x + ty^3(ax^2 + by^5) + x^{2i}y^2f_{2-i}(x^2, y^5)$ $b \neq 0 i \leq 1$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(8, 19, 5, 2)$ Diff= $(0, \frac{1}{2}, 0, 0)$	4
111	$z^2x + x^i y^2 f_{4-i}(x, y^3)$ $i \leq 2$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(14, 11, 3, 1)$ $\mathbb{P}(3n + 4, n, 2)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	6
112	$z^2x + x^4y^2 + zy^{2n}$ $n = 2; 5$	$t + g(z, x, y^{1/2}) \subset \mathbb{P}(8n + 1, 3n + 1, 2n - 1, 5)$ $\mathbb{P}(3n + 1, 2n - 1, 5)$, Diff= $(\frac{2}{3}, 0, 0, \frac{1}{2})$	6,10
113	$z^2x + x^4y^2 + atxy^4 + zy^5$	$t^3 + g(z, x, y) \subset \mathbb{P}(14, 17, 8, 5)$ Diff = $\emptyset$	4
114	$z^2x + x^4y^2 + zy^n$ $n = 7; 9$	$t + g(z, x, y) \subset \mathbb{P}(8n + 2, 3n + 2, 2n - 2, 5)$ $\mathbb{P}(3n + 2, 2n - 2, 5)$, Diff= $(\frac{2}{3}, 0, 0, 0)$	9,15
115	$z^2x + x^4y^2 + atxy^6 + bzy^8$	$t^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(11, 13, 7, 5)$ Diff= $(0, 0, 0, \frac{1}{2})$	5
116	$z^2x + x^4y^2 + ty^n$ $n = 5; 7; 9, 11$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(4n, 9n + 4, 3n - 4, 8)$ Diff= $(0, \frac{1}{2}, 0, 0)$	7,10,16
117	$z^2x + x^4y^2 + txy^n$ $n = 5, 7$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(4n - 2, 9n - 2, 3n - 4, 5)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10
118	$z^2x + x^4y^2 + tzy^{2n}$ $n = 1; 2$	$t^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(8n + 1, 9n + 2, 6n - 1, 7)$ Diff= $(0, 0, 0, \frac{1}{2})$	6,7
119	$z^2x + x^4y^2 + tzy^3$	$t^3 + g(z, x, y) \subset \mathbb{P}(26, 31, 16, 7)$ Diff = $\emptyset$	8

120	$z^2x + x^iy^j(x+y)^k f_{7-i-j-k}(x,y) + azy^4$ $j \leq 3 \& k \leq 3 \& (a \neq 0 i = 0)$	$t + g(z, x, y) \subset \mathbb{P}(7, 3, 1, 1)$ $\mathbb{P}(3, 1, 1), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	3
121	$z^2x + x^7 + y^n$ $n = 8; 10$	$t + z^2x + x^7 + y \subset \mathbb{P}(7, 3, 1, 7)$ $\mathbb{P}(7, 3, 1), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{n-1}{n})$	9,21
122	$z^2x + x^7 + aty^6 + by^9$	$t^3 + z^2x + x^7 + aty^2 + by^3 \subset \mathbb{P}(7, 9, 3, 7)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	7
123	$z^2x + ty^2(z + \underline{a}x^3) + \underline{b}x^7 + cx^2y^6$	$t^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(7, 9, 3, 5)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	6
124	$z^2x + x^7 + x^2y^7$	$t + z^2x + x^7 + x^2y \subset \mathbb{P}(7, 3, 1, 5)$ $\mathbb{P}(3, 1, 5), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{6}{7})$	15
125	$z^2x + x^7 + zy^5 + bx^3y^5$	$t + z^2x + x^7 + zy + bx^3y \subset \mathbb{P}(7, 3, 1, 4)$ $\mathbb{P}(3, 1, 4), \text{Diff}=(\frac{2}{3}, 0, 0, \frac{4}{5})$	6
126	$z^2x + x^7 + ty^5$	$t^3 + z^2x + x^7 + ty \subset \mathbb{P}(7, 9, 3, 14)$ $\text{Diff}=(0, 0, 0, \frac{4}{5})$	7
127	$z^2x + x^7 + txy^5$	$t^3 + z^2x + x^7 + txy \subset \mathbb{P}(7, 9, 3, 11)$ $\text{Diff}=(0, 0, 0, \frac{4}{5})$	11
128	$z^2x + x^6y + atx^2y^3 + y^8$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(16, 41, 7, 6)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	4
129	$z^2x + ty^2(\underline{a}x^3 + by^4) + \underline{c}x^{3i}y \cdot$ $f_{2-i}(x^3, y^4), b \neq 0 i = 0$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(9, 23, 4, 3)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	6
130	$z^2x + x^{2i}y^{3j+1}f_{3-i-j}(x^2, y^3)$ $i \leq 1 \& j \leq 1$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(20, 17, 3, 2)$ $\mathbb{P}(17, 3, 2), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, 0)$	12
131	$z^2x + x^6y + zy^n$ $n = 5; 6$	$t + g(z, x, y) \subset \mathbb{P}(12n + 1, 5n + 1, 2n - 1, 7)$ $\mathbb{P}(5n + 1, 2n - 1, 7), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	12,21
132	$z^2x + x^6y + ty^n$ $n = 5; 7$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(6n, 15n + 2, 3n - 2, 12)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	11,24
133	$z^2x + x^6y + txy^5$	$t^3 + g(z, x, y) \subset \mathbb{P}(29, 37, 13, 9)$ $\text{Diff} = \emptyset$	9
134	$z^2x + x^6y + tzy^2$	$t^3 + g(z, x, y) \subset \mathbb{P}(25, 32, 11, 9)$ $\text{Diff} = \emptyset$	9
135	$z^2x + x^5y^2 + y^n$ $n = 8; 10; 11$	$t + g(z, x, y) \subset \mathbb{P}(5n, 2n + 1, n - 2, 5)$ $\mathbb{P}(2n + 1, n - 2, 5), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	9,12,15
136	$z^2x + x^5y^2 + aty^6 + by^9$	$t^3 + g(z, x, y) \subset \mathbb{P}(15, 19, 7, 5)$ $\text{Diff} = \emptyset$	5
137	$z^2x + x^5y^2 + azy^{n-1} + bx^2y^n$ $n = 6 \ a \neq 0; 7$	$t + g(z, x, y) \subset \mathbb{P}(5n - 4, 2n - 1, n - 2, 3)$ $\mathbb{P}(2n - 1, n - 2, 3), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	6,9
138	$z^2x + x^5y^2 + txy^5$	$t^3 + g(z, x, y) \subset \mathbb{P}(23, 29, 11, 7)$ $\text{Diff} = \emptyset$	7
139	$z^2x + x^4y^3 + y^n$ $n = 8; 10; 14$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(4n, 3n + 3, n - 3, 4)$ $\mathbb{P}(3n + 3, n - 3, 4), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, 0)$	12,16,24
140	$z^2x + x^4y^3 + aty^8 + by^{12}$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(16, 39, 9, 4)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	8
141	$z^2x + x^4y^3 + ax^2y^8 + by^{13}$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(26, 21, 5, 2)$ $\mathbb{P}(21, 5, 2), \text{Diff}=(\frac{2}{3}, \frac{1}{2}, 0, 0)$	12
142	$z^2x + x^4y^3 + zy^n$ $n = 5; 7, 8$	$t + g(z, x, y) \subset \mathbb{P}(8n + 3, 3n + 3, 2n - 3, 5)$ $\mathbb{P}(3n + 3, 2n - 3, 5), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	9,15
143	$z^2x + x^4y^3 + atxy^5 + bzy^6$	$t^3 + g(z, x, y) \subset \mathbb{P}(17, 21, 9, 5)$ $\text{Diff} = \emptyset$	5

144	$z^2x + x^4y^3 + ty^n$ $n = 5; 7$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(4n, 9n + 6, 3n - 6, 8)$ Diff = $(0, \frac{1}{2}, 0, 0)$	8,10
145	$z^2x + x^4y^3 + ty^9$	$t^3 + zx + x^4y + ty^3 \subset \mathbb{P}(12, 29, 7, 8)$ Diff = $(0, \frac{1}{2}, 0, \frac{2}{3})$	16
146	$z^2x + x^4y^3 + txy^6$	$t^3 + zx + x^4y + txy^2 \subset \mathbb{P}(7, 17, 4, 5)$ Diff = $(0, \frac{1}{2}, 0, \frac{2}{3})$	10
147	$z^2x + x^4y^3 + tzy^2$	$t^3 + g(z, x, y) \subset \mathbb{P}(19, 24, 9, 7)$ Diff = $\emptyset$	7
148	$z^2x + x^4y^3 + tzy^3$	$t^3 + z^2x + x^4y + tzy \subset \mathbb{P}(9, 11, 5, 7)$ Diff = $(0, 0, 0, \frac{2}{3})$	7
149	$z^2x + x^i y^j f_{8-i-j}(x, y)$ $i \leq 1 \& j \leq 4$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(8, 7, 1, 1)$ $\mathbb{P}(7, 1, 1)$, Diff = $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	6
150	$z^2x + x^8 + aty^6 + by^9$	$t^3 + zx + x^8 + aty^2 + by^3 \subset \mathbb{P}(8, 21, 3, 8)$ Diff = $(0, \frac{1}{2}, 0, \frac{2}{3})$	16
151	$z^2x + x^8 + zy^5$	$t + z^2x + x^8 + zy \subset \mathbb{P}(16, 7, 2, 9)$ $\mathbb{P}(7, 2, 9)$, Diff = $(\frac{2}{3}, 0, 0, \frac{4}{5})$	15
152	$z^2x + x^8 + x^2y^7$	$t + zx + x^8 + x^2y \subset \mathbb{P}(8, 7, 1, 6)$ $\mathbb{P}(7, 1, 6)$, Diff = $(\frac{2}{3}, \frac{1}{2}, 0, \frac{6}{7})$	36
153	$z^2x + x^8 + txy^5$	$t^3 + zx + x^8 + txy \subset \mathbb{P}(8, 21, 3, 13)$ Diff = $(0, \frac{1}{2}, 0, \frac{4}{5})$	26
154	$z^2x + x^8 + tzy^2$	$t^3 + z^2x + x^8 + tzy \subset \mathbb{P}(16, 21, 6, 11)$ Diff = $(0, 0, 0, \frac{1}{2})$	11
155	$z^2x + x^7y + aty^6 + by^9$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(21, 55, 8, 7)$ Diff = $(0, \frac{1}{2}, 0, 0)$	14
156	$z^2x + x^7y + x^2y^7$	$t + zx + x^7y + x^2y^7 \subset \mathbb{P}(47, 41, 6, 5)$ $\mathbb{P}(41, 6, 5)$, Diff = $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	36
157	$z^2x + x^7y + zy^5$	$t + g(z, x, y) \subset \mathbb{P}(71, 31, 9, 8)$ $\mathbb{P}(31, 9, 8)$, Diff = $(\frac{2}{3}, 0, 0, 0)$	12
158	$z^2x + x^7y + txy^5$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(34, 89, 13, 11)$ Diff = $(0, \frac{1}{2}, 0, 0)$	22
159	$z^2x + x^6y^2 + aty^6 + by^9$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(18, 47, 7, 6)$ Diff = $(0, \frac{1}{2}, 0, 0)$	12
160	$z^2x + x^6y^2 + x^2y^7$	$t + zx + x^6y^2 + x^2y^7 \subset \mathbb{P}(38, 33, 5, 4)$ $\mathbb{P}(33, 5, 4)$, Diff = $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	24
161	$z^2x + x^6y^2 + zy^5$	$t + g(z, x, y) \subset \mathbb{P}(62, 27, 8, 7)$ $\mathbb{P}(27, 8, 7)$, Diff = $(\frac{2}{3}, 0, 0, 0)$	12
162	$z^2x + x^6y^2 + txy^5$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(28, 73, 11, 9)$ Diff = $(0, \frac{1}{2}, 0, 0)$	18
163	$z^2x + x^6y^2 + tzy^2$	$t^3 + z^2x + x^6y + tzy \subset \mathbb{P}(13, 17, 5, 9)$ Diff = $(0, 0, 0, \frac{1}{2})$	9
164	$z^2x + \underline{a}x^5y^3 + \underline{b}tx^5 + cty^6 + dy^9$ $ c + d \neq 0$	$t^3 + g(z^{1/2}, x, y^{1/3}) \subset \mathbb{P}(5, 13, 2, 5)$ Diff = $(0, \frac{1}{2}, 0, \frac{2}{3})$	10
165	$z^2x + x^5y^3 + y^{10}$	$t + zx + x^5y^3 + y^{10} \subset \mathbb{P}(50, 43, 7, 5)$ $\mathbb{P}(43, 7, 5)$, Diff = $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	30
166	$z^2x + x^5y^3 + x^2y^7$	$t + zx + x^5y^3 + x^2y^7 \subset \mathbb{P}(29, 25, 4, 3)$ $\mathbb{P}(25, 4, 3)$, Diff = $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	18
167	$z^2x + x^5y^3 + zy^5$	$t + g(z, x, y) \subset \mathbb{P}(53, 23, 7, 6)$ $\mathbb{P}(23, 7, 6)$, Diff = $(\frac{2}{3}, 0, 0, 0)$	9

168	$z^2x + x^5y^3 + txy^5$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(22, 57, 9, 7)$ Diff= $(0, \frac{1}{2}, 0, 0)$	14
169	$z^2x + \underline{a}x^4y^4 + \underline{b}tx^4y + cty^6 + dy^9$ $ c + d \neq 0$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(12, 31, 5, 4)$ Diff= $(0, \frac{1}{2}, 0, 0)$	8
170	$z^2x + x^4y^4 + y^{11}$	$t + zx + x^4y^4 + y^{11} \subset \mathbb{P}(44, 37, 7, 4)$ $\mathbb{P}(37, 7, 4)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	24
171	$z^2x + x^4y^4 + zy^5$	$t + g(z, x, y) \subset \mathbb{P}(44, 19, 6, 5)$ $\mathbb{P}(19, 6, 5)$, Diff= $(\frac{2}{3}, 0, 0, 0)$	9
172	$z^2x + x^4y^4 + zy^6$	$t + z^2x + x^4y^2 + zy^3 \subset \mathbb{P}(26, 11, 4, 5)$ $\mathbb{P}(11, 4, 5)$, Diff= $(\frac{2}{3}, 0, 0, \frac{1}{2})$	15
173	$z^2x + x^4y^4 + ty^7$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(28, 71, 13, 8)$ Diff= $(0, \frac{1}{2}, 0, 0)$	16
174	$z^2x + x^4y^4 + txy^5$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(16, 41, 7, 5)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10
175	$z^2x + \underline{a}x^4y^4 + \underline{b}tx^5 + tzy^2$	$t^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(10, 13, 4, 7)$ Diff= $(0, 0, 0, \frac{1}{2})$	7
176	$z^2x + tx^3 + y^n$ $n = 7; 11; 13; 17$	$t^3 + zx + tx^3 + y \subset \mathbb{P}(3, 7, 2, 9)$ $\mathbb{P}(3, 7, 2)$, Diff= $(0, \frac{1}{2}, 0, \frac{n-1}{n})$	7,11,14 18
177	$z^2x + tx^3 + aty^8 + bx^3y^4 + cy^{12}$ $ a + c \neq 0$	$t^3 + g(z, x, y^{1/2}) \subset \mathbb{P}(6, 7, 4, 3)$ Diff= $(0, 0, 0, \frac{1}{2})$	3
178	$z^2x + tx^3 + y^{2n}$ $n = 4, 5; 7, 8$	$t^3 + z^2x + tx^3 + y^2 \subset \mathbb{P}(6, 7, 4, 9)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	5,9
179	$z^2x + tx^3 + x^2y^n$ $n = 7, 9$	$t^3 + zx + tx^3 + x^2y \subset \mathbb{P}(3, 7, 2, 5)$ Diff= $(0, \frac{1}{2}, 0, \frac{n-1}{n})$	10
180	$z^2x + tx^3 + atzy^n + bx^2y^{2n}$ $n = 2 \ a \neq 0; n = 3 \ a \neq 0, 4$	$t^3 + g(z, x, y^{1/n}) \subset \mathbb{P}(6, 7, 4, 5)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	4,5
181	$z^2x + tx^3 + zy^n$ $n = 4; 5, 6, 7; 8, 9, 10$	$t^3 + z^2x + tx^3 + zy \subset \mathbb{P}(6, 7, 4, 11)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	4,5,7 11
182	$z^2x + tx^3y + y^{2n+1}$ $n = 3; 5, 6$	$t^3 + g(z, x, y) \subset \mathbb{P}(6n + 3, 7n + 5, 4n - 1, 9)$ Diff = $\emptyset$	5,9
183	$z^2x + tx^3y + y^n$ $n = 8; 10; 14$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(3n, 7n + 3, 2n - 3, 9)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10,12,18
184	$z^2x + tx^3y + aty^6 + bx^3y^4 + cy^9$ $ a + c \neq 0$	$t^3 + g(z, x, y) \subset \mathbb{P}(9, 11, 5, 3)$ Diff = $\emptyset$	3
185	$z^2x + tx^3y + aty^8 + bx^3y^5 + cy^{12}$ $ a + c \neq 0$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(12, 29, 7, 3)$ Diff= $(0, \frac{1}{2}, 0, 0)$	6
186	$z^2x + tx^3y + atzy^n + bx^2y^{2n+1}$ $n = 2 \ a \neq 0, 3$	$t^3 + g(z, x, y) \subset \mathbb{P}(6n + 1, 7n + 2, 4n - 1, 5)$ Diff = $\emptyset$	5
187	$z^2x + tx^3y + x^2y^8$	$t^3 + zx + tx^3y + x^2y^8 \subset \mathbb{P}(22, 53, 13, 5)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10
188	$z^2x + tx^3y + zy^n$ $n = 4; 5, 6; 7, 8$	$t^3 + g(z, x, y) \subset \mathbb{P}(6n + 1, 7n + 3, 4n - 3, 11)$ Diff = $\emptyset$	5,7,11
189	$z^2x + tx^5 + y^8$	$t^3 + z^2x + tx^5 + y^2 \subset \mathbb{P}(10, 13, 4, 15)$ Diff= $(0, 0, 0, \frac{3}{4})$	5
190	$z^2x + tx^5 + x^2y^7$	$t^3 + zx + tx^5 + x^2y \subset \mathbb{P}(5, 13, 2, 11)$ Diff= $(0, \frac{1}{2}, 0, \frac{6}{7})$	22
191	$z^2x + tx^5 + zy^5$	$t^3 + z^2x + tx^5 + zy \subset \mathbb{P}(10, 13, 4, 17)$ Diff= $(0, 0, 0, \frac{4}{5})$	10

192	$z^2x + tx^4y + y^n$ $n = 8; 10$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(4n, 10n + 3, 2n - 3, 12)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10,24
193	$z^2x + tx^4y + x^2y^7$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(26, 67, 11, 8)$ Diff= $(0, \frac{1}{2}, 0, 0)$	16
194	$z^2x + tx^4y + zy^5$	$t^3 + g(z, x, y) \subset \mathbb{P}(41, 53, 17, 14)$ Diff = $\emptyset$	7
195	$z^2x + tx^3y^2 + y^8$	$t^3 + g(z, x, y) \subset \mathbb{P}(24, 31, 10, 9)$ Diff = $\emptyset$	5
196	$z^2x + tx^3y^2 + y^{10}$	$t^3 + z^2x + tx^3y + y^5 \subset \mathbb{P}(15, 19, 7, 9)$ Diff= $(0, 0, 0, \frac{1}{2})$	9
197	$z^2x + tx^3y^2 + y^{11}$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(33, 83, 16, 9)$ Diff= $(0, \frac{1}{2}, 0, 0)$	18
198	$z^2x + tx^3y^2 + x^2y^7$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(17, 43, 8, 5)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10
199	$z^2x + tx^3y^2 + zy^5$	$t^3 + g(z, x, y) \subset \mathbb{P}(32, 41, 14, 11)$ Diff = $\emptyset$	7
200	$z^2x + tx^3y^2 + zy^6$	$t^3 + z^2x + tx^3y + zy^3 \subset \mathbb{P}(19, 24, 9, 11)$ Diff= $(0, 0, 0, \frac{1}{2})$	11
201	$z^2y + x^7 + y^n$ $n = 8; 10; 16; 20$	$t + zy + x + y^n \subset \mathbb{P}(n, n - 1, n, 1)$ $\mathbb{P}(n - 1, n, 1)$, Diff= $(\frac{2}{3}, \frac{1}{2}, \frac{6}{7}, 0)$	15,21,36 42
202	$z^2y + x^7 + ty^{n+1}(az + by^{3n+1}) +$ $cy^{6n+3}, n = 1, 2$	$t^3 + g(z, x^{1/7}, y) \subset \mathbb{P}(2n + 1, 3n + 1, 6n + 3, 1)$ $\mathbb{P}(2n + 1, 3n + 1, 1)$, Diff= $(0, 0, \frac{6}{7}, 0)$	7
203	$z^2y + x^7 + y^{2n+1}$ $n = 5, 6; 8, 9$	$t + z^2y + x + y^{2n+1} \subset \mathbb{P}(2n + 1, n, 2n + 1, 1)$ $\mathbb{P}(n, 2n + 1, 1)$, Diff= $(\frac{2}{3}, 0, \frac{6}{7}, 0)$	15,21
204	$z^2y + x^7 + aty^{4n} + by^{6n}$ $n = 2; 3$	$t^3 + g(z^{1/2}, x^{1/7}, y) \subset \mathbb{P}(2n, 6n - 1, 6n, 1)$ $\mathbb{P}(2n, 6n - 1, 1)$, Diff= $(0, \frac{1}{2}, \frac{6}{7}, 0)$	8,14
205	$z^2y + x^i(x + y^2)^j f_{7-i-j}(x, y^2)$ $i \leq 5 \& j \leq 5$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(14, 13, 2, 1)$ $\mathbb{P}(13, 2, 1)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	6
206	$z^2y + x^7 + xy^n$ $n = 7; 11; 13; 17$	$t + zy + x^7 + xy^n \subset \mathbb{P}(7n, 7n - 6, n, 6)$ $\mathbb{P}(7n - 6, n, 6)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	15,21,30 36
207	$z^2y + x^7 + ax^4y^n + bxy^{2n}$ $n = 4; 8$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(7n, 7n - 3, n, 3)$ $\mathbb{P}(7n - 3, n, 3)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	12,18
208	$z^2y + ty(ax^4 + bx^2y^3 + cy^6) +$ $x^{2i+1}f_{3-i}(x^2, y^3), b + c \neq 0 i \leq 1$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(7, 19, 3, 2)$ Diff= $(0, \frac{1}{2}, 0, 0)$	4
209	$z^2y + x^7 + ax^4y^{2n+1} + bxy^{4n+2}$ $n = 2, 3$	$t + g(z, x, y) \subset \mathbb{P}(14n + 7, 7n + 2, 2n + 1, 3)$ $\mathbb{P}(7n + 2, 2n + 1, 3)$, Diff= $(\frac{2}{3}, 0, 0, 0)$	9
210	$z^2y + x^{2i+1}f_{3-i}(x^2, y^5)$ $i \leq 2$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(35, 33, 5, 2)$ $\mathbb{P}(33, 5, 2)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	12
211	$z^2y + x^7 + atx^3y^n + bx^2y^{3n}$ $n = 2 \ b \neq 0; 4$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(7n, 21n - 5, 3n, 5)$ Diff= $(0, \frac{1}{2}, 0, 0)$	4,10
212	$z^2y + x^7 + x^2y^{2n+1}$ $n = 3; 5, 6$	$t + g(z, x, y) \subset \mathbb{P}(14n + 7, 7n + 1, 2n + 1, 5)$ $\mathbb{P}(7n + 1, 2n + 1, 5)$, Diff= $(\frac{2}{3}, 0, 0, 0)$	9,15
213	$z^2y + x^7 + x^2y^n$ $n = 8; 14$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(7n, 7n - 5, n, 5)$ $\mathbb{P}(7n - 5, n, 5)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	16,30
214	$z^2y + x^7 + atx^3y^3 + bx^2y^9$	$t^3 + g(z, x, y) \subset \mathbb{P}(21, 29, 9, 5)$ Diff = $\emptyset$	5
215	$z^2y + x^7 + x^3y^n$ $n = 5; 7; 11$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(7n, 7n - 4, n, 4)$ $\mathbb{P}(7n - 4, n, 4)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	10,18,24

216	$z^2y + x^7 + atx^2y^6 + bx^3y^9$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(21, 59, 9, 4)$ Diff= $(0, \frac{1}{2}, 0, 0)$	8
217	$z^2y + x^7 + ty^n$ $n = 5; 9, 11; 13$	$t^3 + zy + x + ty^n \subset \mathbb{P}(n, 3n - 2, 3n, 2)$ $\mathbb{P}(n, 3n - 2, 2)$, Diff= $(0, \frac{1}{2}, \frac{6}{7}, 0)$	10, 22, 28
218	$z^2y + x^7 + txy^n$ $n = 4; 6, 8; 10$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(7n, 21n - 11, 3n, 11)$ Diff= $(0, \frac{1}{2}, 0, 0)$	8, 16, 22
219	$z^2y + x^7 + txy^{2n+1}$ $n = 2; 3, 4$	$t^3 + g(z, x, y) \subset \mathbb{P}(14n + 7, 21n + 5, 6n + 3, 11)$ Diff = $\emptyset$	5, 11
220	$z^2y + x^7 + tx^2y^n$ $n = 3, 5; 7$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(7n, 21n - 8, 3n, 8)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10, 16
221	$z^2y + x^8 + ty^2(az + by^4) + cy^9$	$t^3 + g(z, x^{1/8}, y) \subset \mathbb{P}(3, 4, 9, 1)$ $\mathbb{P}(3, 4, 1)$, Diff= $(0, 0, \frac{7}{8}, 0)$	8
222	$z^2y + x^8 + ax^4y^5 + by^{10}$	$t + g(z, x^{1/2}, y) \subset \mathbb{P}(20, 9, 5, 2)$ $\mathbb{P}(9, 5, 2)$, Diff= $(\frac{2}{3}, 0, \frac{1}{2}, 0)$	12
223	$z^2y + x^8 + y^{11}$	$t + z^2y + x + y^{11} \subset \mathbb{P}(11, 5, 11, 1)$ $\mathbb{P}(11, 5, 1)$, Diff= $(\frac{2}{3}, 0, \frac{7}{8}, 0)$	24
224	$z^2y + x^8 + xy^n$ $n = 8; 10$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(8n, 8n - 7, n, 7)$ $\mathbb{P}(8n - 7, n, 7)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	18, 42
225	$z^2y + x^8 + xy^9$	$t^3 + zy + x^8 + xy^9 \subset \mathbb{P}(24, 65, 9, 7)$ Diff= $(0, \frac{1}{2}, 0, 0)$	14
226	$z^2y + x^8 + x^2y^7$	$t + g(z, x, y) \subset \mathbb{P}(56, 25, 7, 6)$ $\mathbb{P}(25, 7, 6)$, Diff= $(\frac{2}{3}, 0, 0, 0)$	9
227	$z^2y + x^8 + ax^5y^4 + bx^2y^8$	$t + g(z^{1/2}, x, y) \subset \mathbb{P}(32, 29, 4, 3)$ $\mathbb{P}(29, 4, 3)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	18
228	$z^2y + x^8 + atx^2y^4 + bx^3y^6$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(16, 43, 6, 5)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10
229	$z^2y + x^8 + x^3y^7$	$t + zy + x^8 + x^3y^7 \subset \mathbb{P}(56, 51, 7, 5)$ $\mathbb{P}(51, 7, 5)$, Diff= $(\frac{2}{3}, \frac{1}{2}, 0, 0)$	30
230	$z^2y + x^2 + ty^7$	$t^3 + z^2y + x^2 + ty^7 \subset \mathbb{P}(14, 19, 21, 4)$ Diff= $(0, 0, \frac{3}{4}, 0)$	16
231	$z^2y + x^8 + txy^n$ $n = 5; 6$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(8n, 24n - 13, 3n, 13)$ Diff= $(0, \frac{1}{2}, 0, 0)$	10, 26
232	$z^2y + zx^4 + y^n$ $n = 8; 10; 14$	$t + z^2y + zx + y^n \subset \mathbb{P}(2n, n - 1, n + 1, 2)$ $\mathbb{P}(n - 1, n + 1, 2)$, Diff= $(\frac{2}{3}, 0, \frac{3}{4}, 0)$	12, 16, 24
233	$z^2y + zx^4 + ty(azy + bx^4 + cy^5) + dx^4y^4 + ey^9, a + c + d + e \neq 0$	$t^3 + g(z, x^{1/4}, y) \subset \mathbb{P}(3, 4, 5, 1)$ Diff= $(0, 0, \frac{3}{4}, 0)$	4
234	$z^2y + zx^4 + x^{2i}y^2f_{3-i}(x^2, y^3)$ $i \leq 2$	$t + g(z, x^{1/2}, y) \subset \mathbb{P}(11, 5, 3, 1)$ $\mathbb{P}(5, 3, 1)$, Diff= $(\frac{2}{3}, 0, \frac{1}{2}, 0)$	6
235	$z^2y + zx^4 + aty^8 + by^{12}$	$t^3 + g(z, x^{1/4}, y) \subset \mathbb{P}(8, 11, 13, 2)$ Diff= $(0, 0, \frac{3}{4}, 0)$	8
236	$z^2y + zx^4 + ax^4y^6 + by^{13}$	$t + z^2y + zx + axy^6 + by^{13} \subset \mathbb{P}(13, 6, 7, 1)$ $\mathbb{P}(6, 7, 1)$, Diff= $(\frac{2}{3}, 0, \frac{3}{4}, 0)$	12
237	$z^2y + zx^4 + xy^{3n+1}$ $n = 2; 3$	$t^3 + g(z, x, y) \subset \mathbb{P}(8n + 3, 12n + 1, 3n + 2, 7)$ Diff = $\emptyset$	4, 7
238	$z^2y + zx^4 + xy^n$ $n = 8; 9, 11; 12$	$t + g(z, x, y) \subset \mathbb{P}(8n + 1, 4n - 3, n + 1, 7)$ $\mathbb{P}(4n - 3, n + 1, 7)$, Diff= $(\frac{2}{3}, 0, 0, 0)$	13, 18, 21
239	$z^2y + zx^4 + x^2y^n$ $n = 6; 10$	$t + g(z, x^{1/2}, y) \subset \mathbb{P}(4n + 1, 2n - 1, n + 1, 3)$ $\mathbb{P}(2n - 1, n + 1, 3)$, Diff= $(\frac{2}{3}, 0, \frac{1}{2}, 0)$	10, 18

240	$z^2y + zx^4 + ax^5y^n + bx^2y^{2n+1}$ $n = 3 \text{ } b \neq 0; 4$	$t + g(z, x, y) \subset \mathbb{P}(8n + 5, 4n + 1, n + 1, 3)$ $\mathbb{P}(4n + 1, n + 1, 3), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	6,9
241	$z^2y + zx^4 + x^3y^n$ $n = 5; 7; 8$	$t + g(z, x, y) \subset \mathbb{P}(8n + 3, 4n - 1, n + 1, 5)$ $\mathbb{P}(4n - 1, n + 1, 5), \text{Diff}=(\frac{2}{3}, 0, 0, 0)$	9,12,15
242	$z^2y + zx^4 + atx^2y^4 + bx^3y^6$	$t^3 + g(z, x, y) \subset \mathbb{P}(17, 23, 7, 5)$ $\text{Diff} = \emptyset$	5
243	$z^2y + zx^4 + ty^n$ $n = 5; 7; 9$	$t^3 + g(z, x^{1/4}, y) \subset \mathbb{P}(2n, 3n - 2, 3n + 2, 4)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	10,13,16
244	$z^2y + zx^4 + txy^n$ $n = 4; 5, 6$	$t^3 + g(z, x, y) \subset \mathbb{P}(8n + 1, 12n - 5, 3n + 2, 13)$ $\text{Diff} = \emptyset$	7,10
245	$z^2y + zx^4 + tx^2y^n$ $n = 3; 5$	$t^3 + g(z, x^{1/2}, y) \subset \mathbb{P}(4n + 1, 6n - 1, 3n + 2, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	7,10
246	$z^2y + tx^5 + y^n$ $n = 8; 14$	$t^3 + zy + tx + y^n \subset \mathbb{P}(n, 3n - 3, 2n, 3)$ $\text{Diff}=(0, \frac{1}{2}, \frac{4}{5}, 0)$	15,30
247	$z^2y + tx^5 + atx^2y^4 + bx^6y^2 +$ $cx^3y^6 + dy^{10}$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(10, 27, 4, 3)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	3
248	$z^2y + tx^5 + y^{11}$	$t^3 + z^2y + tx + y^{11} \subset \mathbb{P}(11, 15, 22, 3)$ $\text{Diff}=(0, 0, \frac{4}{5}, 0)$	11
249	$z^2y + tx^5 + aty^8 + bx^5y^4 + cy^{12}$	$t^3 + g(z^{1/2}, x, y^{1/5}) \subset \mathbb{P}(4, 11, 8, 1)$ $\text{Diff}=(0, \frac{1}{2}, \frac{4}{5}, 0)$	10
250	$z^2y + tx^5 + xy^n$ $n = 8; 10; 12$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(5n, 15n - 13, 2n, 13)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	16,22,26
251	$z^2y + tx^5 + xy^9$	$t^3 + g(z, x, y) \subset \mathbb{P}(45, 61, 18, 13)$ $\text{Diff} = \emptyset$	9
252	$z^2y + tx^5 + x^2y^n$ $n = 6; 8; 10$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(5n, 15n - 11, 2n, 11)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	12,18,22
253	$z^2y + tx^5 + x^2y^7$	$t^3 + g(z, x, y) \subset \mathbb{P}(35, 47, 14, 11)$ $\text{Diff} = \emptyset$	7
254	$z^2y + tx^5 + x^3y^5$	$t^3 + g(z, x, y) \subset \mathbb{P}(25, 33, 10, 9)$ $\text{Diff} = \emptyset$	5
255	$z^2y + tx^5 + x^3y^8$	$t^3 + zy + tx^5 + x^3y^8 \subset \mathbb{P}(40, 111, 16, 9)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	18
256	$z^2y + tx^5 + x^4y^n$ $n = 4; 6$	$t^3 + g(z^{1/2}, x, y) \subset \mathbb{P}(5n, 15n - 7, 2n, 7)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	10,14

(0) $n = 1 \& k \leq 1 \& (a \neq 0 \parallel (i = 0 \& j \leq 1))$, $n = 2 \& k \leq 1 \& (a \neq 0 \parallel (i \leq 1 \& j \leq 4))$.

(1) The singularity is $t^3 + z^2x + \underline{a}tx^i(x + y^2)^j f_{3-i-j}(x, y^2) + \underline{b}x^k y(x + y^2)^l f_{4-k-l}(x, y^2)$. The common exceptionality condition for two cases is $(i = 0 \parallel k \leq 1)$. The first case is $(b \neq 0 \parallel j \leq 2)$ and the coefficient $\underline{a}$ is absent (see the comments to the tables). If the coefficient $\underline{b}$ is absent then we have the second case.

14. Singularity $-tz^i f_{2-i}(t, z) + g(z, x, y)(i \leq 1)$

1	$x^4 + y^n$ $n = 5; 7; 11$	$tz^i f_{2-i}(t, z) + x + y \subset \mathbb{P}(1, 1, 3, 3)$ $\mathbb{P}(1, 1, 3), \text{Diff}=(0, 0, \frac{3}{4}, \frac{n-1}{n})$	5,8,12
2	$f_3(t, z) + y^2 f_2(t, z) +$ (1) $(t + z)(x^2 y + y^4) + f_2(x^2, y^3)$	$tz^i f_{2-i}(t, z) + g(z, x^{1/2}, y) \subset \mathbb{P}(2, 2, 3, 1)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	2
3	$x^j f_{4-j}(x, y^2)$ $j \leq 2$	$tz^i f_{2-i}(t, z) + g(z, x, y^{1/2}) \subset \mathbb{P}(4, 4, 3, 3)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	3
4	$t^3 + ktz^2 + lz^3 + (at + bz)y^6 +$	$tz^i f_{2-i}(t, z) + g(z, x^{1/4}, y^{1/3}) \subset \mathbb{P}(1, 1, 3, 1)$	4

	$ctzy^3 + dy^9 + x^4$	$\mathbb{P}^2, \text{Diff}=(0, 0, \frac{3}{4}, \frac{2}{3})$	
5	$x^{2j}f_{2-j}(x^2, y^5)$ $j \leq 1$	$tz^i f_{2-i}(t, z) + x^j f_{2-j}(x, y) \subset \mathbb{P}(2, 2, 3, 3)$ $\text{Diff}=(0, 0, \frac{1}{2}, \frac{4}{5})$	6
6	$x^4 + xy^n$ $n = 4, 5; 7, 8$	$tz^i f_{2-i}(t, z) + x^4 + xy \subset \mathbb{P}(4, 4, 3, 9)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5,9
7	$x^4 + zy^n$ $n = 3 \ i = 0; 5; 7$	$tz^i f_{2-i}(t, z) + x + zy \subset \mathbb{P}(1, 1, 3, 2)$ $\mathbb{P}(1, 1, 2), \text{Diff}=(0, 0, \frac{3}{4}, \frac{n-1}{n})$	4,5,8
8	$x^4 + zxy^n$ $n = 2 \ i = 0; n = 3, 4$	$tz^i f_{2-i}(t, z) + x^4 + zxy \subset \mathbb{P}(4, 4, 3, 5)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	2,5
9	$x^j f_{5-j}(x, y)$ $j = 1$	$tz^i f_{2-i}(t, z) + x^j f_{5-j}(x, y) \subset \mathbb{P}(5, 5, 3, 3)$ $\text{Diff} = \emptyset$	3
10	$t^3 + ktz^2 + lz^3 + (at + bz)y^4 +$ $ctzy^2 + dy^6 + x^5$	$tz^i f_{2-i}(t, z) + g(z, x^{1/5}, y^{1/2}) \subset \mathbb{P}(1, 1, 3, 1)$ $\mathbb{P}^2, \text{Diff}=(0, 0, \frac{4}{5}, \frac{1}{2})$	5
11	$x^5 + y^7$	$tz^i f_{2-i}(t, z) + x + y \subset \mathbb{P}(1, 1, 3, 3)$ $\mathbb{P}(1, 1, 3), \text{Diff}=(0, 0, \frac{4}{5}, \frac{6}{7})$	15
12	$x^5 + xy^5$	$tz^i f_{2-i}(t, z) + x^5 + xy \subset \mathbb{P}(5, 5, 3, 12)$ $\text{Diff}=(0, 0, 0, \frac{4}{5})$	6
13	$x^5 + x^2y^4$	$tz^i f_{2-i}(t, z) + x^5 + x^2y \subset \mathbb{P}(5, 5, 3, 9)$ $\text{Diff}=(0, 0, 0, \frac{3}{4})$	9
14	$x^5 + zxy^3$	$tz^i f_{2-i}(t, z) + x^5 + zxy \subset \mathbb{P}(5, 5, 3, 7)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	7
15	$t^3 + ktz^2 + lz^3 + (at + bz)y^4 +$ $ctzy^2 + dy^6 + x^4y$	$tz^i f_{2-i}(t, z) + g(z, x^{1/4}, y) \subset \mathbb{P}(2, 2, 5, 1)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	4
16	$x^4y + ax^2y^4 + by^7$ $b \neq 0$	$tz^i f_{2-i}(t, z) + g(z, x^{1/2}, y) \subset \mathbb{P}(7, 7, 9, 3)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	6
17	$x^4y + y^8$	$tz^i f_{2-i}(t, z) + g(z, x^{1/4}, y) \subset \mathbb{P}(8, 8, 21, 3)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	12
18	$x^4y + xy^n$ $n = 5; 6$	$tz^i f_{2-i}(t, z) + g(z, x, y) \subset \mathbb{P}(4n - 1, 4n - 1, 3n - 3, 9)$ $\text{Diff} = \emptyset$	6,9
19	$x^4y + zy^5$	$tz^i f_{2-i}(t, z) + g(z, x^{1/4}, y) \subset \mathbb{P}(5, 5, 13, 2)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	8
20	$x^4y + zxy^3$	$tz^i f_{2-i}(t, z) + x^4y + zxy^3 \subset \mathbb{P}(11, 11, 7, 5)$ $\text{Diff} = \emptyset$	5
21	$zx^3 + y^n$ $n = 5 \ i = 0; 7; 8$	$tz^i f_{2-i}(t, z) + zx + y \subset \mathbb{P}(1, 1, 2, 3)$ $\mathbb{P}(1, 1, 2), \text{Diff}=(0, 0, \frac{2}{3}, \frac{n-1}{n})$	5,7,9
22	$t^3 + ktz^2 + lz^3 + tx^3 + aty^4 + sy^6$ $+ z(bx^3 + cy^4) + dx^3y^2 + etzy^2$	$tz^i f_{2-i}(t, z) + g(z, x^{1/3}, y^{1/2}) \subset \mathbb{P}(1, 1, 2, 1)$ $\text{Diff}=(0, 0, \frac{2}{3}, \frac{1}{2})$	3
23	$zx^3 + xy^n$ $n = 4 \ i = 0; 5; 6$	$tz^i f_{2-i}(t, z) + zx^3 + xy \subset \mathbb{P}(3, 3, 2, 7)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	4,5,7
24	$zx^3 + x^2y^n$ $n = 3 \ i = 0; 4$	$tz^i f_{2-i}(t, z) + zx^3 + x^2y \subset \mathbb{P}(3, 3, 2, 5)$ $\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	3,5

(1) See example 3.26.

(2) If $d = e = 0$ then we require that the polynomial without monomial y^6 is irreducible.

15. Singularity $-t^2z + g(t, z, x, y)$

1	$z^i x^j f_{4-i-j}(x, y) + y^n$ $n = 5 \ i = 0 \& j \leq 2; 7$	$tz + z^i x^j f_{4-i-j}(x, y) + y \subset \mathbb{P}(3, 1, 1, 4)$ $\mathbb{P}(3, 1, 1), \text{Diff}=(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	5,8
2	$f_4(z, x) + y^3 f_2(z, x) +$ $+ y^6$	$tz + g(z, x, y^{1/3}) \subset \mathbb{P}(3, 1, 1, 2)$ $\text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	4
3	$z^i x^j f_{4-i-j}(x, y) + ty^n$	$t^2z + z^i x^j f_{4-i-j}(x, y) + ty \subset \mathbb{P}(3, 2, 2, 5)$	3,5

	$n = 3 \ i = 0 \& j \leq 1; 4$	$\text{Diff}=(0, 0, 0, \frac{n-1}{n})$	
4	$z^4 + x^3y + y^n$ $n = 5; 9; 11; 15$	$tz + z^4 + xy + y^n \subset \mathbb{P}(3n, n, 4n - 4, 4)$ $\text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	7,15,16 24
5	$z^4 + x^3y + az^2y^n + by^{2n}$ $n = 3 \ b \neq 0; 7$	$tz + g(z, x^{1/3}, y) \subset \mathbb{P}(3n, n, 4n - 2, 2)$ $\text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	4,12
6	$z^4 + x^i y f_{3-i}(x, y^n)$ $n = 2; 4$	$tz + g(z, x, y) \subset \mathbb{P}(9n + 3, 3n + 1, 4n, 4)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	7,8
7	$z^i f_{4-i}(z, y^2) + x^3y + aty^5$ $a \neq 0 \ \ i \leq 1$	$t^2z + g(z, x^{1/3}, y) \subset \mathbb{P}(3, 2, 7, 1)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	3
8	$z^i f_{4-i}(z, y^3) + x^3y$ $i \leq 2$	$tz + z^i f_{4-i}(z, y^3) + xy \subset \mathbb{P}(9, 3, 11, 1)$ $\text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	6
9	$z^4 + x^3y + xy^n$ $n = 6; 8$	$tz + g(z, x, y) \subset \mathbb{P}(9n - 3, 3n - 1, 4n - 4, 8)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,16
10	$z^4 + x^3y + zxy^4$	$tz + g(z, x, y) \subset \mathbb{P}(33, 11, 13, 5)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
11	$z^4 + x^3y + aty^7 + bxyz^5$	$t^2z + g(z, x, y) \subset \mathbb{P}(21, 14, 17, 5)$ $\text{Diff} = \emptyset$	5
12	$z^4 + x^3y + ty^n$ $n = 3; 6; 8, 9$	$t^2z + g(z, x^{1/3}, y) \subset \mathbb{P}(3n, 2n, 8n - 5, 5)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	6,9,15
13	$z^4 + x^3y + txy^n$ $n = 2; 3, 4$	$t^2z + g(z, x, y) \subset \mathbb{P}(9n - 3, 6n - 2, 8n - 5, 7)$ $\text{Diff} = \emptyset$	5,7
14	$z^4 + x^i y^j f_{5-i-j}(x, y)$ $i \leq 2 \& j \leq 2$	$tz + g(z, x, y) \subset \mathbb{P}(15, 5, 4, 4)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8
15	$z^4 + x^5 + zxy^3$	$tz + z^4 + x^5 + zxy \subset \mathbb{P}(15, 5, 4, 11)$ $\text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	22
16	$z^4 + x^3y^2 + az^2y^3 + by^6$	$tz + z^4 + xy^2 + az^2y^3 + by^6 \subset \mathbb{P}(9, 3, 8, 2)$ $\text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	12
17	$z^4 + x^3y^2 + y^7$	$tz + z^4 + xy^2 + y^7 \subset \mathbb{P}(21, 7, 20, 4)$ $\text{Diff}=(\frac{1}{2}, 0, \frac{2}{3}, 0)$	24
18	$z^4 + x^3y^2 + ty^4$	$t^2z + z^4 + xy + ty^2 \subset \mathbb{P}(6, 4, 11, 5)$ $\text{Diff}=(0, 0, \frac{2}{3}, \frac{1}{2})$	15
19	$z^4 + tx^3 + y^5$	$t^2z + z^4 + tx + y \subset \mathbb{P}(3, 2, 5, 8)$ $\mathbb{P}(3, 2, 5), \text{Diff}=(0, 0, \frac{2}{3}, \frac{4}{5})$	15
20	$z^4 + tx^3 + xy^4$	$t^2z + z^4 + tx^3 + xy \subset \mathbb{P}(9, 6, 5, 19)$ $\text{Diff}=(0, 0, 0, \frac{3}{4})$	8
21	$z^4 + tx^3 + x^2y^3$	$t^2z + z^4 + tx^3 + x^2y \subset \mathbb{P}(9, 6, 5, 14)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	7
22	$z^4 + tx^3 + zxy^3$	$t^2z + z^4 + tx^3 + zxy \subset \mathbb{P}(9, 6, 5, 13)$ $\text{Diff}=(0, 0, 0, \frac{2}{3})$	13
23	$z^4 + tx^2y + y^n$ $n = 5; 7$	$t^2z + g(z, x^{1/2}, y) \subset \mathbb{P}(3n, 2n, 5n - 8, 8)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	10,16
24	$z^4 + tx^2y + az^2y^3 + by^6$	$t^2z + g(z, x^{1/2}, y) \subset \mathbb{P}(9, 6, 11, 4)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	8
25	$z^4 + tx^2y + xy^5$	$t^2z + g(z, x, y) \subset \mathbb{P}(27, 18, 17, 11)$ $\text{Diff} = \emptyset$	11
26	$z^3x + x^3y + y^{2n+1}$ $n = 2; 4, 5$	$t^2z + g(z, x, y) \subset \mathbb{P}(7n + 3, 4n + 3, 6n, 9)$ $\text{Diff} = \emptyset$	3,9
27	$z^3x + x^3y + y^n$	$tz + g(z, x, y) \subset \mathbb{P}(7n - 1, 2n + 1, 3n - 3, 9)$	8,11,18

	$n = 6; 8; 12$	$\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	
28	$z^3x + az^3y^3 + x^iyf_{3-i}(x, y^3)$ $i \leq 2$	$tz + g(z, x, y) \subset \mathbb{P}(23, 7, 9, 3)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
29	$z^3x + x^3y + z^2y^5$	$t^2z + g(z, x, y) \subset \mathbb{P}(18, 11, 14, 5)$ $\text{Diff} = \emptyset$	5
30	$z^3x + x^3y + z^2y^6$	$tz + g(z, x, y) \subset \mathbb{P}(43, 13, 17, 5)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
31	$z^3x + x^3y + ty^n$ $n = 4; 6$	$t^2z + g(z, x, y) \subset \mathbb{P}(7n - 1, 4n + 1, 6n - 4, 11)$ $\text{Diff} = \emptyset$	5, 11
32	$z^3x + x^iy^jf_{5-i-j}(x, y)$ $i \leq 1 \& j \leq 2$	$tz + g(z, x, y) \subset \mathbb{P}(11, 4, 3, 3)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
33	$z^3x + x^4y + y^6$	$tz + g(z, x, y) \subset \mathbb{P}(53, 19, 15, 12)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	24
34	$z^3x + x^3y^2 + y^7$	$tz + g(z, x, y) \subset \mathbb{P}(47, 16, 15, 9)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	18
35	$z^3x + x^3y^2 + z^2y^3$	$tz + g(z, x, y) \subset \mathbb{P}(23, 8, 7, 5)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
36	$z^3x + x^3y^2 + ty^4$	$t^2z + z^3x + x^3y + ty^2 \subset \mathbb{P}(13, 9, 8, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
37	$z^3x + tx^3 + y^n$ $n = 5; 6$	$t^2z + z^3x + tx^3 + y \subset \mathbb{P}(7, 5, 4, 19)$ $\mathbb{P}(7, 5, 4), \text{Diff}=(0, 0, 0, \frac{n-1}{n})$	5, 19
38	$z^3x + tx^2y + y^6$	$t^2z + g(z, x, y) \subset \mathbb{P}(29, 20, 18, 13)$ $\text{Diff} = \emptyset$	9
39	$z^3y + x^5 + y^6$	$tz + z^3y + x + y^6 \subset \mathbb{P}(13, 5, 18, 3)$ $\mathbb{P}(13, 5, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	30
40	$z^3y + x^5 + zxy^3$	$tz + g(z, x, y) \subset \mathbb{P}(29, 11, 8, 7)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
41	$z^3y + z^2x^2 + azxy^3 + x^3y^2 + cy^6$	$tz + g(z, x, y) \subset \mathbb{P}(13, 5, 4, 3)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
42	$z^3y + tx^3 + y^n$ $n = 5; 6$	$t^2z + g(z, x^{1/3}, y) \subset \mathbb{P}(2n + 1, 2n - 2, 4n - 1, 6)$ $\text{Diff}=(0, 0, \frac{2}{3}, 0)$	6, 9
43	$z^2x^2 + x^iy^jf_{5-i-j}(x, y)$ $i \leq 1 \& j \leq 2$	$tz + g(z, x, y) \subset \mathbb{P}(7, 3, 2, 2)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
44	$z^2x^2 + x^5 + azxy^3 + by^6$	$tz + z^2x^2 + x^5 + azxy + by^2 \subset \mathbb{P}(7, 3, 2, 5)$ $\text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	10
45	$z^2x^2 + x^5 + xy^5$	$tz + z^2x^2 + x^5 + xy \subset \mathbb{P}(7, 3, 2, 8)$ $\text{Diff}=(\frac{1}{2}, 0, 0, \frac{4}{5})$	16
46	$z^2x^2 + x^5 + ty^4$	$t^2z + z^2x^2 + x^5 + ty \subset \mathbb{P}(7, 6, 4, 13)$ $\text{Diff}=(0, 0, 0, \frac{3}{4})$	13
47	$z^2x^2 + x^4y + azxy^3 + by^6$	$tz + g(z, x, y) \subset \mathbb{P}(17, 7, 5, 4)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8
48	$z^2x^2 + x^4y + ty^4$	$t^2z + z^2x + x^2y + ty^4 \subset \mathbb{P}(11, 9, 13, 5)$ $\text{Diff}=(0, 0, \frac{1}{2}, 0)$	10
49	$z^2x^2 + x^3y^2 + y^7$	$tz + g(z, x, y) \subset \mathbb{P}(31, 11, 10, 6)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	12
50	$z^2x^2 + x^3y^2 + az^2y^3 + bxy^5$	$tz + g(z, x, y) \subset \mathbb{P}(19, 7, 6, 4)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8
51	$z^2x^2 + x^3y^2 + ty^4$	$t^2z + z^2x^2 + x^3y + ty^2 \subset \mathbb{P}(8, 6, 5, 7)$	7

		$\text{Diff}=(0,0,0,\frac{1}{2})$	
52	$z^2x^2+tx^2y+y^n$ $n=5;7$	$t^2z+g(z,x^{1/2},y)\subset\mathbb{P}(2n-1,n+2,3n-4,5)$ $\text{Diff}=(0,0,\frac{1}{2},0)$	6,10
53	$z^2x^2+tx^2y+y^6$	$t^2z+g(z,x,y)\subset\mathbb{P}(11,8,7,5)$ $\text{Diff}=\emptyset$	5
54	$z^2x^2+tx^2y+xy^5$	$t^2z+g(z,x,y)\subset\mathbb{P}(17,12,11,7)$ $\text{Diff}=\emptyset$	7
55	$z^2y^2+x^5+y^n$ $n=7;9$	$tz+z^2y^2+x+y^n\subset\mathbb{P}(n+2,n-2,2n,2)$ $\mathbb{P}(n+2,n-2,3), \text{Diff}=(\frac{1}{2},0,\frac{4}{5},0)$	15,20
56	$z^2y^2+x^5+azy^5+by^8$	$tz+z^2y^2+x+azy^5+by^8\subset\mathbb{P}(5,3,8,1)$ $\mathbb{P}(5,3,1), \text{Diff}=(\frac{1}{2},0,\frac{4}{5},0)$	10
57	$z^2y^2+x^5+xy^n$ $n=5;7$	$tz+g(z,x,y)\subset\mathbb{P}(5n+8,5n-8,2n,8)$ $\text{Diff}=(\frac{1}{2},0,0,0)$	11,16
58	$z^2y^2+x^5+ax^3y^3+bx^2y^6$	$tz+g(z,x,y)\subset\mathbb{P}(19,11,6,4)$ $\text{Diff}=(\frac{1}{2},0,0,0)$	8
59	$z^2y^2+x^5+azxy^3+bx^2y^4$	$tz+g(z,x,y)\subset\mathbb{P}(13,7,4,3)$ $\text{Diff}=(\frac{1}{2},0,0,0)$	6
60	$z^2y^2+x^5+x^2y^5$	$tz+g(z,x,y)\subset\mathbb{P}(31,19,10,6)$ $\text{Diff}=(\frac{1}{2},0,0,0)$	12
61	$z^2y^2+x^5+ty^5$	$t^2z+z^2y^2+x+ty^5\subset\mathbb{P}(7,8,22,3)$ $\mathbb{P}(7,8,3), \text{Diff}=(0,0,\frac{4}{5},0)$	11
62	$z^2y^2+x^5+ty^6$	$t^2z+z^2y^2+x+ty^3\subset\mathbb{P}(4,5,13,3)$ $\mathbb{P}(4,5,3), \text{Diff}=(0,0,\frac{4}{5},\frac{1}{2})$	15
63	$z^2y^2+x^5+txy^3$	$t^2z+g(z,x,y)\subset\mathbb{P}(23,24,14,11)$ $\text{Diff}=\emptyset$	7
64	$z^2y^2+x^5+txy^4$	$t^2z+z^2y^2+x^5+txy^2\subset\mathbb{P}(14,17,9,11)$ $\text{Diff}=(0,0,0,\frac{1}{2})$	11
65	$z^2y^2+x^5+tx^2y^2$	$t^2z+z^2y^2+x^5+tx^2y\subset\mathbb{P}(8,9,5,7)$ $\text{Diff}=(0,0,0,\frac{1}{2})$	7
66	$z^2xy+x^iy^jf_{5-i-j}(x,y)$ $i\leq 1\&j\leq 1$	$tz+g(z,x,y)\subset\mathbb{P}(7,3,2,2)$ $\text{Diff}=(\frac{1}{2},0,0,0)$	4
67	$z^2xy+x^5+y^n$ $n=6;7$	$tz+g(z,x,y)\subset\mathbb{P}(6n+5,4n-5,2n,10)$ $\text{Diff}=(\frac{1}{2},0,0,0)$	8,20
68	$z^2xy+x^5+ty^4$	$t^2z+g(z,x,y)\subset\mathbb{P}(29,27,17,14)$ $\text{Diff}=\emptyset$	7
69	$z^2x+x^i(x+y^3)^jf_{4-i-j}(x,y^3)$ $i\leq 2\&j\leq 3$	$tz+g(z,x,y^{1/3})\subset\mathbb{P}(5,3,2,2)$ $\text{Diff}=(\frac{1}{2},0,0,\frac{2}{3})$	4
70	$z^2x+x^4+y^n$ $n=5;7;9;11;13,15$	$tz+z^2x+x^4+y\subset\mathbb{P}(5,3,2,8)$ $\mathbb{P}(5,3,2), \text{Diff}=(\frac{1}{2},0,0,\frac{n-1}{n})$	5,7,10 11,16
71	$z^2x+x^4+ax^2y^n+by^{2n}$ $n=3\ b\neq 0;5,7$	$tz+z^2x+x^4+ax^2y+by^2\subset\mathbb{P}(5,3,2,4)$ $\text{Diff}=(\frac{1}{2},0,0,\frac{n-1}{n})$	3,8
72	$z^2x+azx^i(x+y^2)^jy^{2k+1}f_{2-i-j-k}(x,y^2)$ $+x^l(x+y^2)^nf_{4-l-n}(x,y^2)$ $(i=0 l\leq 1)\&(j=0 n\leq 2)$	$tz+g(z,x,y)\subset\mathbb{P}(5,3,2,1)$ $\text{Diff}=(\frac{1}{2},0,0,0)$	2
73	$z^2x+x^4+ty^n$ $n=4,5;6;7;8,9,10$	$t^2z+z^2x+x^4+ty\subset\mathbb{P}(5,6,4,11)$ $\text{Diff}=(0,0,0,\frac{n-1}{n})$	5,6,7 11
74	$z^2x+x^4+txy^n$ $n=4,5,6$	$t^2z+z^2x+x^4+txy\subset\mathbb{P}(5,6,4,7)$ $\text{Diff}=(0,0,0,\frac{n-1}{n})$	7

74 (2)	$z^2x + zf_3(x, y) + f_5(x, y)$	$tz + g(z, x, y) \subset \mathbb{P}(3, 2, 1, 1)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	2
75 (3)	$\frac{z^2x + zx^3 + x^5}{n=3} + atxy^n + by^{2n}$ $n=3, b \neq 0; 5$	$t^2z + g(z, x, y^{1/n}) \subset \mathbb{P}(3, 4, 2, 5)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	3,5
76 (3)	$\frac{z^2x + zx^3 + x^5}{n=7} + y^n$ $n=7; 9$	$tz + \frac{z^2x + zx^3 + x^5}{n=7} + y \subset \mathbb{P}(3, 2, 1, 5)$ $\mathbb{P}(3, 2, 1)$, Diff= $(\frac{1}{2}, 0, 0, \frac{n-1}{n})$	7,10
77 (3)	$\frac{z^2x + zx^3 + x^5}{n=4^*} + ty^n$ $n=4^*; 5; 6$	$t^2z + \frac{z^2x + zx^3 + x^5}{n=4^*} + ty \subset \mathbb{P}(3, 4, 2, 7)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	4,5,7
78	$z^2x + x^4y + y^n$ $n=6; 8; 10; 12$	$tz + g(z, x, y) \subset \mathbb{P}(5n-1, 3n+1, 2n-2, 8)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	7,10,12 16
79	$z^2x + x^4y + ax^2y^{n+1} + by^{2n+1}$ $n=3, b \neq 0; 5$	$tz + g(z, x, y) \subset \mathbb{P}(5n+2, 3n+2, 2n, 4)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	4,8
80	$z^2x + x^i(x+y)^j y^k f_{4-i-j}(x, y^2)$ $i \leq 2 \& j \leq 3$	$tz + g(z, x, y) \subset \mathbb{P}(11, 7, 4, 2)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	4
81	$z^2x + x^4y + ty^n$ $n=4; 5; 6; 7, 8$	$t^2z + g(z, x, y) \subset \mathbb{P}(5n-1, 6n+1, 4n-3, 11)$ Diff = $\emptyset$	5,6,7 11
82	$z^2x + x^4y + txy^n$ $n=4, 5$	$t^2z + g(z, x, y) \subset \mathbb{P}(5n-2, 6n-1, 4n-3, 7)$ Diff = $\emptyset$	7
83	$z^2x + x^i(x+y)^j y^k f_{6-i-j-k}(x, y)$ $i \leq 1 \& j \leq 2 \& k \leq 2$	$tz + g(z, x, y) \subset \mathbb{P}(7, 5, 2, 2)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	4
84	$z^2x + x^6 + y^7$	$tz + z^2x + x^6 + y \subset \mathbb{P}(7, 5, 2, 12)$ $\mathbb{P}(7, 5, 2)$, Diff= $(\frac{1}{2}, 0, 0, \frac{6}{7})$	8
85	$z^2x + x^6 + x^2y^5$	$tz + z^2x + x^6 + x^2y \subset \mathbb{P}(7, 5, 2, 8)$ Diff= $(\frac{1}{2}, 0, 0, \frac{4}{5})$	16
86	$z^2x + x^6 + ty^5$	$t^2z + z^2x + x^6 + ty \subset \mathbb{P}(7, 10, 4, 17)$ Diff= $(0, 0, 0, \frac{4}{5})$	10
87	$z^2x + x^6 + txy^4$	$t^2z + z^2x + x^6 + txy \subset \mathbb{P}(7, 10, 4, 13)$ Diff= $(0, 0, 0, \frac{3}{4})$	13
88	$z^2x + x^5y + y^n$ $n=7; 8$	$tz + g(z, x, y) \subset \mathbb{P}(6n-1, 4n+1, 2n-2, 10)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	8,10
89	$z^2x + x^5y + ty^5$	$t^2z + g(z, x, y) \subset \mathbb{P}(29, 41, 17, 14)$ Diff = $\emptyset$	7
90	$z^2x + x^4y^2 + y^n$ $n=7; 9$	$tz + g(z, x, y) \subset \mathbb{P}(5n-2, 3n+2, 2n-4, 8)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	11,16
91	$z^2x + x^4y^2 + ax^2y^5 + by^8$	$tz + g(z, x, y) \subset \mathbb{P}(19, 13, 6, 4)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	8
92	$z^2x + x^4y^2 + ty^5$	$t^2z + g(z, x, y) \subset \mathbb{P}(23, 32, 14, 11)$ Diff = $\emptyset$	7
93	$z^2x + x^4y^2 + ty^6$	$t^2z + z^2x + x^4y + ty^3 \subset \mathbb{P}(14, 19, 9, 11)$ Diff= $(0, 0, 0, \frac{1}{2})$	11
94	$z^2x + x^4y^2 + txy^4$	$t^2z + z^2x + x^4y + txy^2 \subset \mathbb{P}(8, 11, 5, 7)$ Diff= $(0, 0, 0, \frac{1}{2})$	7
95	$z^2x + tx^3 + y^n$ $n=5; 6; 7; 8; 9; 10; 11, 12$	$t^2z + z^2x + tx^3 + y \subset \mathbb{P}(4, 5, 3, 13)$ $\mathbb{P}(4, 5, 3)$, Diff= $(0, 0, 0, \frac{n-1}{n})$	5,6,7,8 9,10,13
96	$z^2x + tx^3 + x^2y^n$ $n=5, 6$	$t^2z + z^2x + tx^3 + x^2y \subset \mathbb{P}(4, 5, 3, 7)$ Diff= $(0, 0, 0, \frac{n-1}{n})$	7
97	$z^2x + tx^4 + y^n$ $n=6; 7; 8$	$t^2z + z^2x + tx^4 + y \subset \mathbb{P}(5, 7, 3, 17)$ $\mathbb{P}(5, 7, 3)$, Diff= $(0, 0, 0, \frac{n-1}{n})$	6,7,17

98	$z^2x + tx^4 + x^2y^5$	$t^2z + z^2x + tx^4 + x^2y \subset \mathbb{P}(5, 7, 3, 11)$ Diff=(0, 0, 0, $\frac{4}{5}$)	11
99	$z^2x + tx^3y + y^n$ $n = 6; 7; 8; 9$	$t^2z + g(z, x, y) \subset \mathbb{P}(4n - 1, 5n + 2, 3n - 4, 13)$ Diff = $\emptyset$	7,9,10 13
100	$z^2x + tx^3y + x^2y^5$	$t^2z + g(z, x, y) \subset \mathbb{P}(17, 23, 11, 7)$ Diff = $\emptyset$	7
101	$z^2y + x^5 + y^n$ $n = 6; 8; 12; 14$	$tz + z^2y + x + y^n \subset \mathbb{P}(n + 1, n - 1, 2n, 2)$ $\mathbb{P}(n + 1, n - 1, 2), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	8,10,16 20
102	$z^2y + x^5 + ty^n(az + by^{3n-1}) +$ $czy^{2n} + dy^{4n-1}, n = 2, 3$	$t^2z + g(z, x^{1/5}, y) \subset \mathbb{P}(n, 2n - 1, 4n - 1, 1)$ $\mathbb{P}(n, 2n - 1, 1), \text{Diff}=(0, 0, \frac{4}{5}, 0)$	5
103	$z^2y + x^5 + azyn + by^{2n-1}$ $n = 5; 7$	$tz + g(z, x^{1/5}, y) \subset \mathbb{P}(n, n - 1, 2n - 1, 1)$ $\mathbb{P}(n, n - 1, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	6,10
104	$z^2y + x^i(x + y^2)^j f_{5-i-j}(x, y^2)$ $i \leq 3 \& j \leq 3$	$tz + g(z, x, y) \subset \mathbb{P}(11, 9, 4, 2)$ Diff=($\frac{1}{2}$, 0, 0, 0)	4
105	$z^2y + x^5 + xy^n$ $n = 5; 7; 9; 11$	$tz + g(z, x, y) \subset \mathbb{P}(5n + 4, 5n - 4, 2n, 8)$ Diff=($\frac{1}{2}$, 0, 0, 0)	6,11,12 16
106	$z^2y + x^5 + ax^3y^n + bxy^{2n}$ $n = 3 \ b \neq 0; 5$	$tz + g(z, x, y) \subset \mathbb{P}(5n + 2, 5n - 2, 2n, 4)$ Diff=($\frac{1}{2}$, 0, 0, 0)	4,8
107	$z^2y + x^5 + x^2y^n$ $n = 4; 8$	$tz + g(z, x, y) \subset \mathbb{P}(5n + 3, 5n - 3, 2n, 6)$ Diff=($\frac{1}{2}$, 0, 0, 0)	7,12
108	$z^2y + \underline{a}tx^3y + \underline{b}x^5 + cty^6 + dzyx^3$ $+ex^2y^5, c + d + e \neq 0$	$t^2z + g(z, x, y) \subset \mathbb{P}(7, 11, 5, 3)$ Diff = $\emptyset$	3
109	$z^2y + x^5 + azxy^4 + bx^2y^7$	$tz + g(z, x, y) \subset \mathbb{P}(19, 16, 7, 3)$ Diff=($\frac{1}{2}$, 0, 0, 0)	6
110	$z^2y + x^5 + ty^n$ $n = 4; 7; 9, 10$	$t^2z + z^2y + x + ty^n \subset \mathbb{P}(n + 1, 2n - 1, 4n + 1, 3)$ $\mathbb{P}(n + 1, 2n - 1, 3), \text{Diff}=(0, 0, \frac{4}{5}, 0)$	5,10,15
111	$z^2y + x^5 + txy^n$ $n = 3, 4; 5; 6, 7$	$t^2z + g(z, x, y) \subset \mathbb{P}(5n + 4, 10n - 3, 4n + 1, 11)$ Diff = $\emptyset$	6,7,11
112	$z^2y + x^5 + tx^2y^n$ $n = 3, 4$	$t^2z + g(z, x, y) \subset \mathbb{P}(5n + 3, 10n - 1, 4n + 1, 7)$ Diff = $\emptyset$	7
113	$z^2y + x^6 + ty^2(az + by^3) + czy^4 +$ dy^7	$t^2z + g(z, x^{1/6}, y) \subset \mathbb{P}(2, 3, 7, 1)$ $\mathbb{P}(2, 3, 1), \text{Diff}=(0, 0, \frac{5}{6}, 0)$	6
114	$z^2y + x^6 + ax^3y^4 + by^8$	$tz + z^2y + x^2 + axy^4 + by^8 \subset \mathbb{P}(9, 7, 8, 2)$ Diff=($\frac{1}{2}$, 0, $\frac{2}{3}$, 0)	12
115	$z^2y + x^6 + xy^n$ $n = 6; 7$	$tz + g(z, x, y) \subset \mathbb{P}(6n + 5, 6n - 5, 2n, 10)$ Diff=($\frac{1}{2}$, 0, 0, 0)	8,20
116	$z^2y + x^6 + azxy^3 + bx^2y^5$	$tz + g(z, x, y) \subset \mathbb{P}(17, 13, 5, 4)$ Diff=($\frac{1}{2}$, 0, 0, 0)	8
117	$z^2y + x^6 + ty^6$	$t^2z + z^2y + x + ty^6 \subset \mathbb{P}(7, 11, 25, 3)$ $\mathbb{P}(7, 11, 3), \text{Diff}=(0, 0, \frac{5}{6}, 0)$	18
118	$z^2y + x^6 + txy^4$	$t^2z + g(z, x, y) \subset \mathbb{P}(29, 44, 17, 14)$ Diff = $\emptyset$	7
119	$z^2y + x^6 + tx^2y^3$	$t^2z + z^2y + x^3 + txy^3 \subset \mathbb{P}(11, 17, 13, 5)$ Diff=(0, 0, $\frac{1}{2}$, 0)	10
120	$z^2y + tx^4 + y^n$ $n = 6; 8; 10$	$t^2z + z^2y + tx + y^n \subset \mathbb{P}(n + 1, 2n - 2, 3n - 1, 4)$ Diff=(0, 0, $\frac{3}{4}$, 0)	8,12,16
121	$z^2y + tx^4 + ty^2(az + by^3) + czy^4 +$ $dx^4y^2 + ey^7, a + b + c + e \neq 0$	$t^2z + g(z, x^{1/4}, y) \subset \mathbb{P}(2, 3, 5, 1)$ Diff=(0, 0, $\frac{3}{4}$, 0)	4

122	$z^2y + tx^4 + azy^5 + by^9$	$t^2z + z^2y + tx + azy^5 + by^9 \subset \mathbb{P}(5, 8, 13, 2)$ Diff= $(0, 0, \frac{3}{4}, 0)$	8
123	$z^2y + tx^4 + xy^n$ $n = 5; 6, 7; 8$	$t^2z + g(z, x, y) \subset \mathbb{P}(4n + 3, 8n - 7, 3n - 1, 13)$ Diff = $\emptyset$	7, 10, 13
124	$z^2y + tx^4 + x^2y^n$ $n = 4; 6$	$t^2z + g(z, x^{1/2}, y) \subset \mathbb{P}(2n + 1, 4n - 3, 3n - 1, 5)$ Diff= $(0, 0, \frac{1}{2}, 0)$	6, 10
125	$z^2y + tx^4 + azxy^3 + bx^2y^5$	$t^2z + g(z, x, y) \subset \mathbb{P}(11, 17, 7, 5)$ Diff = $\emptyset$	5
126	$z^2y + tx^4 + x^3y^4$	$t^2z + g(z, x, y) \subset \mathbb{P}(17, 27, 11, 7)$ Diff = $\emptyset$	7

(1) The singularity is $t^2z + z^i x^j f_{4-i-j}(x, y^2) + az^k x^l y^3 f_{2-k-l}(x, y) + by^6$. The exceptionality condition is

$$(i = 0 \parallel k = 0) \& \left((b \neq 0 \& (j, l \leq 2)) \parallel (b = 0 \& (j, l \leq 1)) \right)$$

(2) There are three cases. A). The singularity is $t^2z + z^2x + azx^i y^{j+1}(x + y)^k f_{2-i-j-k}(x, y) + x^{i_1} y(x + y)^{k_1} f_{4-i_1-k_1}(x, y)$. The exceptionality condition is $(i = 0 \parallel i_1 = 0) \& (k = 0 \parallel k_1 \leq 1)$.

B). The singularity is $t^2z + z^2x + zx^3 + azx^i y^{j+1}(x + y)^k f_{2-i-j-k}(x, y) + bx^{i_1} y^{j_1+1}(x + y)^{k_1} f_{4-i_1-j_1-k_1}(x, y)$. The exceptionality condition is $(i = 0 \parallel i_1 = 0) \& (j_1 \leq 2 \& k_1 \leq 3 \& i_1 \leq 3)$.

C). The singularity is $t^2z + z^2x + zx^3 + ax^5 + bxz^i y^{j+1}(x + y)^k f_{2-i-j-k}(x, y) + cx^{i_1} y^{j_1+1}(x + y)^{k_1} f_{4-i_1-j_1-k_1}(x, y)$, where $a \neq 0$. The exceptionality condition is $(i = 0 \parallel i_1 = 0) \& i_1 \leq 3$.

(3). The following notation $\frac{z^2x + zx^3 + x^5}{x(z + x^2)^2}$ means one of the following polynomials: $z^2x + zx^3$; $z^2x + zx^3 + ax^5$, where $a \neq \frac{1}{4}$; $x(z + x^2)^2$. The symbol (*) means that the third case is impossible.

16. Singularity $-t^2x + g(t, z, x, y)$

1	$z^4 + x^3y + y^n$ $n = 5; 9$	$t^2x + g(z^{1/2}, x, y) \subset \mathbb{P}(2n + 1, 3n, 2n - 2, 6)$ Diff= $(0, \frac{1}{2}, 0, 0)$	3, 12
2	$z^4 + x^3y + az^2y^3 + by^6$	$t^2x + g(z, x, y) \subset \mathbb{P}(13, 9, 10, 6)$ Diff = $\emptyset$	3
3	$z^4 + x^i y f_{3-i}(x, y^2) + atz^2y$	$t^2x + g(z^{1/2}, x, y) \subset \mathbb{P}(5, 7, 4, 2)$ Diff= $(0, \frac{1}{2}, 0, 0)$	4
4	$z^i f_{4-i}(z, y^2) + x^3y$ $i \leq 2$	$tx + g(z, x, y) \subset \mathbb{P}(17, 6, 7, 3)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	6
5	$z^4 + x^3y + zy^n$ $n = 4; 5; 7$	$tx + g(z, x, y) \subset \mathbb{P}(8n + 3, 3n, 4n - 3, 9)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	5, 6, 18
6	$z^4 + x^3y + zxy^4$	$tx + g(z, x, y) \subset \mathbb{P}(33, 11, 13, 5)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	10
7	$z^4 + x^3y + ty^n$ $n = 3; 4; 5; 6$	$t^2x + z + x^3y + ty^n \subset \mathbb{P}(2n + 1, 6n + 1, 2n - 1, 4)$ $\mathbb{P}(2n + 1, 2n - 1, 4)$, Diff= $(0, \frac{3}{4}, 0, 0)$	4, 5, 9 16
8	$z^4 + x^3y + tzy^n$ $n = 2, 3$	$t^2x + g(z, x, y) \subset \mathbb{P}(8n + 3, 6n + 1, 8n - 2, 10)$ Diff = $\emptyset$	5
9	$z^4 + x^i y^j f_{5-i-j}(x, y) + aty^3$ $(a \neq 0 \parallel i = 0) \& j \leq 2$	$t^2x + g(z^{1/4}, x, y) \subset \mathbb{P}(2, 5, 1, 1)$ $\mathbb{P}(2, 1, 1)$, Diff= $(0, \frac{3}{4}, 0, 0)$	4
10	$z^4 + az^2y^3 + by^6 + x^5$	$t^2x + z^2 + azy + by^2 + x^5 \subset \mathbb{P}(4, 5, 2, 5)$	10

		$\text{Diff}=(0, \frac{1}{2}, 0, \frac{2}{3})$	
11	$z^4 + x^5 + zy^4$	$t^2x + z^4 + x^5 + zy \subset \mathbb{P}(8, 5, 4, 15)$ $\text{Diff}=(0, 0, 0, \frac{3}{4})$	5
12	$z^4 + ax^5 + tzy^2 + \underline{c}zx^2y^2$	$t^2x + g(z, x, y^{1/2}) \subset \mathbb{P}(8, 5, 4, 7)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	7
13	$z^4 + az^2y^3 + by^6 + \underline{d}x^4y + \underline{c}zx^2y^2$ $ a + b \neq 0$	$tx + g(z, x, y) \subset \mathbb{P}(19, 6, 5, 4)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8
14	$z^4 + x^4y + ty^4$	$t^2x + z + x^4y + ty^4 \subset \mathbb{P}(13, 33, 7, 5)$ $\mathbb{P}(13, 7, 5), \text{Diff}=(0, \frac{3}{4}, 0, 0)$	20
15	$z^4 + x^3y^2 + az^2y^3 + by^6$	$t^2x + g(z^{1/2}, x, y) \subset \mathbb{P}(7, 9, 4, 3)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	6
16	$z^4 + x^3y^2 + y^7$	$t^2x + z + x^3y^2 + y^7 \subset \mathbb{P}(8, 21, 5, 3)$ $\mathbb{P}(8, 5, 3), \text{Diff}=(0, \frac{3}{4}, 0, 0)$	12
17	$z^4 + x^3y^2 + aty^4 + bxy^5$	$t^2x + g(z^{1/2}, x, y) \subset \mathbb{P}(10, 13, 6, 4)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	8
18	$z^4 + x^3y^2 + zy^n$ $n = 4; 5$	$t^2x + g(z, x, y) \subset \mathbb{P}(4n + 3, 3n, 4n - 6, 9)$ $\text{Diff} = \emptyset$	5,9
19	$z^4 + x^3y^2 + tzy^2 + bzy^3$	$t^2x + g(z, x, y) \subset \mathbb{P}(11, 7, 6, 5)$ $\text{Diff} = \emptyset$	5
20	$z^4 + zx^2y + y^n$ $n = 5; 7; 9; 11$	$tx + g(z, x, y) \subset \mathbb{P}(5n + 4, 2n, 3n - 4, 8)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,10,12 16
21	$z^4 + zx^2y + az^2y^n + by^{2n}$ $n = 3 \text{ } b \neq 0; 5$	$tx + g(z, x, y) \subset \mathbb{P}(5n + 2, 2n, 3n - 2, 4)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4,8
22	$z^i f_{4-i}(z, y^2) + zx^2y + ax^2y^3$ $(a = 0 \& i \leq 2) (a \neq 0 \& i \leq 3)$	$tx + g(z, x, y) \subset \mathbb{P}(11, 4, 5, 2)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
23	$z^4 + zx^2y + ty^n$ $n = 3, 4; 5; 6, 7$	$t^2x + g(z, x, y) \subset \mathbb{P}(5n + 4, 4n + 1, 6n - 4, 11)$ $\text{Diff} = \emptyset$	5,7,11
24	$z^4 + zx^2y + tzy^n$ $n = 3, 4$	$t^2x + g(z, x, y) \subset \mathbb{P}(5n + 3, 4n + 1, 6n - 2, 7)$ $\text{Diff} = \emptyset$	7
25	$z^4 + az^2y^3 + by^6 + zx^4$	$tx + z^4 + az^2y + by^2 + zx^4 \subset \mathbb{P}(13, 4, 3, 8)$ $\text{Diff}=(\frac{1}{2}, 0, 0, \frac{2}{3})$	16
26	$z^4 + zx^4 + tzy^2$	$t^2x + z^4 + zx^4 + tzy \subset \mathbb{P}(13, 8, 6, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
27	$z^4 + az^2y^3 + by^6 + zx^3y$	$tx + g(z, x, y) \subset \mathbb{P}(29, 9, 7, 6)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	12
28	$z^4 + zx^2y^2 + y^7$	$tx + g(z, x, y) \subset \mathbb{P}(43, 14, 13, 8)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	16
29	$z^4 + zx^2y^2 + ty^4$	$t^2x + z^4 + zx^2y + ty^2 \subset \mathbb{P}(14, 9, 8, 11)$ $\text{Diff}=(0, 0, 0, \frac{1}{2})$	11
30	$z^3y + x^5 + y^n$ $n = 6; 8$	$t^2x + g(z^{2/3}, x, y) \subset \mathbb{P}(2n, 5n - 5, n, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9,12
31	$z^3y + x^5 + ty^4 + bx^2y^4$	$t^2x + g(z^{2/3}, x, y) \subset \mathbb{P}(8, 17, 4, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	6
32	$z^3y + z^2(x^2 + y^3) + zx^2y^2 +$ (1) $zy^5 + x^4y + x^2y^4 + y^7$	$tx + g(z, x, y) \subset \mathbb{P}(11, 4, 3, 2)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4
34	$z^i y f_{3-i}(z, y^2) + x^6$	$tx + g(z, x, y) \subset \mathbb{P}(35, 12, 7, 6)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	12
35	$z^3y + x^6 + x^2y^5$	$tx + zy + x^6 + x^2y^5 \subset \mathbb{P}(25, 26, 5, 4)$	24

		$\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	
36	$z^3y + x^6 + zxy^4$	$tx + g(z, x, y) \subset \mathbb{P}(55, 19, 11, 9)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	18
37	$z^3y + x^6 + ty^4$	$t^2x + zy + x^6 + ty^4 \subset \mathbb{P}(20, 41, 8, 7)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
38	$z^3y + x^6 + tzy^2$	$t^2x + g(z, x, y) \subset \mathbb{P}(25, 17, 10, 9)$ $\text{Diff} = \emptyset$	9
39	$z^i y f_{3-i}(z, y^2) + \underline{a}x^4y^2 + \underline{b}zx^4$ $i \leq 2$	$tx + g(z, x, y) \subset \mathbb{P}(23, 8, 5, 4)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8
40	$z^3y + x^{2i}y^2f_{2-i}(x^2, y^3)$ $i \leq 1$	$tx + g(z^{1/3}, x, y) \subset \mathbb{P}(13, 14, 3, 2)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	12
41	$z^3y + x^4y^2 + y^9$	$tx + zy + x^4y^2 + y^9 \subset \mathbb{P}(29, 32, 7, 4)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	24
42	$z^3y + x^4y^2 + zxy^4$	$tx + g(z, x, y) \subset \mathbb{P}(31, 11, 7, 5)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10
43	$z^3y + x^4y^2 + ty^n$ $n = 4; 5$	$t^2x + g(z^{1/3}, x, y) \subset \mathbb{P}(3n + 2, 8n - 3, 2n - 2, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9,15
44	$z^3y + x^4y^2 + tzy^2$	$t^2x + g(z, x, y) \subset \mathbb{P}(19, 13, 8, 7)$ $\text{Diff} = \emptyset$	7
45	$z^3y + zx^4 + y^n$ $n = 6; 8$	$tx + g(z, x, y) \subset \mathbb{P}(10n - 1, 4n - 4, 2n + 1, 12)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	8,24
46	$z^3y + z^2x^2 + y^n$ $n = 5; 9$	$tx + g(z, x, y) \subset \mathbb{P}(5n - 2, 2n - 2, n + 2, 6)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,12
47	$z^3y + z^2x^2 + atzy^2 + bzy^3 +$ $cx^3y^2 + dy^6, d \neq 0 (ba \neq 0 ac \neq 0)$	$t^2x + g(z, x, y) \subset \mathbb{P}(7, 5, 4, 3)$ $\text{Diff} = \emptyset$	3
48	$z^3y + z^2x^2 + azxy^4 + bx^3y^3 +$ $cy^8, a + c \neq 0$	$tx + g(z, x, y) \subset \mathbb{P}(19, 7, 5, 3)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6
49	$z^3y + z^2x^2 + azyn^{n+1} + bx^2y^n$ $n = 3 \text{ } ab \neq 0; 5$	$tx + g(z, x, y) \subset \mathbb{P}(5n + 2, 2n, n + 2, 4)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4,8
50	$z^3y + z^2x^2 + ty^n$ $n = 4; 5$	$t^2x + g(z, x, y) \subset \mathbb{P}(5n - 2, 4n - 3, 2n + 2, 7)$ $\text{Diff} = \emptyset$	5,7

- (1) There are two cases. The common exceptionality condition is $|b| + |l| \neq 0$.
A). The singularity is $t^2x + z^3y + z^2x^2 + azx^2y^2 + bzy^5 + cx^4y + dx^2y^4 + ly^7$.
B). The singularity is $t^2x + z^3y + x^4y + az^2x^2 + czx^2y^2 + bzy^5 + dx^2y^4 + ly^7$.

17. Singularity $-t^2y + g(t, z, x, y)$

1	$z^4 + x^5 + az^2y^3 + by^6$	$t^2y + z^4 + x + az^2y^3 + by^6 \subset \mathbb{P}(5, 3, 12, 2)$ $\mathbb{P}(5, 3, 2), \text{Diff}=(0, 0, \frac{4}{5}, 0)$	6
2	$z^4 + x^5 + y^{2n+1}$ $n = 3; 4$	$t^2y + z + x + y^{2n+1} \subset \mathbb{P}(n, 2n + 1, 2n + 1, 1)$ $\mathbb{P}(n, 2n + 1, 1), \text{Diff}=(0, \frac{3}{4}, \frac{4}{5}, 0)$	16,20
3	$f_4(z, y^2) + x^5$	$ty + f_4(z, y^2) + x \subset \mathbb{P}(7, 2, 8, 1)$ $\mathbb{P}(7, 2, 1), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	10
4	$z^4 + x^5 + xy^n$ $n = 5; 7$	$t^2y + g(z^{1/2}, x, y) \subset \mathbb{P}(5n - 4, 5n, 2n, 8)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	12,16
5	$z^4 + x^{2i+1}f_{2-i}(x^2, y^3)$	$t^2y + g(z^{1/2}, x, y) \subset \mathbb{P}(13, 15, 6, 4)$ $\text{Diff}=(0, \frac{1}{2}, 0, 0)$	8
6	$z^4 + x^5 + az^2xy^2 + bx^2y^4 +$ $czy^5 + dzx^3y, a + b + c \neq 0$	$ty + g(z, x, y) \subset \mathbb{P}(17, 5, 4, 3)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6

7	$z^4 + x^5 + x^2y^5$	$t^2y + z + x^5 + x^2y^5 \subset \mathbb{P}(11, 25, 5, 3)$ $\mathbb{P}(11, 5, 3), \text{Diff}=(0, \frac{3}{4}, 0, 0)$	12
8	$z^4 + x^5 + zy^n$ $n = 4; 7$	$ty + z^4 + x + zy^n \subset \mathbb{P}(4n - 3, n, 4n, 3)$ $\mathbb{P}(4n - 3, n, 3), \text{Diff}=(\frac{1}{2}, 0, \frac{4}{5}, 0)$	10,30
9	$z^4 + x^5 + zxy^n$ $n = 3; 4; 5$	$ty + g(z, x, y) \subset \mathbb{P}(20n - 11, 5n, 4n, 11)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	6,18,22
10	$z^4 + x^5 + zx^2y^n$ $n = 2; 3$	$ty + g(z, x, y) \subset \mathbb{P}(20n - 7, 5n, 4n, 7)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,14
11	$z^4 + zx^4 + az^2y^3 + by^6$	$t^2y + z^4 + zx + az^2y^3 + by^6 \subset \mathbb{P}(5, 3, 9, 2)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	8
12	$z^4 + zx^4 + y^7$	$t^2y + z^4 + zx + y^7 \subset \mathbb{P}(12, 7, 21, 4)$ $\text{Diff}=(0, 0, \frac{3}{4}, 0)$	16
13	$z^4 + zx^4 + xy^n$ $n = 5; 6$	$ty + g(z, x, y) \subset \mathbb{P}(16n - 13, 4n, 3n, 13)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,26
14	$z^4 + zx^4 + x^3y^3$	$ty + g(z, x, y) \subset \mathbb{P}(41, 12, 9, 7)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
15	$z^3x + x^5 + azx^2y^n + by^{3n}$ $n = 2 \text{ } b \neq 0; 4$	$ty + g(z, x, y) \subset \mathbb{P}(15n - 5, 4n, 3n, 5)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	4,10
16	$z^3x + x^5 + y^{2n+1}$ $n = 3; 5$	$t^2y + g(z^{1/3}, x, y) \subset \mathbb{P}(5n, 8n + 4, 2n + 1, 5)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	7,15
17	$z^3x + x^5 + y^n$ $n = 8; 14$	$ty + g(z^{1/3}, x, y) \subset \mathbb{P}(5n - 5, 4n, n, 5)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	15,30
18	$z^3x + x^i f_{5-i}(x, y^2) + az^3y^2$ $a \neq 0 i \leq 3$	$ty + g(z^{1/3}, x, y) \subset \mathbb{P}(9, 8, 2, 1)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	6
19	$z^3x + x^5 + x^2y^n$ $n = 4; 8$	$ty + g(z^{1/3}, x, y) \subset \mathbb{P}(5n - 3, 4n, n, 3)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	10,18
20	$z^3x + x^5 + x^2y^5$	$t^2y + zx + x^5 + x^2y^5 \subset \mathbb{P}(11, 20, 5, 3)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
21	$z^3x + x^5 + zy^n$ $n = 4; 6; 8; 10$	$ty + g(z, x, y) \subset \mathbb{P}(15n - 11, 4n, 3n, 11)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	7,14,16 22
22	$z^3x + x^5 + zy^7$	$t^2y + g(z, x, y) \subset \mathbb{P}(47, 28, 21, 11)$ $\text{Diff} = \emptyset$	11
23	$z^3x + x^5 + z^2y^3$	$t^2y + g(z, x, y) \subset \mathbb{P}(19, 12, 9, 7)$ $\text{Diff} = \emptyset$	6
24	$z^3x + x^5 + z^2y^n$ $n = 4; 6$	$ty + g(z, x, y) \subset \mathbb{P}(15n - 7, 4n, 3n, 7)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	12,14
25	$z^3x + x^6 + y^7$	$t^2y + zx + x^6 + y^7 \subset \mathbb{P}(18, 35, 7, 6)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	9
26	$z^3x + x^6 + ax^3y^4 + by^8$	$ty + g(z^{1/3}, x, y) \subset \mathbb{P}(21, 20, 4, 3)$ $\text{Diff}=(\frac{1}{2}, \frac{2}{3}, 0, 0)$	18
27	$z^3x + x^6 + x^2y^5$	$t^2y + zx + x^6 + x^2y^5 \subset \mathbb{P}(13, 25, 5, 4)$ $\text{Diff}=(0, \frac{2}{3}, 0, 0)$	12
28	$z^3x + x^6 + zy^n$ $n = 5; 6$	$ty + g(z, x, y) \subset \mathbb{P}(18n - 13, 5n, 3n, 13)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	10,26
29	$z^3x + x^6 + zx^2y^3$	$ty + g(z, x, y) \subset \mathbb{P}(47, 15, 9, 7)$ $\text{Diff}=(\frac{1}{2}, 0, 0, 0)$	14
30	$z^3x + x^6 + z^2y^3$	$t^2y + g(z, x, y) \subset \mathbb{P}(23, 15, 9, 8)$ $\text{Diff} = \emptyset$	8

31	$z^3x + zx^4 + y^8$	$ty + g(z, x, y) \subset \mathbb{P}(77, 24, 16, 11)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	16
32	$z^3x + zx^4 + x^2y^n$ $n = 4; 6$	$ty + g(z, x, y) \subset \mathbb{P}(11n - 7, 3n, 2n, 7)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	8,14
33	$z^3x + zx^4 + x^3y^4$	$ty + g(z, x, y) \subset \mathbb{P}(39, 12, 8, 5)$ Diff= $(\frac{1}{2}, 0, 0, 0)$	10
34	$z^3x + tx^3 + y^n$ $n = 6; 10$	$t^2y + g(z^{1/3}, x, y) \subset \mathbb{P}(3n - 3, 5n - 1, n + 1, 6)$ Diff= $(0, \frac{2}{3}, 0, 0)$	9,18
35	$z^3x + tx^3 + ax^3y^3 + by^7$ $b \neq 0$	$t^2y + g(z^{1/3}, x, y) \subset \mathbb{P}(9, 17, 4, 3)$ Diff= $(0, \frac{2}{3}, 0, 0)$	6
36	$z^3x + tx^3 + ax^4y^2 + bx^2y^5 +$ $cy^8, b + c \neq 0$	$t^2y + g(z^{1/3}, x, y) \subset \mathbb{P}(7, 13, 3, 2)$ Diff= $(0, \frac{2}{3}, 0, 0)$	6
37	$z^3x + tx^3 + az^2y^3 + bx^3y +$ $cx^2y^4, a + c \neq 0$	$t^2y + g(z, x, y) \subset \mathbb{P}(11, 7, 5, 4)$ Diff = $\emptyset$	4
38	$z^3x + tx^3 + x^2y^6$	$t^2y + g(z^{1/3}, x, y) \subset \mathbb{P}(17, 31, 7, 4)$ Diff= $(0, \frac{2}{3}, 0, 0)$	12
39	$z^3x + tx^3 + zy^n$ $n = 4; 6$	$t^2y + g(z, x, y) \subset \mathbb{P}(9n - 7, 5n - 1, 3n + 2, 13)$ Diff = $\emptyset$	7,10
40	$z^3x + tx^3 + z^2y^4$	$t^2y + g(z, x, y) \subset \mathbb{P}(31, 19, 13, 8)$ Diff = $\emptyset$	8

REFERENCES

- [1] *Arnold V. I., Gusein-Zade S. M., Varchenko A. N.* Singularities of differentiable maps. Vol I, Birkhauser Boston, Boston, Mass., 1985.
- [2] *Iano-Fletcher A. R.* Working with weighted complete intersections, in “Explicit birational geometry of 3-folds”, 73–173, London Math. Soc. Lecture Note Ser., **281**, Cambridge Univ. Press, Cambridge, 2000.
- [3] *Fujino O.* The indices of log canonical singularities, Amer.J.Math. e-print math.AG/9909035
- [4] *Ishii S.* On isolated Gorenstein singularities, Math. Ann., **270**, (1985) 541–554.
- [5] *Ishii S.* The quotient of log-canonical singularities by finite groups, Advanced Studies in Pure Math. 29, (2000) 135–161
- [6] *Ishii S., Prokhorov Yu. G.* Hypersurface exceptional singularities, Intern. Journal of Math. 12 **6** (2001) 661–687.
- [7] *Kawamata Y.* Crepant blowin-ups of 3-dimensional canonical singularities and its application to degenerations of surfaces, Ann. Math 127(1988), 93–163.
- [8] *Kawamata Y., Matsuda K., Matsuki K.* Introduction to the minimal model program // Algebraic Geometry, Sendai. Adv. Stud. Pure Math. 1987. **10**. 283–360
- [9] *Kollar J. et al* Flips and abundance for algebraic threefolds, Astérisque 211,(1992).
- [10] *Kollár J.* Singularities of pairs, Proc. Symp. Pure Math. **62** (1995) 221–287
- [11] *Kudryavtsev S. A.* On exceptional terminal singularities, Math. Surveys, V. 56. No. 1 (2001), e-print math.AG/0111303.
- [12] *Kudryavtsev S. A.* On purely log terminal blow-ups, Math. Notes **69** No. 6 (2001) e-print math.AG/0111278.
- [13] *Lazzeri F.* On the monodromy of isolated singularities, Astérisque 7,8, (1973), 269–276
- [14] *Markushevich D. G.* Canonical singularities of three-dimensional hypersurfaces, Math. USSR-Izv. 26 (1985), 315–345.
- [15] *Oda T.* Convex bodies and algebraic geometry. An introduction to the theory of toric varieties, Ergeb. Math. Grenzgeb. (3) 1-5, Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, (1988).
- [16] *Prokhorov Yu. G.* On the existence of complements of canonical divisor for Mori conic bundles, Russian Acad. Sci. Sb. Math. 188 No. 11 (1997), 99–120.
- [17] *Prokhorov Yu. G.* Mori conic bundles with a reduced log-terminal boundary, J.Math.Sci. 91,(1999), 1051–1059.
- [18] *Prokhorov Yu. G.* Blow-ups of canonical singularities, Algebra. Proc. Internat. Conf. on the Occasion of the 90th birthday of A. G. Kurosh, Moscow, Russia, May 25–30, 1998, Yu. Bahturin ed., Walter de Gruyter, Berlin (2000), 301–317
- [19] *Prokhorov Yu. G., Markushevich D. G.* Klein’s group defines an exceptional singularity of dimension 3, J.Math.Sci. 91,(1999), 1060–1067.
- [20] *Prokhorov Yu. G.* Lectures on complements on log surfaces, MSJ Memoirs 10, (2001).
- [21] *Reid M.* Canonical threefolds, Géométrie Algébrique Angers, A.Beaville ed., Sijthoff and Noordhoff (1980), 273–310.
- [22] *Reid M.* Young person’s guide to canonical singularities, Proc. Symp. in Pure Math. 46,(1987), 343–416.
- [23] *Shokurov V. V.* 3-fold log flips, Russian Acad. Sci. Izv. Math. **40** (1993) 93–202 & **43** (1994) 527–558

- [24] *Shokurov V. V.* Complements on surfaces, J. of Math. Sci. 102 No. 2 (2000), 3876-3932.
- [25] *Varchenko A. N.* Zeta-function of monodromy and Newton's diagram, Invent. Math. 37,(1976), 253-262.
- [26] *Yonemura T.* Hypersurface simple K3-singularities , Tohoku Math. J. **42**, (1990), 351-380.

DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS, MOSCOW STATE
LOMONOSOV UNIVERSITY, 117234 MOSCOW, RUSSIA
E-mail address: kudryav@mech.math.msu.su